

Two Extended New Approaches to Vacuum, Matter & Fields

Alex Kaivarainen

*University of Turku, Department of Physics,
Vesilinnantie 5, FIN-20014, Turku, Finland,
H2o@karelia.ru www.karelia.ru/~ alexk*

Bo Lehnert

*Alfven Laboratory,
Royal Institute of Technology,
Stockholm, S-10044, Sweden
bo.lehnert@alfvenlab.kth.se*

Abstract

There are unsolved problems in physics leading to difficulties with Maxwell's equations that are not removed by and not directly associated with quantum mechanics. Recently a number of extended and modified theories have therefore been put forward. For a manifold of these the concepts of the vacuum state have played a fundamental rôle. The two theories by the authors (AK and BL) are related to this state. The first theory consists of a revised formulation of electromagnetics with an electric vacuum charge density(BL), and the second is a unified theory of Bivacuum, matter and fields(AK).

In the first theory by BL a space-charge current density arises from the condition of Lorentz invariance, in addition to the conventional displacement current. This leads both to steady electromagnetic states and to new wave types. The steady states provide models for the leptons, having an internal structure associated with vortex motion around the axis of symmetry at the velocity of light. For the electron the radial forces are kept in balance such as to prevent it from “exploding” under the action of its net self-charge. The electron model has the deduced character of a point-charge-like state with a very small radius, in which the infinite self-energy problem is removed, and a more surveyable alternative to renormalization is implemented. The electron charge is deduced from the theory, and it deviates by only one percent from its experimental value. Two individual photon models are further investigated. Both consist of axisymmetric configurations having the capacity of a boson particle of limited spatial extent and nonzero angular momentum. One model leads to the geometry of needle-shaped radiation. These photon models behave simultaneously as particles and as waves.

In the second theory by AK a Bivacuum model is established as a consequence of a new interpretation and development of Dirac theory, pointing to equal probability of positive and negative energy. Bivacuum is considered as the infinite dynamic superfluid matrix of virtual dipoles, named Bivacuum fermions ($\mathbf{BVF}^\uparrow$)ⁱ and Bivacuum bosons ($\mathbf{BVB}^\pm$)ⁱ, formed by correlated torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) (donuts), as a collective excitations of subquantum particles and antiparticles of opposite energy, charge and magnetic moments, separated by an energy gap. The spatial and energetic parameters of the torus and antitorus of primordial Bivacuum, i.e. in the absence of matter and field influence, correspond to electrons, muons and tauons ($i = e, \mu, \tau$). The positive and negative Virtual Pressure Waves ($\mathbf{VPW}^\pm$) and Virtual Spin Waves ($\mathbf{VirSW}^{S=\pm 1/2}$) are

the result of emission and absorption of positive and negative Virtual Clouds ($\mathbf{VC}^\pm$), resulting from transitions of $\mathbf{V}^+$ and $\mathbf{V}^-$ between different states of excitation. The formation of real sub-elementary fermions and their fusion to stable triplets of elementary fermions, corresponding to the electron and proton rest mass and charge origination, become possible at a certain symmetry shift between the $\mathbf{V}^+ \Updownarrow \mathbf{V}^-$ parameters. This shift appears at the velocity of angular rotation of pairs of $[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow]$ around a common axis, being determined by the Golden Mean condition: $(\mathbf{v}/\mathbf{c})^2 = \phi = 0.618$. The photon in such an approach is a result of fusion (annihilation) of two triplets of particle and antiparticle: (*electron + positron*) or (*proton + antiproton*). It represents a sextet of sub-elementary fermions and antifermions with axial structural symmetry.

New expressions for total, potential and kinetic energies of de Broglie waves of elementary particles, extending the relativity formalism, were obtained. The dynamic mechanism of [corpuscle (C) $\rightleftharpoons$ wave (W)] duality is proposed. It involves the modulation of the internal (hidden) quantum beats frequency between the asymmetric 'actual' (torus) and 'complementary' (antitorus) states of sub-elementary fermions or antifermions by the external - empirical de Broglie wave frequency of the whole particles. It is demonstrated, that the different kind of Bivacuum matrix excitations, accompanied [*Corpuscle* $\rightleftharpoons$ *Wave*] pulsations of sub-elementary fermions in triplets of fermions and their fact rotation are responsible for electric, magnetic and gravitational fields origination. The zero-point vibrations of particle and evaluated zero-point velocity of these vibrations are also a result of [*recoil* $\rightleftharpoons$ *antirecoil*] effects, accompanied [$\mathbf{C} \rightleftharpoons \mathbf{W}$] pulsations. The electromagnetic field, is a result of Corpuscle - Wave pulsation of photon and its fast rotation with angle velocity (ω_{rot}), equal to pulsation frequency. The clockwise or anticlockwise direction of photon rotation as respect to direction of its propagation, corresponds to its two polarization.

The Maxwell displacement current and the additional currents, increasing the refraction index of Bivacuum, are also the consequences of the Bivacuum dipole ($\mathbf{BVF}^\updownarrow$ and $\mathbf{BVB}^\pm$) vibrations and transitions, induced by an interaction with matter and fields. The nonzero contributions of the rest mass of photons and neutrinos in this perturbed Bivacuum follow from an elevated refraction index.

It is demonstrated that the dimensionless 'pace of time' ($d\mathbf{t}/\mathbf{t} = -d\mathbf{T}_k/\mathbf{T}_k$) and time itself ($\mathbf{t}$) for each closed system are determined by the change of this system's kinetic energy. They are positive if the particle dynamics of the system is slowing down, and negative in the opposite case. The concept of Virtual Replica (VR) or virtual hologram of any material object is developed.

The **VR** or virtual hologram is a result of interference of Virtual Pressure Waves ($\mathbf{VPW}_q^+$ and $\mathbf{VPW}_q^-$) and Virtual Spin waves ($\mathbf{VirSW}_q^{S=\pm 1/2}$) of the Bivacuum, representing "*reference waves*" with "*object waves*": $\mathbf{VPW}_m^\pm$ and $\mathbf{VirSW}_m^{\pm 1/2}$, modulated by de Broglie waves of molecules.

A possible Mechanism of Quantum entanglement between remote elementary particles via Virtual Guides of spin, momentum and energy ($\mathbf{VirG}_{S,M,E}$) is proposed. The consequences of the Unified theory (UT) of Bivacuum, matter and fields are in good agreement with known experimental data. The theory of AK makes an attempt to explain and unify the electric, magnetic and gravitation fields in the framework of the proposed mechanism of corpuscle-wave pulsations of elementary fermions, accompanied by different kinds of their dynamic interaction with the Bivacuum superfluid matrix.

The theories of BL and AK reinforce each other, by having several points in common. There are the similarities with the Dirac theory, symmetry with respect to the concepts of matter and antimatter, the non-appearance of magnetic monopoles, a space-charge vacuum current in addition to the displacement current, models of the leptons having an internal vortex-like structure and force balance, photon models with a small but nonzero rest mass and a slightly reduced velocity of propagation, and particle-wave properties, explaining self-interference even of single electrons or photons.

PACS indexing codes: 03; 04; 11

CONTENTS

ABSTRACT

I. Introduction

II. Electromagnetics with a Vacuum Charge Density (by Bo Lehnert)

A. Basis of the Theory

A.1 The Extended Field Equations

A.2 Comparison with the Dirac Theory

A.3 The Quantization Procedure

A.4 Momentum and Energy

B. New Aspects and Fundamental Applications

B.1 Steady Phenomena

B.2 Time-dependent Phenomena

C. General Features of Steady Axisymmetric States

C.1 The Generating Function

C.2 Quantum Conditions

D. A Model of the Electron

D.1 Generating Function and Integrated Field Quantities

D.2 The Magnetic Flux

D.3 Quantum Conditions of the Electron

D.4 Comparison with the Conventional Renormalization Procedure

D.5 Variational Analysis of the Integrated Charge

D.6 Force Balance of the Electron Model

E. Axisymmetric Wave Modes and Their Photon Models

E.1 Elementary Normal Modes

E.2 Wave Packets

E.3 Spatially Integrated Field Quantities

E.4 Features of Present Photon Model

F. Summary of Obtained Main Results in Section II

III. Unified Theory of Bivacuum, Particles Duality, Fields and Time (by A.Kaivarainen)

A.1. Properties of Bivacuum dipoles - Bivacuum fermions and Bivacuum bosons

*A.2 The basic (carrying) Virtual Pressure Waves ($VPW^\pm$) and
Virtual spin waves ($VirSW^{\pm 1/2}$) of Bivacuum*

A.3 Virtual Bose condensation (VirBC), as a base of Bivacuum nonlocality

B. Basic postulates of Unified Theory and their consequences

*B.1 The relation between the external and internal parameters of Bivacuum
fermions & quantum roots of Golden mean. The rest mass and charge origi-
nation*

C. Fusion of triplets of elementary particles from sub-elementary fermions and
antifermions at Golden mean conditions

*C.1 Correlation between our new model of hadrons and conventional quark
model of
protons, neutrons and mesons*

D. Total, potential and kinetic energies of elementary de Broglie waves

E. The dynamic mechanism of corpuscle-wave duality

F. The nature of electrostatic, magnetic and gravitational potentials, based
on Unified theory

F.1 Electromagnetic dipole radiation, as a consequence of charge oscillation

*F.2 The different kind of Bivacuum dipoles perturbation, induced by dynam-
ics of elementary particles*

*F.3 The new approach to quantum gravity and antigravity. The equality of
gravitational and inertial mass of particles*

F.4. Possible nature of neutrino and antineutrino

F.5 The Bivacuum dipoles symmetry shift and linear and circular ordering,

as a background of electrostatic and magnetic fields origination

F.6 Interpretation of Maxwell displacement current, based on Bivacuum model

F.7 New kind of current in secondary Bivacuum, additional to displacement one,

increasing the refraction index of Bivacuum

F.8 The mechanism, increasing the refraction index of Bivacuum

G. The Principle of least action, as a consequence of Bivacuum basic Virtual Pressure Waves ($\text{VPW}_{q=1}^{\pm}$) resonance interaction with particles. Possible source of energy for overunity devices.

G.1 The new approach to problem of Time

H. Virtual Replica (VR) of matter in Bivacuum

H.1 Bivacuum perturbations, induced by the oscillation of the total energy of de Broglie waves,

accompanied their thermal vibrations and recoil $\rightleftharpoons$ antirecoil effects

H.2 Modulation of basic Virtual Pressure Waves ($\text{VPW}_q^{\pm}$) and Virtual Spin Waves ($\text{VirSW}_q^{\pm 1/2}$) of Bivacuum by translational and librational dynamics (de Broglie waves) of the object molecules, correspondingly

I. Possible Mechanism of Quantum entanglement between remote elementary particles via Virtual Guides of spin, momentum and energy ($\text{VirG}_{\text{S,M,E}}$)

I.1 The role of tuning force ($\mathbf{F}_{\text{VPW}^{\pm}}$) of virtual pressure waves $\text{VPW}_q^{\pm}$ of Bivacuum in entanglement

J. Experimental data, confirming Unified theory (UT)

J.1 Radiation of accelerating charges, magnetic moment of the electron

J.2 The double turn (720^0), as a condition of the fermions spin state reversibility

J.3 Michelson-Morley experiment, as the evidence for the Virtual Replica of the Earth

J.4 Interaction of particles with their Virtual Replicas, as a background of two slit experiments explanation

K. The main Conclusions of Unified Theory (section III)

IV. Common Points and Main Features of the two Present Theories

V. General Conclusions

I. INTRODUCTION

Conventional field theory based on quantum mechanics and Maxwell's equations has been very successful in its applications to numerous problems in physics, and has sometimes manifested itself in an extremely good agreement with experimental results. Nevertheless there are certain areas within which these joint theories do not seem to provide fully adequate descriptions of physical reality. Thus, as already pointed out by Feynman[1], there are unsolved problems leading to difficulties with Maxwell's equations that are not removed by and not directly associated with quantum mechanics. Physics is therefore a far from completed field of research. Consequently, a number of extended and modified theories have recently been put forward, and for a manifold of these the concepts of the vacuum state have played a fundamental rôle. It would lead too far to present a detailed description of all these approaches within this review. The latter is thus limited to two theories by the authors (AK and BL) being related to the properties of the vacuum state. The first theory consists of a revised formulation of electromagnetics with a vacuum charge density(BL), and the second is a unified theory of Bivacuum, matter, fields and time (AK).

Among the present shortcomings of conventional theory, a number of examples can be given here. Turning first to charged leptons in the capacity of elementary particles such as the electron, there are the following problems:

- In conventional theory these particles are treated strictly as mathematical points, with no internal structure. This is unsatisfactory from the physical point of view.
- According to Maxwell's equations the electron cannot be kept in equilibrium by its own static forces. It tends to "explode" under the action of its self-charge, and has therefore to be kept in balance by some forces of nonelectromagnetic character, as stated by Jackson[2] among others. This explanation in terms of unspecified forces is incomplete.

- The infinite self-energy of the point charge is a serious problem. It has so far been solved in terms of a renormalization procedure, where the difference between two “infinities” gives a finite result. However, a more satisfactory procedure from the physical point of view has been called for by Ryder[3].
- There is so far no explanation by conventional theory why the free elementary electronic charge has a discrete minimum value, as found from experiments.

When further considering the photons and their wave phenomena, there are additional problems as follows:

- As pointed out by Heitler[4] among others, the photon in the form of a plane conventional electromagnetic wave does not possess any angular momentum (spin). For a nonzero spin, derivatives are therefore needed in the transverse direction of a photon configuration which has the capacity of a boson particle. In the case of conventional electromagnetic theory, however, such transverse derivatives lead to divergent solutions in space, as already found by Thomson[5] and being further discussed by several investigators. Such solutions become physically unacceptable. An even more serious complication is due to the fact that there does not arise any spin of an axisymmetric wave deduced from Maxwell’s equations for which there is a vanishing electric field divergence.
- There are no explicit solutions from conventional theory which become consistent with needle-like radiation. Such radiation has been suggested by Einstein, and it becomes associated with the photoelectric effect where a photon “knocks out” an atomic electron, as well as with the observed dot-shaped marks on a screen in two-slit experiments.
- The particle and wave nature of the photon remain in conventional theory as two separate concepts. In two-slit experiments the individual photon behaves on the other hand in terms of both concepts at the same time, as demonstrated by Tsuchiya et. al.[6].

These problems have been tackled in terms of an extended Lorentz invariant theory described in Section II of this review and based on the concept of a nonzero electric charge

density and electric field divergence in the vacuum (Lehnert [7–15]). This gives rise to a space-charge current density in the vacuum state, in addition to the displacement current, and it leads both to steady electromagnetic states and to new types of wave phenomena.

The steady states provide models of the leptons, such as the electron and the neutrino. These states possess an internal structure which comes out from the theory, thereby being associated with a vortex motion around the axis of symmetry at the velocity of light, and with related electric and magnetic field components. This vortex-like structure is not hypothetical, but it comes out from the solutions of the extended basic equations. There is an “electromagnetic confinement” by which the radial forces are kept in balance, also in the case of the electron which is then prevented from “exploding” under the action of its net self-charge. The electron model has further the deduced character of a point-charge-like state with a very small radius, in which the infinite self-energy problem is removed, and a physically more surveyable alternative to the renormalization process is being implemented. The deduced elementary electronic charge deviates by only one percent from its experimental value, and there are proposed mechanisms for the possible removal of this small deviation.

An application of the theory to photon physics further leads to several options of photon models. One resulting axisymmetric configuration has the capacity of a boson particle of limited spatial extent and a nonzero angular momentum. Another option leads to the geometry of a strongly limited transverse dimension, in the form of needle-shaped radiation. Thereby the elaborated photon models behave simultaneously as particles and as waves.

The Dirac equation points to equal probability of positive and negative energy[16]. In the Dirac vacuum its realm of negative energy is saturated by an infinite number of electrons. However, it was assumed that these electrons, following the Pauli principle, have not any gravitational or viscosity effects. Positrons in the Dirac model represent the “holes”, originated as a result of the electron jumps in the realm of positive energy, over the energetic gap $\Delta = 2m_0c^2$. Currently it becomes clear that the Dirac model of the vacuum is not general enough to explain all known experimental data, e.g. the boson emergency. The model of the Bivacuum presented in Section III of this review appears to be more advanced. However, it uses the same starting point of equal probability of positive and negative energy, being confined in each of the Bivacuum dipoles. These virtual dipoles are correlated pairs of torus and antitorus, representing, in turn, the collective vortical excitations of subquantum particles and antiparticles and separated by an energy gap.

The theory of AK explains the absence of magnetic monopoles, through the absence of a symmetry shift between the internal opposite actual quantities $|\mu_+|$ and the complementary $|\mu_-|$ magnetic moments of the virtual torus and antitorus which form Bivacuum fermions. This is in contrast to the mass and charge symmetry shifts, providing the rest mass and charge origination.

In a book by Bohm and Hiley[17] the electron is considered as a particle with well-defined position and momentum which are, however, under the influence of a special wave (quantum potential). In accordance with these authors an elementary particle is a *sequence of incoming and outgoing waves*, which are very close to each other. But the particle itself does not have a wave nature. The interference pattern in double-slit experiments is after Bohm the result of a periodically “bunched” character of a quantum potential. There is a basic difference in this respect with a model of duality in Section III, where it is assumed that the wave and corpuscle phases are realized alternatively at a high Compton frequency. This frequency is modulated by the empirical de Broglie wave frequency of particles, thereby making possible their interference with their own ‘virtual replica’, formed by Bivacuum Virtual Pressure Waves (VPW $^\pm$). In 1950 Wheeler and Misner[18] published “Geometrodynamics”, a new description of space-time properties based on topology. They supposed that elementary particles and antiparticles, their spins, and positive and negative charges can be presented as interconnected black and white holes. The connecting tube exists in another space-time than the holes themselves and the process of energy transmission looks as instantaneous. In conventional space-time the two ends of the tube, termed “wormholes”, can be a vast distance apart. This is one of the possible explanations of quantum non-locality. The “wormhole” idea has common features with our model of entanglement between coherent remote elementary particles via Virtual Guides of spin, momentum, and energy (VirG_{SME}) - microtubules, formed by pairs of torus and antitorus.

Sidhart[19, 20] considered the elementary particle as a *relativistic vortex* of Compton radius, from which he recovered its mass and quantized spin. A particle was pictured as a fluid vortex, steadily circulating at the light velocity along a 2D ring or spherical 3D shell with radius $L = \hbar/2mc$. Inside such a vortex the notions of negative energy, superluminal velocities and nonlocality are acceptable without contradiction with conventional theory. This view has some common features with the earlier models of extended electromagnetic steady states [7–15] described in Section II, and with the models of elementary and sub-

elementary particles and antiparticles[21–23] described in Section III. If the measurements are averaged over time $t \sim mc^2/\hbar$ and over space $L \sim \hbar/mc$, the imaginary part of the particle position disappears, and we are back in usual physics[19].

Aspden[24] introduced in his aether theory a basic unit, named Quon, as a pair of virtual muons of opposite charges, i.e. a muon and an antimuon. This idea is close to the model of Bivacuum dipoles consisting of a torus and an antitorus of opposite energy/mass, charge, and magnetic moment with Compton radii of the electron, muon and tauon[21–23]. In the book entitled “The physics of creation”, Aspden used the Thomson model of the electron, as a sphere with the radius a , charge e , and energy $2e^2/3a = mc^2$. This model strongly differs from that presented in the unified theory of Section III.

Barut and Bracken[25] considered the *zitterbewegung*, a rapidly oscillating imaginary part of particle position leading from the Dirac theory[16], as a harmonic oscillator in the Compton wavelength region of the particle. Also Einstein[26–28] and Schrödinger[29] considered the oscillation of the electron at the frequency $\nu = m_0c^2/\hbar$ and the amplitude $\zeta_{\max} = \hbar/(2m_0c)$. It was demonstrated by Schrödinger that the position of the free electron can be presented as $x = \bar{x} + \zeta$ where $\bar{x}$ characterizes the average position of the free electron, and ζ is its instantaneous position related to the oscillations. Hestness[30] further proposed that the *zitterbewegung* arises from self-interaction, resulting from wave-particle duality. These ideas are close to our explanation of zero-point oscillations of elementary particles, as a consequence of their recoil and antirecoil vibrations, induced by the particles corpuscle-wave pulsations.

Puthoff[31] developed the idea of “vacuum engineering”, using the hypothesis of a polarizable vacuum. The electric permittivity ε_0 and the magnetic permeability μ_0 are interrelated in symmetric vacuum, as $\varepsilon_0\mu_0 = 1/c^2$. It is conceived that a change of the vacuum refraction index $n = c/v = \varepsilon^{1/2}$ in gravitational or electric fields is accompanied by a variation of a number of space-time parameters. Fock[32] earlier explained the bending of a light beam, induced by gravitation near massive bodies, also by a change of the refraction in the vacuum, i.e. in another way than that of the general theory of relativity. However, the mechanism of this vacuum polarization and its related refraction index remains obscure.

The transformation of a neutron after scattering on a neutrino to a proton and electron in accordance with the electro-weak (EW) theory developed by Glashow[33], Weinberg[34], and Salam[35], is mediated by a negative W^- particle. The reverse reaction in EW theory

is mediated by a positive massless W^+ boson. Scattering of the electron on a neutrino, not being accompanied by charge transfer, is mediated by a third massless neutral boson Z^0 . For the explanation in EW theory of spontaneous symmetry violation of intermediate vector bosons, such as charged $W^\pm$ and neutral Z^0 with spin 1, the Higgs field was introduced and accompanied by a big mass in the origination of these exchange particles. The EW theory also needs the quantum of the Higgs field, namely Higgs bosons with big mass, zero charge and integer spin. The experimental discovery of the heavy W^+ and Z^0 particles in 1983 was considered as a confirmation of the EW theory. However, the Higgs field and Higgs bosons are still not found. The Unified theory of Section III proposes another explanation of spin, mass and charge origination, related to the symmetry shift of Bivacuum dipoles.

Thomson, Heaviside and Searl supposed that mass is an electrical phenomenon [24]. In the theory of Rueda and Haisch [36] it was proposed that the inertia is a reaction force, originating in a course of dynamic interaction between the electromagnetic zero-point field (ZPF) of the vacuum and the charges of elementary particles. However, it is not clear from their theory how the charges themselves are being created. In EW theory, mentioned above, two parameters of $W^\pm$ particles, such as charge and mass, are considered to be independent. The theory of AK tries to unify these parameters with the Bivacuum dipole symmetry shift, induced by rotation of their pairs around a common axis, fusion to triplets and external translational motion of triplets [21–23, 37]. This theory also makes an attempt to explain and unify the electric, magnetic and gravitation fields in the framework of the proposed mechanism of correlated corpuscle - wave pulsations of sub-elementary fermions in triplets, accompanied by different kinds of Bivacuum superfluid matrix perturbation. The internal time (t) and its pace (dt/t) of the each closed system of particles are parameters, characterizing both - the average velocity and acceleration of these particles. Such new approach to time problem differs from the conventional one, following from relativistic theory, however, they are compatible.

The conventional quantum field theory do not explain a lot of fundamental phenomena, like corpuscle-wave duality, origination of mass, charge, etc. This approach face a serious problem of singularity and related energy divergence. Our Unified theory goes from Bivacuum dipoles to particles and then to fields and contains the following main stages:

1. The Bivacuum dipoles symmetry shift, induced by their external translational-rotational motion, turning them to sub-elementary fermions at Golden mean conditions

$(\mathbf{v}/\mathbf{c})^2 = 0.618$; **2.** Origination of the rest mass, charge and fusion of triplets of sub-elementary fermions, representing elementary fermions. This process, like the previous one can be considered, as a kind of self-organization in Bivacuum; **3.** The correlated **Corpuscle** $\rightleftharpoons$ **Wave** pulsation of sub-elementary particles, composing elementary particles; **4.** Different kind of Bivacuum medium perturbations, induced by **C** $\rightleftharpoons$ **W** pulsation of unpaired and paired sub-elementary fermions, responsible for excitation of the electric, magnetic and gravitational potentials.

II. ELECTROMAGNETICS WITH A VACUUM CHARGE DENSITY

The vacuum is not merely an empty space. Its zero-point energy is related to electromagnetic vacuum fluctuations. Moreover, the observed electron-positron pair formation from an energetic photon indicates that electric charges can be created out of an electrically neutral vacuum state. It can therefore be conceived that a current density related to an electric space charge density and a nonzero electric field divergence can arise in the vacuum, in addition to the displacement current by Maxwell. As a consequence, the theory of this Section consists of a modified form extended beyond Maxwell's equations in the vacuum state, and having the latter equations as a special case. This approach is thus based on two mutually independent hypotheses:

- A nonzero charge density and a related electric field divergence can exist in the vacuum state. This should in principle not become less conceivable than the nonzero curl of the magnetic field strength for a conventional electromagnetic wave in the vacuum. All these concepts can be regarded as intrinsic properties of the electromagnetic field.
- The resulting extended (revised) form of the field equations should remain Lorentz invariant. Physical experience supports such a statement, as long as there are no observations getting into conflict with it.

This section presents a condensed review of the theory, and for its details reference is made to earlier publications[7–15].

A. Basis of the Theory

The basic concepts of the theory are first presented here, with resulting field equations, main features, and the related quantization procedure.

1. The Extended Field Equations

In presence of a charge density and a related current density in the vacuum, the extended field equations are now written in the four-dimensional form of a Proca-Type equation

$$\square A_\mu \equiv \left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) A_\mu = \mu_0 J_\mu \quad (2.1)$$

$$\mu = 1, 2, 3, 4$$

given in SI units In this equation

$$A_\mu = (\mathbf{A}, i\phi/c) \quad (2.2)$$

where $\mathbf{A}$ and ϕ are the magnetic vector potential and the electrostatic potential in three-space, and

$$J_\mu = (\mathbf{j}, ic\bar{\rho}) \quad (2.3)$$

is the additional four-current density which includes the corresponding three-space current density $\mathbf{j}$ and the electric charge density $\bar{\rho}$.

Maxwell's equations in the vacuum are recovered as a special case in which the current density J_μ disappears.

The current density (2.3) appearing in the right-hand side of the Proca-type equation (2.1) is here required to transform as a four-vector. This implies that the square of J_μ should become invariant to a transition from one inertial frame K to another such frame K'. Then Eq.(2.3) yields

$$j^2 - c^2 \bar{\rho}^2 = j'^2 - c^2 \bar{\rho}'^2 = \text{const} \quad (2.4)$$

In addition, the corresponding three-space current density $\mathbf{j}$ is required to exist only when there is also an electric charge density $\bar{\rho}$ associated with the nonzero electric field divergence. This implies that the constant in Eq.(2.4) vanishes. Consequently the three-space current density becomes

$$\mathbf{j} = \bar{\rho} \mathbf{C} = \varepsilon_0 (\nabla \cdot \mathbf{E}) \mathbf{C} \quad \mathbf{C}^2 = c^2 \quad (2.5)$$

where $\mathbf{C}$ is a velocity vector having a modulus equal to the velocity c of light. The final form of the four-current density is then given by

$$J_\mu = (\mathbf{j}, ic\bar{\rho}) = \varepsilon_0(\nabla \cdot \mathbf{E})(\mathbf{C}, ic) \quad (2.6)$$

In analogy with the direction to be determined for the current density in conventional theory, the unit vector $\mathbf{C}/c$ will depend on the geometry of the particular configuration to be considered. This unit vector can in principle have components which vary from point to point in space, whereas the modulus of $\mathbf{C}$ remains constant according to the invariance condition (2.5). Both curl $\mathbf{C}$ and div $\mathbf{C}$ can also differ from zero, but in all cases treated here we restrict ourselves to

$$\nabla \cdot \mathbf{C} = 0 \quad (2.7)$$

and to a time-independent velocity vector $\mathbf{C}$.

In the three-dimensional representation the extended equations of the vacuum state now become

$$\nabla \times \mathbf{B}/\mu_0 = \varepsilon_0(\nabla \cdot \mathbf{E})\mathbf{C} + \varepsilon_0\partial\mathbf{E}/\partial t \quad (2.8)$$

$$\nabla \times \mathbf{E} = -\partial\mathbf{B}/\partial t \quad (2.9)$$

$$\mathbf{B} = \nabla \times \mathbf{A} \quad \nabla \cdot \mathbf{B} = 0 \quad (2.10)$$

$$\mathbf{E} = -\nabla\phi - \partial\mathbf{A}/\partial t \quad (2.11)$$

$$\nabla \cdot \mathbf{E} = \bar{\rho}/\varepsilon_0 \quad (2.12)$$

where the first term of the right-hand member of Eq.(2.8) and Eq.(2.12) are the new parts being introduced by the present theory. The nonzero electric field divergence introduces an asymmetry in the appearance of the electric and magnetic fields in these equations.

The equations (2.8)–(2.12) are gauge invariant [10, 15], but not those which explicitly include a particle mass [3].

The presence in Eqs.(2.12) and (2.8) of the dielectric constant ε_0 and the magnetic permeability μ_0 of the conventional vacuum state may require further explanation. Liquid and solid matter consist of atoms and molecules which often behave as electric and magnetic dipoles. Such matter can thus become electrically polarized and magnetized when external electric and magnetic fields are being imposed. Then the constants ε_0 and μ_0 have to be

replaced by modified values ε and μ , which take these polarization effects into account. One consequence of this is that a conventional plane electromagnetic wave would propagate in such matter at a velocity $(1/\varepsilon\mu)^{1/2}$ being smaller than the velocity $c = (1/\varepsilon_0\mu_0)^{1/2}$ in empty vacuum space. In the present extended theory on the vacuum state, however, no electrically polarized and magnetized atoms or molecules are present. There are only electromagnetic fluctuations due to the zero-point field. In other words, the vacuum state is here conceived as a background of empty space upon which are superimposed zero point electromagnetic wave-like fluctuations, as well as phenomena of a more regular character which could take the form of the wave phenomena and field configurations to be treated in the present theory. In this respect Eq.(2.12) becomes identical with that used in the conventional theory on hot plasmas which contain freely moving charged particles of both polarities, and where electric charge separation and a resulting electric field can arise, such as in the case of plasma oscillations.

The question may also be raised why only $\nabla \cdot \mathbf{E}$ and not $\nabla \cdot \mathbf{B}$ should be permitted to become nonzero in an extended approach. It has then to be noticed that the nonzero electric field divergence has its support in the observed pair formation and the vacuum fluctuations, whereas a nonzero magnetic field divergence is so far a possible but not proved theoretical supposition. With Dirac[38] we shall therefore leave the magnetic monopole concept as an open question, but not include it at the present stage in the theory.

As shown later, this theory will lead to a small but nonzero photon rest mass. This raises a further question about the gauge invariance. According to earlier investigations it was concluded that the gauge invariance does not require the photon rest mass to be zero[9, 39]. In fact, the field equations (2.8)-(2.9) can easily be seen to become gauge invariant. A new gauge with a magnetic vector potential $\mathbf{A}'$ and an electrostatic potential ϕ' is defined by

$$\mathbf{A}' = \mathbf{A} + \nabla\psi \quad \phi' = \phi - \partial\psi/\partial t \quad (2.13)$$

When being inserted into Eqs.(2.10) and (2.11), these gauge relations lead to field strengths $\mathbf{E}' = \mathbf{E}$ and $\mathbf{B}' = \mathbf{B}$ which thus become invariant. The field equations (2.8) and (2.9) only contain the field strengths and therefore become gauge invariant.

2. Comparison with the Dirac Theory

In the theory of the electron by Dirac[40] the relativistic wave function ψ has four components in spin space. The three-space charge and current densities become[41]

$$\bar{\rho} = e\bar{\psi}\psi \quad (2.14)$$

and

$$\mathbf{j} = ce(\bar{\psi}\boldsymbol{\alpha}_i\psi) \quad i = 1, 2, 3 \quad (2.15)$$

where $\boldsymbol{\alpha}_i$ are the Dirac matrices of the three spatial directions, and the expression for $\bar{\psi}$ is obtained from that of ψ by replacing the included unit vectors and functions by their complex conjugate values. These equations are then seen to be similar to the corresponding relation (2.5) for the current density in the present theory where $|\mathbf{C}| = c$. The angular momentum in the Dirac theory emerges from the spin matrices $\boldsymbol{\alpha}_i$, whereas the relation $\mathbf{C}^2 = c^2$ leads to the two spin directions of the same momentum when the present theory is applied to an axially symmetric configuration such as that in a model of the electron.

It should further be observed that the present theory does not have to include the concepts of a net electric charge and a particle rest mass from the beginning. Such concepts will first come out from spatial integration of the electric charge density and the electromagnetic energy density. When relating this approach to the Dirac theory, wave functions have thus to be considered which only represent states without an initially included net charge and rest mass. With u as an arbitrary function and U as a constant, this yields a charge density[42]

$$\bar{\rho} = 2e\bar{U}U\bar{u}u \quad (2.16)$$

and the corresponding current density components

$$j_z = \pm c\bar{\rho}; \quad j_x = 0 \quad \text{and} \quad j_y = 0 \quad (2.17)$$

where a bar over U and u indicates the complex conjugate value. Other analogous forms can be chosen where instead $j_y = \pm c\bar{\rho}$ or $j_x = \pm c\bar{\rho}$.

This result indicates that there is a connection between the present theory and that by Dirac. But the former theory sometimes applies to a larger class of phenomena, in the capacity of both "bound" and of "free" states. Thereby the elementary charge is a given quantity in Dirac's theory, whereas the total net charge of the present approach is deduced from the field equations in a steady axisymmetric state, as shown later.

3. The Quantization Procedure

The relevant quantum conditions to be imposed are those on the angular momentum, the magnetic moment, and on the magnetic flux of the steady or time-dependent configuration to be analyzed. The rigorous and most complete way to proceed is then to quantize the field equations already from the outset.

As a first step, however, a simplification will instead be made here, by first determining the general solutions of the basic field equations, and then imposing the quantum conditions afterwards. This is at least justified by the fact that the quantized electrodynamic equations become equivalent to the field equations (2.1) in which the potentials A_μ and currents J_μ are replaced by their expectation values, as shown by Heitler[4]. In this connection Gersten[43, 44] states that Maxwell's equations should be used as a guideline for proper interpretations of quantum theories. This should also apply to the inhomogeneous equations (2.1) being used here. The present theory may therefore not be too far from the truth, by representing the most probable states in a first approximation to a rigorous quantum-theoretical approach. Thus the theory could be designated as a type of extended quantum electrodynamics ("EQED").

4. Momentum and Energy

In analogy with conventional electromagnetic theory, the basic equations (2.8) and (2.9) can be subject to vector and scalar multiplication to result in new but still identical forms. This leads to the momentum equation

$$\nabla \cdot {}^2\mathbf{S} = \bar{\rho}(\mathbf{E} + \mathbf{C} \times \mathbf{B}) + \varepsilon_0 \frac{\partial}{\partial t}(\mathbf{E} \times \mathbf{B}) \quad (2.18)$$

where ${}^2\mathbf{S}$ is the electromagnetic stress tensor[45], and where

$$\mathbf{f} = \bar{\rho}(\mathbf{E} + \mathbf{C} \times \mathbf{B}) \quad (2.19)$$

is the total local volume force. The corresponding integral form is

$$\int {}^2\mathbf{S} \cdot \mathbf{n} \, dS = \mathbf{F}_e + \mathbf{F}_m + \frac{\partial}{\partial t} \int \mathbf{g} \, dV \quad (2.20)$$

Here dS and dV are surface and volume elements, $\mathbf{n}$ is the surface normal,

$$\begin{aligned}\mathbf{F}_e &= \int \bar{\rho} \mathbf{E} dV \\ \mathbf{F}_m &= \int \bar{\rho} \mathbf{C} \times \mathbf{B} dV\end{aligned}\tag{2.21}$$

are the integrated electric and magnetic volume forces, and

$$\mathbf{g} = \varepsilon_0 \mathbf{E} \times \mathbf{B} = \frac{1}{c^2} \mathbf{S}\tag{2.22}$$

can be interpreted as an electromagnetic momentum density, with $\mathbf{S}$ denoting the Poynting vector. The latter can be conceived to represent the magnitude and direction of the energy flow in space.

One further obtains the energy equation

$$-\nabla \cdot \mathbf{S} = -\left(\frac{1}{\mu_0}\right) \nabla \cdot (\mathbf{E} \times \mathbf{B}) = \bar{\rho} \mathbf{E} \cdot \mathbf{C} + \frac{1}{2} \varepsilon_0 \frac{\partial}{\partial t} (\mathbf{E}^2 + c^2 \mathbf{B}^2)\tag{2.23}$$

with its integral form

$$\int \mathbf{S} \cdot \mathbf{n} dS + \int \bar{\rho} \mathbf{E} \cdot \mathbf{C} dV = -\frac{1}{2} \varepsilon_0 \int \frac{\partial}{\partial t} (\mathbf{E}^2 + c^2 \mathbf{B}^2) dV\tag{2.24}$$

Using the basic equations (2.8)–(2.12) and some vector identities, expressions can be obtained for the energy densities, as given by

$$w_f = \frac{1}{2} (\varepsilon_0 \mathbf{E}^2 + \mathbf{B}^2 / \mu_0)\tag{2.25}$$

$$w_s = \frac{1}{2} (\bar{\rho} \phi + \mathbf{j} \cdot \mathbf{A}) = \frac{1}{2} \bar{\rho} (\phi + \mathbf{C} \cdot \mathbf{A})\tag{2.26}$$

Here w_f can be interpreted as a “field energy density” being associated with the field strengths $\mathbf{E}$ and $\mathbf{B}$, and w_s as a “source energy density” being associated with the sources $\bar{\rho}$ and $\mathbf{j}$ of the electromagnetic field. Conditions for the integral forms of w_f and w_s to become equal or to be different have been investigated, as shown in detail elsewhere[10].

B. New Aspects and Fundamental Applications

The introduction of an additional degree of freedom in the form of a nonzero electric field divergence, and the resulting transition from a homogeneous d’Alembert equation to an inhomogeneous Proca-type equation, lead to a number of new aspects and applications.

These include the new class of steady states, and an extended class of time-dependent states with new wave phenomena. Within these classes there are special applications to models of the electron, neutrino, and photon, as well as to string-shaped geometry, and to total reflection of dissipative plane waves. This condensed review is limited to models of the electron and photon.

The analysis which follows mainly concerns axially symmetric systems, but asymmetric configurations can in principle also be treated by the theory.

1. *Steady Phenomena*

In a time-independent case the space-charge current density makes possible the existence of steady electromagnetic states which are absent within the frame of Maxwell's equations. From relations (2.7)-(2.12) the basic steady-state equations

$$c^2 \nabla \times \nabla \times \mathbf{A} = -\mathbf{C}(\nabla^2 \phi) = (\bar{\rho}/\epsilon_0)\mathbf{C} \quad (2.27)$$

are then obtained. On the basis of the electromagnetic forces given by Eqs.(2.18) and (2.21), the corresponding steady electromagnetic equilibrium of the electron will later be discussed.

Axisymmetric states are of special interest and will be treated in detail in the following subsections. These states can roughly be pictured as a result of "self-confined" circulating electromagnetic radiation, on which relevant quantum conditions are being imposed. Among these "bound" states two subclasses are of special interest:

- Particle-shaped states where the geometrical configuration is bounded both in the axial and in the radial directions. There are states both with a nonzero as well as with a zero net electric charge to be investigated. The corresponding solutions and models may have some bearing on and contribute to the understanding of such truly elementary particles as the leptons.
- String shaped states where the geometrical configuration is uniform in the axial direction and becomes piled up near the axis of symmetry. These equilibria can in an analogous manner reproduce several desirable features of the earlier proposed hadron string model.

2. Time-dependent Phenomena

The basic equations for time-dependent states combine to

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \nabla^2\right)\mathbf{E} + (c^2 \nabla + \mathbf{C} \frac{\partial}{\partial t})(\nabla \cdot \mathbf{E}) = 0 \quad (2.28)$$

for the electric field. From the corresponding solution the magnetic field $\mathbf{B}$ can be obtained by means of Eq.(2.9). A divergence operation on Eq.(2.8) further yields

$$\left(\frac{\partial}{\partial t} + \mathbf{C} \cdot \nabla\right)(\nabla \cdot \mathbf{E}) = 0 \quad (2.29)$$

in combination with Eq.(2.7). In some cases this equation becomes useful to the analysis, but it does not introduce more information than that already contained in Eq.(2.28).

Three limiting cases can be identified on the basis of Eq.(2.28):

- When $\nabla \cdot \mathbf{E} = 0$ and $\nabla \times \mathbf{E} \neq 0$ the result is a conventional *transverse* electromagnetic wave, henceforth denoted as an “EM wave”.
- When $\nabla \cdot \mathbf{E} \neq 0$ and $\nabla \times \mathbf{E} = 0$ a purely *longitudinal* electric space-charge wave arises, here being denoted as an “S wave”.
- When both $\nabla \cdot \mathbf{E} \neq 0$ and $\nabla \times \mathbf{E} \neq 0$ a hybrid *nontransverse* electromagnetic space-charge wave appears, here denoted as an “EMS wave”.

In a general case these various modes can become superimposed, also the EMS modes with different velocity vectors $\mathbf{C}$. That the basic equations can give rise both to modes with a nonvanishing and a vanishing electric field divergence is not less conceivable than the analogous property of the conventional basic equations governing a magnetized plasma. The frame of the latter accommodates both longitudinal electrostatic waves and transverse Alfvén waves. These are, like the EM, S, and EMS waves, separate modes even if they originate from the same basic formalism.

The applications which follow are related to the EMS waves. The conditions under which the S wave may exist are not clear at this stage.

C. General Features of Steady Axisymmetric States

The analysis is now restricted to particle-shaped axisymmetric states. For these a frame (r, θ, φ) of spherical coordinates is introduced where all relevant quantities are independent

of the angle φ . The analysis is further restricted to a current density $\mathbf{j} = (0, 0, C\bar{\rho})$ and a magnetic vector potential $\mathbf{A} = (0, 0, A)$. Here $C = \pm c$ represents the two possible spin directions. The basic equations (27) then take the form

$$\frac{(r_0\rho)^2\bar{\rho}}{\varepsilon_0} = D\phi = [D + (\sin\theta)^{-2}](CA) \quad (2.30)$$

where the dimensionless radial coordinate

$$\rho = r/r_0 \quad (2.31)$$

has been introduced, r_0 is a characteristic radial dimension, and the operator D is given by

$$\begin{aligned} D &= D_\rho + D_\theta \\ D_\rho &= -\frac{\partial}{\partial\rho}(\rho^2\frac{\partial}{\partial\rho}) \\ D_\theta &= -\frac{\partial^2}{\partial\theta^2} - \frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\theta} \end{aligned} \quad (2.32)$$

1. The Generating Function

The general solution of Eqs.(2.30) in particle-shaped geometry can now be obtained in terms of a *generating function*

$$F(r, \theta) = CA - \phi = G_0.G(\rho, \theta) \quad (2.33)$$

where G_0 stands for a characteristic amplitude and G for a normalized dimensionless part. This yields the solution

$$CA = -(\sin\theta)^2 DF \quad (2.34)$$

$$\phi = -[1 + (\sin\theta)^2 D]F \quad (2.35)$$

$$\bar{\rho} = -(\frac{\varepsilon_0}{r_0^2\rho^2})D[1 + (\sin\theta)^2 D]F \quad (2.36)$$

which is easily seen by direct insertion to satisfy Eqs.(2.30). Starting from an arbitrary function F , the corresponding spatial distributions of the potentials CA and ϕ and of the space-charge density $\bar{\rho}$ are thus generated, i.e. those which simultaneously satisfy the set (2.30) of equations. If one would instead start from a given distribution of only one of the field quantities CA , ϕ , or $\bar{\rho}$, this would not provide a simple solution such as that obtained from the set (2.33)–(2.36).

The quantities (2.34)–(2.36) are uniquely determined by the generating function. The latter can be chosen such that all field quantities decrease rapidly to zero at large radial distances from the origin. In this way the charge and current density, as well as all associated features of an electromagnetic field configuration, can be confined to a limited region of space.

In the analysis of particle-shaped states the functions

$$f(\rho, \theta) = -(\sin \theta)D[1 + (\sin \theta)^2 D]G \quad (2.37)$$

$$g(\rho, \theta) = -[1 + 2(\sin \theta)^2 D]G \quad (2.38)$$

will now be introduced. Using expressions (2.34)–(2.36), (2.26), (2.37) and (2.38), integrated field quantities can be obtained which represent a net electric charge q_0 , magnetic moment M_0 , mass m_0 , and angular momentum s_0 . The magnetic moment is obtained from the local contributions provided by the current density (2.5). The mass and the angular momentum are deduced from the local contributions of w_s/c^2 being given by the source energy density (2.26) and the energy relation by Einstein. The current density (2.5) then behaves as a convection current being common to all contributions from the charge density. The corresponding mass flow originates from the velocity vector $\mathbf{C}$ which has the same direction for positive as for negative charge elements. The integrated field quantities finally become

$$q_0 = 2\pi\epsilon_0 r_0 G_0 J_q \quad I_q = f \quad (2.39)$$

$$M_0 = \pi\epsilon_0 C r_0^2 G_0 J_M \quad I_M = \rho(\sin \theta)f \quad (2.40)$$

$$m_0 = \pi(\epsilon_0/c^2)r_0 G_0^2 J_m \quad I_m = fg \quad (2.41)$$

$$s_0 = \pi(\epsilon_0 C/c^2)r_0^2 G_0^2 J_s \quad I_s = \rho(\sin \theta)fg \quad (2.42)$$

These relations include normalized integrals defined by

$$J_k = \int_{\rho_k}^{\infty} \int_0^{\pi} I_k \, d\rho d\theta \quad k = q, M, m, s \quad (2.43)$$

where ρ_k are small radii of circles centered around the origin $\rho = 0$ when G is divergent there, and $\rho_k = 0$ when G is convergent at $\rho = 0$. The case $\rho_k \neq 0$ will later be treated in detail. So far the integrals (2.39)–(2.43) are not uniquely determined but depend on the distribution of the function G in space. They will first become fully determined when further

conditions are being imposed, such as those of a quantization. The general forms (2.39)–(2.43) thus provide the integrated quantities with a certain degree of flexibility, being related to the extra degree of freedom which has been introduced through the nonzero electric field divergence.

At this point a further step is taken by imposing the restriction of a separable generating function

$$G(\rho, \theta) = R(\rho).T(\theta) \quad (2.44)$$

The integrands of the normalized form (2.43) then become

$$I_q = \tau_0 R + \tau_1 (D_\rho R) + \tau_2 D_\rho (D_\rho R) \quad (2.45)$$

$$I_M = \rho(\sin \theta) I_q \quad (2.46)$$

$$I_m = \tau_0 \tau_3 R^2 + (\tau_0 \tau_4 + \tau_1 \tau_3) R (D_\rho R) + \tau_1 \tau_4 (D_\rho R)^2 + \\ + \tau_2 \tau_3 R D_\rho (D_\rho R) + \tau_2 \tau_4 (D_\rho R) [D_\rho (D_\rho R)] \quad (2.47)$$

$$I_s = \rho(\sin \theta) I_m \quad (2.48)$$

where

$$\tau_0 = -(\sin \theta)(D_\theta T) - (\sin \theta) D_\theta [(\sin^2 \theta)(D_\theta T)] \quad (2.49)$$

$$\tau_1 = -(\sin \theta) T - (\sin \theta) D_\theta [(\sin^2 \theta) T] - \sin^3 \theta (D_\theta T) \quad (2.50)$$

$$\tau_2 = -(\sin^3 \theta) T \quad (2.51)$$

$$\tau_3 = -T - 2(\sin^2 \theta)(D_\theta T) \quad (2.52)$$

$$\tau_4 = -2(\sin^2 \theta) T \quad (2.53)$$

Among the possible forms to be adopted for the radial function $R(\rho)$ and the polar function $T(\theta)$, we will here consider the following cases:

- The radial part R can become convergent or divergent at the origin $\rho = 0$ but it always goes strongly towards zero at large distances from it.
- The polar part T is always finite and has finite derivatives. It can become symmetric or antisymmetric with respect to the “equatorial plane” (midplane) defined by $\theta = \pi/2$.

We first consider the radial part R and the corresponding normalized integrals J_q and J_M of the integrated charge and magnetic moment. After a number of partial integrations this yields the following conclusions:

- Convergent radial functions R lead to a class of electrically neutral particle states where q_0 and M_0 both vanish, whereas m_0 and s_0 are nonzero.
- For the present particle-shaped geometry to result in electrically charged states, the *divergence* of the radial function becomes a necessary but not sufficient condition. This leads to the subsequent question whether the corresponding integrals (2.43) would then be able to form the basis of a steady state having *finite* and nonzero values of all the integrated field quantities (2.39)–(2.42). This will later be shown to become possible.

We then turn to the polar part T of the generating function. The top-bottom symmetry properties of the polar function T are now considered with respect to the equatorial plane. The integrals (2.43) which are evaluated in the interval $0 \leq \theta \leq \pi$ become nonzero for symmetric integrands I_k , but vanish for antisymmetric ones. These symmetry properties are evident from expressions (2.45)–(2.53). Thus, the product of two symmetric or of two antisymmetric functions becomes symmetric, whereas a product of a symmetric function with an antisymmetric one becomes antisymmetric. The symmetry or antisymmetry of T leads to a corresponding symmetry or antisymmetry of $D_\theta T$, $D_\theta[(\sin \theta)^2 T]$, and $D_\theta[(\sin \theta)^2 (D_\theta T)]$. Therefore all functions (2.49)–(2.53) are either symmetric or antisymmetric in the same way as T . The symmetry rules with respect to T thus lead to two general conclusions:

- The charge and the magnetic moment vanish for antisymmetric forms of T , irrespective of the form of R .
- The mass and the angular momentum generally differ from zero, both when T is symmetric and when it is antisymmetric.

To sum up, it is thus found that a model of the electron can be based on R being divergent at the origin and T having top-bottom symmetry with respect to the equatorial plane, whereas a model of the neutrino can be based on any of the remaining alternatives being considered here.

Finally a few words should be said about matter and antimatter particle models. With $C = \pm c$ and $G_0 = \pm|G_0|$ it is seen from Eqs.(2.39)–(2.53) that there are pairs of solutions having the same geometry of their local distributions, but having opposite signs of the corresponding integrated field quantities in places where G_0 and C appear linearly. This could become the basis for models of charged and neutral particles of matter as well as of antimatter. A minor difference between matter and antimatter will then not be included in the models.

2. Quantum Conditions

The angular momentum condition to be imposed on the models of the electron in the capacity of a fermion particle, as well as of the neutrino, is combined with Eq.(2.42) to result in

$$s_0 = \pi(\varepsilon_0 C/c^2)r_0^2 G_0^2 J_s = \pm h/4\pi \quad (2.54)$$

This condition becomes compatible with the two signs of $C = \pm c$, as obtained from the Lorentz invariance.

In particular, for a charged particle such as the electron, muon, tauon or their antiparticles, Eqs.(2.39) and (2.42) combine to

$$\begin{aligned} q^* &\equiv |q_0/e| = (f_0 J_q^2 / 2 J_s)^{1/2} \\ f_0 &= 2\varepsilon_0 c h / e^2 \end{aligned} \quad (2.55)$$

where q^* is a dimensionless charge being normalized with respect to the experimentally determined electronic charge “ e ”, and $f_0 \approx 137.036$ is the inverted value of the fine-structure constant.

According to Dirac[40], Schwinger[46] and Feynman[47] the quantum condition on the magnetic moment M_0 of a charged particle such as the electron becomes

$$M_0 m_0 / q_0 s_0 = 1 + \delta_M \quad \delta_M = 1/2\pi f_0 \quad (2.56)$$

which shows excellent agreement with experiments. Here the unity term of the right-hand member is due to Dirac who obtained the correct Landé factor by considering the electron to be situated in an imposed external magnetic field. This leads to a magnetic moment being twice as large as that expected from an elementary proportionality relation between

the electron spin and the magnetic moment[3]. Further, in Eq.(2.56) the term δ_M is a small quantum mechanical correction due to Schwinger and Feynman.

Conditions (2.54) and (2.56) can also be made plausible by elementary physical arguments based on the present picture of a particle-shaped state of “self-confined” radiation. In the latter picture there is a circulation of radiation at the velocity of light around the axis of symmetry. If this circulation takes place at the average characteristic radial distance r_0 , the corresponding frequency of revolution would become $\nu \cong c/2\pi r_0$. In combination with the energy relations $W = m_0 c^2$ and $W = h\nu$ by Einstein and Planck, one then obtains

$$r_0 m_0 \cong \frac{h}{2\pi c} \quad (2.57)$$

where the radius r_0 has the same form as the Compton radius. We shall later return to this concept, at the end of Section D.1. The result (2.57) should, however, only be taken as a crude physical indication where the radius in Eq.(2.57) comes out to be far too large as compared to observations. With the total current $q_0\nu$, the magnetic moment further becomes

$$M_0 = \pi r_0^2 q_0 \nu = \frac{1}{2} q_0 c r_0 = \frac{q_0 h}{4\pi m_0} \quad (2.58)$$

which agrees with conditions (2.54) and (2.56) when δ_M is neglected. Thereby the angular momentum becomes $|s_0| = r_0 c m_0 / 2 = h/4\pi$ when its inwardly peaked spatial distribution corresponds to an equivalent radius $r_0/2$.

The unequal integrands of the magnetic moment and the angular momentum in Eqs.(2.40) and (2.42) could also be reconcilable with a proportionality relation between s_0 and M_0 which leads to the correct Landé factor. Another possibility of obtaining its correct value is to modify Eq.(2.57) to represent the first “subharmonic” given by a path length $4\pi r_0$ around the axis. In an axisymmetric case this would, however, become a rather far-fetched proposal. In any case, when using the correct value of this factor, the electronic charge deduced later in Section D turns out to be close to its experimental value. This would not be the case for a Landé factor being half as large.

Finally, in a charged particle-shaped state with a nonzero magnetic moment, the electric current distribution will generate a total magnetic flux Γ_{tot} and a corresponding total magnetic energy. In such a state the quantized value of the angular momentum further depends on the type of configuration being considered. It thus becomes $|s_0| = h/4\pi$ for a fermion,

but $|s_0| = h/2\pi$ for a boson. We now consider the electron to be a system also having a quantized charge q_0 . As in a number of other physical systems, the flux should then become quantized as well, and be given by the two quantized concepts s_0 and q_0 , in a relation having the dimension of magnetic flux. This leads to the quantum condition[10, 12]

$$\Gamma_{tot} = |s_0/q_0| \quad (2.59)$$

D. A Model of the Electron

A model of the electron will now be elaborated which in principle also applies to the muon, tauon and to corresponding antiparticles[48]. According to the previous analysis in Section C.1, a charged particle-shaped state only becomes possible by means of a generating function having a radial part R which is divergent at the origin, and a polar part T with top-bottom symmetry. In its turn, the divergence of R leads to the question how to obtain finite and nonzero integrated field quantities (2.39)–(2.43). The following analysis will show this to become possible, by shrinking the characteristic radius r_0 to the very small values of a “point-charge-like” state. It does on the other hand not imply that r_0 has to become strictly equal to zero, which would end up into the unphysical situation of a structureless point.

1. Generating Function and Integrated Field Quantities

The generating function to be considered has the parts

$$R = \rho^{-\gamma} e^{-\rho} \quad \gamma > 0 \quad (2.60)$$

$$\begin{aligned} T &= 1 + \sum_{v=1}^n \{a_{2v-1} \sin[(2v-1)\theta] + a_{2v} \cos(2v\theta)\} = \\ &= 1 + a_1 \sin \theta + a_2 \cos 2\theta + a_3 \sin 3\theta + \dots \end{aligned} \quad (2.61)$$

Concerning the radial part (2.60), it may at a first glance appear to be somewhat special and artificial. Under general conditions one could thus have introduced a negative power series of ρ instead of the single term $\rho^{-\gamma}$. However, for a limited number of terms in such a series,

that with the largest negative power will in any case dominate near the origin. Moreover, due to the following analysis the same series has to contain one term only, with a locked special value[12] of the radial parameter γ . The exponential factor in expression (2.60) has been included to secure the convergence of any moment with R at large distances from the origin. This factor will not appear in the end results of the analysis.

The polar part (2.61) represents a general form of axisymmetric geometry having top-bottom symmetry with respect to the equatorial plane. Here it should be noticed that all sine and cosine terms can be rewritten into a corresponding power series of $\sin \theta$, as well as the entire resulting series.

The radial form given by Eq.(2.60) can now be inserted into the integrands (2.45)–(2.48). Since the radial integrals (2.43) are extended to very low limits ρ_k , they become dominated by contributions from terms of the strongest negative power. Keeping these contributions only, the integrands reduce to

$$I_k = I_{k\rho} I_{k\theta} \quad k = q, M, m, s \quad (2.62)$$

where

$$I_{q\rho} = R \quad I_{M\rho} = \rho R \quad (2.63)$$

$$I_{m\rho} = R^2 \quad I_{s\rho} = \rho R^2$$

and

$$I_{q\theta} = \tau_0 - \gamma(\gamma - 1)\tau_1 + \gamma^2(\gamma - 1)^2\tau_2 \quad (2.64)$$

$$I_{M\theta} = (\sin \theta) I_{q\theta} \quad (2.65)$$

$$I_{m\theta} = \tau_0\tau_3 - \gamma(\gamma - 1)(\tau_0\tau_4 + \tau_1\tau_3) + \gamma^2(\gamma - 1)^2(\tau_1\tau_4 + \tau_2\tau_3) - \gamma^3(\gamma - 1)^3\tau_2\tau_4 \quad (2.66)$$

$$I_{s\theta} = (\sin \theta) I_{m\theta} \quad (2.67)$$

$I_{M\theta} = (\sin \theta) I_{q\theta}$ The corresponding integrals (2.43) then become

$$J_k = J_{k\rho} J_{k\theta} \quad (2.68)$$

where

$$J_{q\rho} = \frac{1}{\gamma - 1} \rho_q^{-(\gamma-1)} \quad (2.69)$$

$$J_{M\rho} = \frac{1}{\gamma - 2} \rho_M^{-(\gamma-2)}$$

$$J_{m\rho} = \frac{1}{2\gamma - 1} \rho_m^{-(2\gamma-1)} \quad (2.70)$$

$$J_{s\rho} = \frac{1}{2(\gamma - 1)} \rho_s^{-2(\gamma-1)}$$

and

$$J_{k\theta} = \int_0^\pi I_{k\theta} d\theta \quad (2.71)$$

In expression (2.69) for the magnetic moment a point-charge-like behavior excludes the range $\gamma < 2$.

The divergences of the so far undetermined radial expressions (2.69)–(2.70) appear in the integrals of Eqs.(2.39)–(2.43) when the lower limits ρ_k approach zero. To outbalance these, one has to introduce a shrinking characteristic radius defined by

$$r_0 = c_0 \varepsilon \quad c_0 > 0 \quad 0 < \varepsilon \ll 1 \quad (2.72)$$

Here c_0 is a positive constant having the dimension of length, and ε is a dimensionless characteristic radius which has the role of a decreasing smallness parameter. Replacing ε by any power of ε would not lead to more generality. Combination of Eqs.(2.69), (2.70), and (2.72) with Eqs.(2.39)–(2.43) gives the result

$$q_0 = 2\pi\varepsilon_0 c_0 G_0 [J_{q\theta}/(\gamma - 1)] (\varepsilon/\rho_q^{\gamma-1}) \quad (2.73)$$

$$M_0 m_0 = \pi^2 (\varepsilon_0^2 C/c^2) c_0^3 G_0^3 \cdot [J_{M\theta} J_{m\theta}/(\gamma - 2)(2\gamma - 1)] \cdot (\varepsilon^3/\rho_M^{\gamma-2} \rho_m^{2\gamma-1}) \quad (2.74)$$

$$s_0 = \pi (\varepsilon_0 C/c^2) c_0^2 G_0^2 \cdot [J_{s\theta}/2(\gamma - 1)] (\varepsilon/\rho_s^{\gamma-1})^2 \quad (2.75)$$

The reason for introducing the compound quantity $M_0 m_0$ in Eq.(2.74) is that this quantity appears as a single entity of all finally obtained relations in the present theory.

For all four quantities q_0 , M_0 , m_0 , s_0 to become finite when ε approaches zero, we must satisfy the conditions[7]

$$\begin{aligned} \rho_q &= \varepsilon^{1/(\gamma-1)} & \rho_M &= \varepsilon^{2/(\gamma-2)} \\ \rho_m &= \varepsilon^{1/(2\gamma-1)} & \rho_s &= \varepsilon^{1/(\gamma-1)} \end{aligned} \quad (2.76)$$

whereas the condition for $M_0 m_0$ to be finite is thus

$$\rho_M^{\gamma-2} \rho_m^{2\gamma-1} = \varepsilon^3 \quad (2.77)$$

Now we only require the configuration represented by q_0 , s_0 and the compound quantity $M_0 m_0$ of Eqs.(2.73)–(2.75) to scale in such a way that the geometry is preserved by becoming independent of ρ_k and ε , within the entire range of small ε . Such a uniform scaling implies that

$$\rho_q = \rho_M = \rho_m = \rho_s = \varepsilon \quad (2.78)$$

and that the radial parameter γ has to approach the value 2 from above, i.e.

$$\gamma(\gamma - 1) = 2 + \delta \quad 0 \leq \delta \ll 1 \quad (2.79)$$

$$\gamma \approx 2 + \delta/3$$

These relations thus hold for condition (2.77) where $M_0 m_0$ is treated as one entity, but not for the two intermediate conditions (2.76) where M_0 and m_0 are treated separately. For both these options $M_0 m_0$ becomes finite in any case.

>From the earlier obtained results of partial integration it is further seen that the contribution from τ_0 in Eq.(2.64) vanishes as well as $J_{M\theta}$ when $\gamma = 2$. In the limit (2.79) the integrands (2.64)–(2.67) can then be replaced by

$$I_{q\theta} = -2\tau_1 + 4\tau_2 \quad (2.80)$$

$$I_{M\theta}/\delta = (\sin \theta)(-\tau_1 + 4\tau_2) \quad (2.81)$$

$$I_{m\theta} = \tau_0\tau_3 - 2(\tau_0\tau_4 + \tau_1\tau_3) + 4(\tau_1\tau_4 + \tau_2\tau_3) - 8\tau_2\tau_4 \quad (2.82)$$

$$I_{s\theta} = (\sin \theta)I_{m\theta} \quad (2.83)$$

With the integrals (2.71) and the definitions

$$A_q \equiv J_{q\theta} \quad A_M \equiv J_{M\theta}/\delta \quad (2.84)$$

$$A_m \equiv J_{m\theta} \quad A_s \equiv J_{s\theta}$$

the integrated quantities (2.73)–(2.75) take the final form

$$q_0 = 2\pi\varepsilon_0 c_0 G_0 A_q \quad (2.85)$$

$$M_0 m_0 = \pi^2 (\varepsilon_0^2 C / c^2) c_0^3 G_0^3 A_M A_m \quad (2.86)$$

$$s_0 = (1/2)\pi (\varepsilon_0 C / c^2) c_0^2 G_0^2 A_s \quad (2.87)$$

for small δ .

Through combination of expressions (2.41), (2.43), (2.58), (2.62), (2.70), (2.72), (2.84), and (2.87), and due to the fact that the ratio A_m/A_s is close to unity, it is readily seen that the radial constant c_0 becomes nearly equal to the Compton wavelength h/m_0c divided by 6π .

2. The Magnetic Flux

According to Eq.(2.34) the magnetic flux function becomes

$$\Gamma = 2\pi r(\sin \theta)A = -2\pi r_0(G_0/c)\rho(\sin \theta)^3 DG \quad (2.88)$$

which vanishes at $\theta = (0, \pi)$. Making use of Eqs.(2.32), (2.60), and (2.72) the flux becomes

$$\Gamma = 2\pi(c_0 G_0/C) \sin^3 \theta \{[\gamma(\gamma - 1) + 2(\gamma - 1)\rho + \rho^2]T - D_\theta T\}(\varepsilon/\rho^{\gamma-1})e^{-\rho} \quad (2.89)$$

This relation shows that the flux increases strongly as ρ decreases towards small values, in accordance with a point-charge-like behavior. To obtain a nonzero and finite magnetic flux when γ approaches the value 2 from above, one has to choose a corresponding dimensionless lower radius limit $\rho_\Gamma = \varepsilon$, in analogy with expressions (2.72) and (2.78). There is then a magnetic flux which intersects the equatorial plane. It is counted from the point $\rho = \rho_\Gamma = \varepsilon$ and outwards, as given by

$$\Gamma_0 = -\Gamma(\rho = \varepsilon, \theta = \pi/2) = 2\pi(c_0 G_0 \varepsilon^{-\delta/3}/C)A_\Gamma \quad (2.90)$$

where

$$A_\Gamma = [D_\theta T - 2T]_{\theta=\pi/2} \quad (2.91)$$

for $\gamma = 2$. The flux (2.90) can be regarded as being generated by a configuration of thin current loops. These have almost all their currents located to a spherical surface with the radius $\rho = \varepsilon$, and the corresponding magnetic field lines cut the equatorial plane at right angles. The described field pattern is based on a separable generating function and applies to this near-field region.

It has to be observed that the flux (2.90) is not necessarily the total flux which is generated by the current system as a whole. There are cases in which magnetic islands are formed above and below the equatorial plane, and where these islands possess an isolated circulating

extra flux which does not intersect the same plane. The total flux Γ_{tot} then consists of the “main flux” $-\Gamma_0$ of Eq.(2.90), plus the extra “island flux” Γ_i , which can be deduced from the function (2.89), as will be shown later. This type of field geometry turns out to prevail in the analysis which follows.

In the analysis of the contribution from the magnetic islands we introduce the normalized flux function defined by Eq.(2.89) in the upper half-plane of the sphere $\rho = \varepsilon \ll 1$. It becomes

$$\Psi \equiv \Gamma(\rho = \varepsilon, \theta)/2\pi(c_0 G_0/C) = \sin^3 \theta (D_\theta T - 2T) \quad (2.92)$$

In the range of increasing θ , the radial magnetic field component B_ρ then vanishes along the lines $\theta = \theta_1$ and $\theta = \theta_2$, whereas the component B_θ vanishes along the lines $\theta = \theta_3$ and $\theta = \theta_4$. Here θ_3 is situated between θ_1 and θ_2 , and θ_4 between θ_2 and $\pi/2$. Consequently, the increase of θ from the axis at $\theta = 0$ first leads to an increasing flux Ψ up to a maximum at the angle $\theta = \theta_1$. Then there follows an interval $\theta_1 < \theta < \theta_2$ of decreasing flux, down to a minimum at $\theta = \theta_2$. In the range $\theta_2 \leq \theta \leq \pi/2$ there is again an increasing flux, up to the value

$$\Psi_0 = \Psi(\pi/2) = A_\Gamma \quad (2.93)$$

which is equal to the main flux.

We further introduce the parts of the integrated magnetic flux defined by

$$\Psi_1 = \Psi(\theta_1) - \Psi(0) = \Psi(\theta_1) \quad (2.94)$$

$$\Psi_2 = \Psi(\pi/2) - \Psi(\theta_2)$$

and associated with the ranges $0 < \theta < \theta_1$ and $\theta_2 < \theta < \pi/2$. The sum of these parts includes the main flux Ψ_0 , plus an outward directed flux from one magnetic island. The contribution from the latter becomes $\Psi_1 + \Psi_2 - \Psi_0$. The total flux which includes the main flux and that from two islands is then given by

$$\Psi_{tot} = 2(\Psi_1 + \Psi_2 - \Psi_0) + \Psi_0 \quad (2.95)$$

which also can be written as

$$\Psi_{tot} = f_{\Gamma f} \Psi_0 \quad (2.96)$$

$$f_{\Gamma f} = [2(\Psi_1 + \Psi_2) - \Psi_0]/\Psi_0$$

where $f_{\Gamma f} > 1$ is a corresponding flux factor.

3. Quantum Conditions of the Electron

The analysis is now at a stage where the relevant quantum conditions can be imposed. For the angular momentum (2.54) and its associated charge relation (2.55) the result becomes

$$q^* = (f_0 A_q^2 / A_s)^{1/2} \quad (2.97)$$

according to Eqs.(2.68)–(2.71), (2.78) and (2.84) in the limit $\gamma = 2$.

For the magnetic moment condition (2.56) reduces to

$$A_M A_m / A_q A_s = 1 + \delta_M \quad (2.98)$$

when applying Eqs.(2.85)–(2.87).

Magnetic flux quantization is expressed by condition (2.59). Combination of Eqs.(2.93), (2.73), (2.75) and (2.84) then yields

$$8\pi f_{\Gamma q} A_\Gamma A_q = A_s \quad (2.99)$$

where $f_{\Gamma q}$ is the flux factor being *required* by the corresponding quantum condition. This factor should not be confused with the factor $f_{\Gamma f}$ of Eq.(2.96) which *results* from the magnetic field geometry. Only when one arrives at a self-consistent solution will these two factors become equal to the common flux factor

$$f_\Gamma = f_{\Gamma f} = f_{\Gamma q} \quad (2.100)$$

4. Comparison with the Conventional Renormalization Procedure

The previous analysis has shown that well-defined convergent and nonzero integrated physical quantities can be obtained in a point-charge-like steady state, thereby avoiding the problem of an infinite self-energy. Attention may here be called to Ryder[3] who has stressed that, despite the success of the conventional renormalization procedure, a more physically satisfactory way is needed concerning the infinite self-energy problem. Possibly the present theory could provide such an alternative, by tackling this problem in a more surveyable manner. The finite result due to a difference between two “infinities” in renormalization theory, i.e. by adding extra counter-terms to the Lagrangian, is then replaced by a finite result

obtained from the product of an “infinity” with a “zero”, as determined by the combination of the present divergent integrands with a shrinking characteristic radius.

As expressed by Eq.(2.72), the latter concept also has an impact on the question of Lorentz invariance of the electron radius. In the limit $\varepsilon = 0$ of a vanishing radius r_0 corresponding to a structureless mass point, the deductions of this chapter will thus in a formal way satisfy the requirement of such an invariance. At the same time the analytic continuation of the obtained solutions can also be applied to the physically relevant case of a very small but nonzero radius of a configuration having an internal structure.

5. *Variational Analysis of the Integrated Charge*

The elementary electronic charge has so far been considered as an independent and fundamental physical constant of nature, being determined by measurements only[48]. However, since it appears to represent the smallest quantum of free electric charge, the question can be raised whether there is a more profound reason for such a minimum charge to exist, possibly in terms of a quantized variational analysis.

The present theory can provide the basis for such an analysis. In a first attempt one would use a conventional procedure including Lagrange multipliers in searching for an extremum of the normalized charge q^* of Eq.(2.97), under the subsidiary quantum conditions (2.98)–(2.100) and (2.96). The available variables are then the amplitudes $a_{2\nu-1}$ and $a_{2\nu}$ of the polar function (2.61). Such an analysis has unfortunately been found to become quite complicated[12], partly due to its nonlinear character and a high degree of the resulting equations. But there is even a more serious difficulty which upsets such a conventional procedure. The latter namely applies only when there are well-defined and localized points of an extremum, in the form of a maximum or minimum or a saddle-point in parameter space, but not when such single points are replaced by a flat plateau which has the effect of an infinite number of extremum points being distributed over the same space.

The plateau behavior is in fact what occurs here[12], and an alternative approach therefore has to be elaborated. The analysis then proceeds by successively including an increasing number of amplitudes $(a_1, a_2, a_3, \dots)$ which are being “swept” (scanned) across their entire range of variation. The flux factor (2.100) has at the same time to be determined in a self-consistent way through an iteration process. For each iteration the lowest occurring value of

q^* can then be determined. Thereby both conditions (2.98) and (2.99) and the flux factors of Eqs.(2.96) and (2.99) include variable parameters. Thus the flux factor $f_{\Gamma f}$ of Eq.(2.99) does not become constant but varies with the amplitudes of the polar function (2.61) when there are magnetic islands which contribute to the magnetic flux. At a first sight this appears also to result in a complicated and work-consuming process. However, the flat plateau behavior is found to simplify the corresponding iteration scheme and the physical interpretation of its results.

In the numerical analysis which follows, solutions with two real roots have always been found. Of these roots only that resulting in the lowest value of q^* will be treated in detail in the following parts of this chapter.

Here the analysis will be illustrated by an example where four amplitudes (a_1, a_2, a_3, a_4) are included in the polar function (2.61). The normalized charge q^* can then be plotted as a function of the two amplitudes a_3 and a_4 which are swept across their entire range of variation, but for a fixed flux factor $f_{\Gamma} = f_{\Gamma q} = 1.82$. The result is shown in Fig.2.1 for the lowest of the two roots obtained for q^* . In fact, the corresponding result of a self-consistent analysis which takes Eq.(2.100) into account, leads only to small modifications which hardly become visible on the scale of Fig.2.1. From the figure is seen that there is a steep barrier in its upper part, from which q^* drops down to a flat plateau being close to the level $q^* = 1$.

When further increasing the number of amplitudes, the results were still found to become plateau-like. The detailed shape of the plateau was seen to become slightly “warped”, thereby locally being partly below and partly above the level $q^* = 1$, i.e. in a range $0.97 \leq q^* \leq 1.03$.

The flat plateau behavior of q^* at an increasing number of included amplitudes can be understood from the fact that the expression (2.97) for the normalized charge solely depends on the profile shape. It therefore becomes a slow function of the higher “multipole” terms in the expansion (2.61).

6. Force Balance of the Electron Model

The fundamental description of a charged particle in conventional theory is deficient in several respects. Thus, an equilibrium cannot be maintained by the classical electrostatic forces, but has been assumed to require forces of a nonelectromagnetic character to be




FIG. 2.1: The normalized electron charge $q^* \equiv |q_0/e|$ as a function of the two amplitudes a_3 and a_4 , for solutions satisfying the subsidiary quantum conditions for a fixed flux factor $f_\Gamma = f_{\Gamma q} = 1.82$, and being based on a polar function T having four amplitudes (a_1, a_2, a_3, a_4) . The profile of q^* consists of a steep “barrier” in the upper part of the figure, and a flat “plateau” in the lower part. The plateau is close to the level $q^* = 1$. The figure only demonstrates the ranges of the real solutions of the first (lowest) root. The deviations of this profile from that obtained for the self-consistent solutions which obey condition (2.100) are hardly visible on the scale of the figure.

present[2, 4, 45]. In other words, the electron would otherwise “explode” under the action of its self-charge. Here it will be shown that a steady electromagnetic equilibrium can under certain conditions be established by the present extended theory.

We now turn to the momentum balance of the present axisymmetric electron model in a frame (r, θ, φ) of spherical coordinates. The integral form of the equivalent forces is in a

steady state given by Eqs.(2.20) and (2.21), on the form

$$\mathbf{F} = \mathbf{F}_e + \mathbf{F}_m = \int \bar{\rho}(\mathbf{E} + \mathbf{C} \times \mathbf{B}) dV \quad (2.101)$$

When first considering the polar direction represented by the angle θ , it is readily seen that the corresponding integrated force F_θ vanishes due to the axial symmetry.

Turning then to the radial direction, the earlier obtained results of Eqs.(2.30)–(2.36) and (2.44) are applied to obtain the radial force component[14]

$$F_r = -2\pi\epsilon_0 G_0^2 \int \int [DG + D(s^2 DG)] \cdot \left[\frac{\partial G}{\partial \rho} - \frac{1}{\rho} s^2 DG \right] \rho^2 s d\rho d\theta \quad (2.102)$$

where $s = \sin \theta$. For the point-charge-like model of Eqs.(2.60)–(2.61) with $G = RT$, $\gamma \longrightarrow 2$ and $R \longrightarrow 1/\rho^2$ at small ρ , we then have

$$\rho^2 DG = D_\theta T - 2T \quad (2.103)$$

to be inserted into the integrand of expression (2.102). The latter can then be represented by the form

$$F_r = I_+ - I_- \quad (2.104)$$

where I_+ and I_- are the positive and negative contributions to the radial force F_r .

Consequently, there will arise an integrated radial force balance in the form of electromagnetic confinement when $I_+ = I_-$. This applies to very small but nonzero values of the characteristic radius (2.72), i.e. when not proceeding all the way to the structureless point case where $r_0 = 0$.

The integrals (2.162) and (2.164) are now applied to the plateau region. It is then found that the ratio I_+/I_- decreases from $I_+/I_- = 1.27$ at $q^* = 0.98$ to $I_+/I_- = 0.37$ at $q^* = 1.01$, thus passing the equilibrium point $I_+/I_- = 1$ at $q^* \cong 0.988$. The remaining degrees of freedom being available in the parameter ranges of the plateau have then been used up by the condition of a radially balanced equilibrium.

To sum up, the variational analysis has resulted in a plateau region at the perimeter of which the normalized charge q^* varies in the range $0.97 < q^* < 1.03$, and where the lowest possible value $q^* \cong 0.97$ is obtained. However, this latter value does not satisfy the requirement of a radial force balance. Such a balance is on the other hand realized within the plateau region, but at the value $q^* \cong 0.99$, i.e. where the deduced charge only deviates by about one percent from the experimental value of the elementary charge.

The reason for this deviation of the deduced charge from the experimentally determined value is not clear at this stage, but a quantum mechanical correction may offer one possibility for its removal. A successful modification of the magnetic moment has earlier been made by Schwinger and Feynman, as given by Eq.(2.56). Without getting into the details of the advanced and work-consuming analysis which leads to this correction, it may here be proposed that also the magnetic flux and its related quantum condition would then have to be modified into the form[15]

$$\begin{aligned}\tilde{f}_{\Gamma q} &= (\bar{A}_s/8\pi\bar{A}_{\Gamma}\bar{A}_q)(1 + \delta_{\Gamma}) \\ \delta_{\Gamma} &= c_{\Gamma}/f_0\end{aligned}\tag{2.105}$$

where c_{Γ} is a so far unspecified constant. For this proposal to lead to an agreement with the experimental value $q^* = 1$, a modified self-consistency relation

$$\tilde{f}_{\Gamma q} = \bar{f}_{\Gamma f}\tag{2.106}$$

would have to be satisfied, and the constant c_{Γ} would then have to be of the order of unity.

At the extremely small dimensions of the present point-charge-like model, it should become justified to consider the integrated force balance instead of its localized counterpart. The quantum mechanical wave nature is then expected to have a smoothing and overlapping effect on the local variations within the configuration. In addition to this, a requirement of the local electrodynamic force of Eq.(2.19) to vanish identically, would result in the unacceptable situation of an overdetermined system of basic equations.

The obtained deviation of q^* by about one percent only from the value $q^* = 1$ can, in itself, be interpreted as an experimental support of the present theory. This is particularly the case as the deduced result has been obtained from two independent aspects, namely the minimization of the charge by a variational analysis, and the determination of the charge from the requirement of a radial balance of forces.

E. Axisymmetric Wave Modes and Their Photon Models

During several decades a number of investigators have discussed the nature of light and photon physics, not only in relation to the propagation of plane wave fronts but also to axisymmetric wave packets, the concept of a photon rest mass, the existence of a magnetic

field component in the direction of propagation, and to an associated angular momentum. The aim is here to elaborate a model of the individual photon in the capacity of a boson particle with an angular momentum (spin), propagating with preserved geometry and limited extensions in a defined direction of space. This leads to the concept of cylindrical waves and wave packets.

1. Elementary Normal Modes

A cylindrical frame (r, φ, z) is introduced where φ is an ignorable coordinate. In this frame the velocity vector is assumed to have the form

$$\mathbf{C} = c(0, \cos \alpha, \sin \alpha) \quad (2.107)$$

with a constant angle α . We further define the operators

$$D_1 = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \quad (2.108)$$

$$D_2 = \frac{\partial}{\partial t} + c(\sin \alpha) \frac{\partial}{\partial z} \quad (2.109)$$

The basic equations (2.28) and (2.29) are then represented by the system

$$\left(D_1 - \frac{1}{r^2}\right) E_r = \frac{\partial}{\partial r} (\nabla \cdot \mathbf{E}) \quad (2.110)$$

$$\left(D_1 - \frac{1}{r^2}\right) E_\varphi = \frac{1}{c} (\cos \alpha) \frac{\partial}{\partial t} (\nabla \cdot \mathbf{E}) \quad (2.111)$$

$$D_1 E_z = \left[\frac{\partial}{\partial z} + \frac{1}{c} (\sin \alpha) \frac{\partial}{\partial t} \right] (\nabla \cdot \mathbf{E}) \quad (2.112)$$

and

$$D_2 (\nabla \cdot \mathbf{E}) = 0 \quad (2.113)$$

Three of the four relations (2.110)–(2.113) thereby form a complete system of equations, because Eq.(2.113) originates from Eq.(2.29). By further defining the operator

$$D_3 = \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \quad (2.114)$$

Eq. (2.110) reduces to

$$D_3 E_r = \frac{\partial^2 E_z}{\partial r \partial z} \quad (2.115)$$

Combination with Eq.(2.111) then yields

$$D_3 \left(D_1 - \frac{1}{r^2}\right) E_\varphi = \frac{1}{c} (\cos \alpha) \frac{\partial^2}{\partial z \partial t} D_1 E_z \quad (2.116)$$

Normal modes depending on z and t as $\exp[i(-\omega t + kz)]$ are now considered, first in the conventional case where $\nabla \cdot \mathbf{E} = 0$, and then for the EMS mode where $\nabla \cdot \mathbf{E} \neq 0$. In fact, there are a number of choices with respect to the form (2.107) as represented by $\pm \cos \alpha$ and $\pm \sin \alpha$ which satisfy the condition of Eq.(2.5), thereby corresponding to the two directions along and around the axis of symmetry.

In the case of a vanishing electric field divergence, Eq.(2.110)–(2.112) reduce to those of an axisymmetric EM mode. Then

$$\left[\bar{D}_\rho - \left(\frac{1}{\rho^2} \right) \right] (E_r, E_\varphi) = 0 \quad (2.117)$$

$$\bar{D}_\rho E_z = 0$$

where

$$\bar{D}_\rho = \frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} \quad (2.118)$$

The solutions of Eqs.(2.117) become

$$E_r = k_{r1}\rho + k_{r2}/\rho \quad (2.119)$$

$$E_\varphi = k_{\varphi1}\rho + k_{\varphi2}/\rho$$

$$E_z = k_{z1} \ln \rho + k_{z2}$$

where $k_{r1}, k_{r2}, k_{\varphi1}, k_{\varphi2}, k_{z1}$ and k_{z2} are constants. The magnetic field is obtained from Eq.(2.9).

A similar divergence of the field at the axis $\rho = 0$ and at large ρ was already realized by Thomson[5] and further discussed by Heitler[4], as well as by Hunter and Wadlinger[49]. It leaves the problem with the radial dependence unresolved. Thus the field does not converge within the entire vacuum space, and cannot be made to vanish at large radial distances as long as $k_{r1}, k_{\varphi1}$ and k_{z1} differ from zero. All parts of the solutions (2.119) further result in integrals of the energy density which become divergent when being extended all over the vacuum space. An additional and even more severe point of concern is due to the electric field divergence of Eq.(2.119) which becomes

$$\nabla \cdot \mathbf{E} = 2k_{r1} + ik(k_{z1} \ln \rho + k_{z2}) \quad (2.120)$$

and has to vanish identically for all (ρ, k) in the case of conventional electromagnetic waves. Thereby the last part of Eq.(2.120) requires the parallel electric field component E_z to

vanish identically, but then it also becomes necessary for the constant k_{r1} in Eq.(2.120) to disappear. There are further analogous solutions for the magnetic field, by which also the axial component B_z vanishes in the conventional case. In its turn, this leads to a Poynting vector (2.22) having only a component in the direction of propagation, thus resulting in a vanishing angular momentum (spin) with respect to the axial direction.

Returning to the axisymmetric EMS mode of Eqs.(2.107)–(2.116), a dispersion relation

$$\omega = kv \quad v = c(\sin \alpha) \quad (2.121)$$

is obtained from Eq.(2.113), having the phase and group velocities v . Here we limit ourselves to positive values of $\cos \alpha$ and $\sin \alpha$. With the dispersion relation (2.121), Eq.(2.116) takes the form

$$\left[\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} - \frac{1}{r^2} - k^2(\cos \alpha)^2 \right] E_\varphi = -(\operatorname{tg} \alpha) \left[\frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} - k^2(\cos \alpha)^2 \right] E_z \quad (2.122)$$

We can now introduce a *generating function*

$$G_0 \cdot G = E_z + (\cot \alpha) E_\varphi \quad (2.123)$$

$$G = R(\rho) e^{i(-\omega t + kz)}$$

where G_0 stands for the amplitude, G for a normalized form, $\rho = r/r_0$ is again a normalized coordinate with respect to the characteristic radius r_0 of the configuration, and $R(\rho)$ is a dimensionless function of ρ . With the operator of Eq.(2.118), the definition

$$D = \bar{D}_\rho - \theta^2(\cos \alpha)^2 \quad \theta = kr_0 \quad (2.124)$$

and using Eqs.(2.121), (2.115), and (2.9), insertion into Eq.(2.122) results in

$$E_r = -iG_0[\theta(\cos \alpha)^2]^{-1} \cdot \frac{\partial}{\partial \rho} [(1 - \rho^2 D)G] = -\frac{1}{r_0} \frac{\partial \phi}{\partial \rho} + i\omega A_r \quad (2.125)$$

$$E_\varphi = G_0(\operatorname{tg} \alpha) \rho^2 D G = i\omega A_\varphi \quad (2.126)$$

$$E_z = G_0(1 - \rho^2 D)G = -ik\phi + i\omega A_z \quad (2.127)$$

and

$$B_r = -G_0[c(\cos \alpha)]^{-1} \cdot \rho^2 D G = -ikA_\varphi \quad (2.128)$$

$$B_\varphi = -iG_0(\sin \alpha)[\theta c(\cos \alpha)^2]^{-1} \cdot \frac{\partial}{\partial \rho} [(1 - \rho^2 D)G] = ikA_r - \frac{1}{r_0} \frac{\partial A_z}{\partial \rho} \quad (2.129)$$

$$B_z = -iG_0[\theta c(\cos \alpha)]^{-1} \left(\frac{\partial}{\partial \rho} + \frac{1}{\rho} \right) \cdot (\rho^2 DG) = \frac{1}{r_0} \frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\varphi) \quad (2.130)$$

This result makes possible a choice of generating functions and corresponding modes which are physically relevant within the entire vacuum space, and which become consistent with the imposed quantum conditions. The solutions (2.125)–(2.127) are reconfirmed by insertion into the basic equations. The result implies that the three equations (2.110)–(2.112) do not become independent of each other.

Due to the dispersion relation (2.121) the EMS mode propagates at phase and group velocities which are smaller than c . Not to get into conflict with the experimental observations by Michelson and Moreley, the condition

$$0 < \cos \alpha \ll 1 \quad (2.131)$$

has then to be taken into account. Since the velocity v of Eq.(2.121) becomes slightly less than c , the condition (2.131) will be related to a very small but nonzero rest mass of the photon, as shown later in more detail.

With the condition (2.131) the present mode has a dominant radially polarized electric field component (2.125), and an associated magnetic field component (2.129). The field configuration of Eqs.(2.125)–(2.130) thus differs in several respects from elliptically, circularly and linearly polarized plane waves.

The components (2.125)–(2.130) are reconcilable with the field configuration by Evans and Vigier[50, 51], in the sense that all components are nonzero and form a helical structure which also includes an axial magnetic field component in the direction of propagation. The present result is, however, not identical with that by Evans and Vigier.

So far the deductions have been performed with respect to the laboratory frame K . In the present case where the phase and group velocities $v < c$, a physically relevant rest frame K' can be defined, but not in the conventional case where there is no rest mass, $\cos \alpha = 0$, and $v = c$. The details of a transformation to the rest frame are demonstrated elsewhere[10].

2. Wave Packets

From a spectrum of normal modes with different wave numbers, a wave packet solution is now being formed which has finite radial and axial extensions and a narrow line width in

wavelength space. Such a line width is consistent with experimental observations. We are free to rewrite the amplitude factor of the generating function (2.123) as

$$G_0 = g_0(\cos \alpha)^2 \quad (2.132)$$

The normal modes are superimposed to result in a wave packet having the amplitude

$$A_k = \left(\frac{k}{k_0^2}\right) e^{-z_0^2(k-k_0)^2} \quad (2.133)$$

in the interval dk and being centered around the wave number k_0 . Further $2z_0$ represents the effective axial length of the packet.

The physically relevant restriction to a narrow line width implies that the amplitude (2.133) drops to $1/e$ of its maximum value for a small deviation of k from the maximum at $k = k_0$. With $\Delta k = k - k_0 = 1/z_0$ this implies that $\Delta k/k_0 = 1/k_0 z_0 = \lambda_0/2\pi z_0 \ll 1$, where $\lambda_0 = 2\pi/k_0$ is the average wave length of the packet. With $\bar{z} = z - vt$ and $v = c(\sin \alpha)$, and after working out the integration with respect to k , the spectral averages of the packet field components become

$$\bar{E}_r = -iE_0[R_5 + (\theta'_0)^2 R_2] \quad (2.134)$$

$$\bar{E}_\varphi = E_0\theta_0(\sin \alpha)(\cos \alpha) \cdot [R_3 - (\theta'_0)^2 R_1] \quad (2.135)$$

$$\bar{E}_z = E_0\theta_0(\cos \alpha)^2[R_4 + (\theta'_0)^2 R_1] \quad (2.136)$$

$$\bar{B}_r = -\left(\frac{1}{c}\right)(\sin \alpha)^{-1} E_\varphi \quad (2.137)$$

$$\bar{B}_\varphi = \left(\frac{1}{c}\right)(\sin \alpha) E_r \quad (2.138)$$

$$\bar{B}_z = -i\left(\frac{1}{c}\right) E_0(\cos \alpha)[R_8 - (\theta'_0)^2 R_7] \quad (2.139)$$

where $\theta_0 = k_0 r_0$, $\theta'_0 = \theta_0(\cos \alpha)$,

$$E_0 = E_0(\bar{z}) = \left(\frac{g_0}{k_0 r_0}\right) \left(\frac{\sqrt{\pi}}{k_0 z_0}\right) \cdot \exp \left[-\left(\frac{\bar{z}}{2z_0}\right)^2 + ik_0 \bar{z} \right] \quad (2.140)$$

and

$$R_1 = \rho^2 R \quad R_2 = \frac{d}{d\rho}(\rho^2 R) \quad R_3 = \rho^2 \bar{D}_\rho R \quad (2.141)$$

$$R_4 = (1 - \rho^2 \bar{D}_\rho) R \quad R_5 = \frac{d}{d\rho}[(1 - \rho^2 \bar{D}_\rho) R] \quad (2.142)$$

$$R_6 = \bar{D}_\rho[(1 - \rho^2 \bar{D}_\rho) R] \quad R_7 = \left(\frac{d}{d\rho} + \frac{1}{\rho}\right)(\rho^2 R) \quad (2.143)$$

with the operator $\bar{D}_\rho$ given by Eq.(2.118).

It should be noticed that expressions (2.137)–(2.139) are approximate, on account of the restriction to a narrow line width. Therefore the condition $\nabla \cdot \bar{\mathbf{B}} = 0$ is only satisfied approximately. In the limit of zero line width we would on the other hand be back to the exact form (2.128)–(2.130) for an elementary normal mode where $\nabla \cdot \mathbf{B} = 0$.

For the wave packet fields (2.134)–(2.139) the components $(\bar{E}_\varphi, \bar{E}_z, \bar{B}_r)$ are in phase with the generating function (2.123), whereas the components $(\bar{E}_r, \bar{B}_\varphi, \bar{B}_z)$ are ninety degrees out of phase with it. In the analysis which follows we choose a generating function which is symmetric with respect to the axial centre $\bar{z} = 0$ of the moving wave packet. Thus

$$G = R(\rho) \cos k\bar{z} \quad (2.144)$$

when the real parts of the forms (2.123) and (2.140) are adopted. Then G and $(\bar{E}_\varphi, \bar{E}_z, \bar{B}_r)$ are symmetric and $(\bar{E}_r, \bar{B}_\varphi, \bar{B}_z)$ are antisymmetric functions of $\bar{z}$ with respect to the centre $\bar{z} = 0$.

3. Spatially Integrated Field Quantities

When elaborating a model for the individual photon, integrals have to be formed to obtain net values of the electric charge q , magnetic moment M , the total mass m which represents the total energy, and of the angular momentum (spin) s . Thereby the limits of $\bar{z}$ are $\pm\infty$, whereas those of ρ become unspecified until further notice.

The integrated charge becomes

$$q = \varepsilon_0 \int \nabla \cdot \bar{\mathbf{E}} \, dV = 2\pi\varepsilon_0 \int \left\{ \frac{\partial}{\partial r} (r \int_{-\infty}^{+\infty} \bar{E}_r \, d\bar{z}) + r [\bar{E}_z]_{-\infty}^{+\infty} \right\} dr = 0 \quad (2.145)$$

because $\bar{E}_r$ is antisymmetric and $\bar{E}_z$ symmetric with respect to $\bar{z}$.

The integrated magnetic moment is related to the component

$$j_\varphi = \bar{\rho} C_\varphi = \varepsilon_0 (\nabla \cdot \bar{\mathbf{E}}) c (\cos \alpha) \quad (2.146)$$

of the space-charge current density. Thus

$$\begin{aligned} M &= \int \int_{-\infty}^{+\infty} \bar{\rho} c (\cos \alpha) \pi r^2 \, dr d\bar{z} = \\ &= \pi \varepsilon_0 c (\cos \alpha) \int \left\{ r \frac{\partial}{\partial r} (r \int_{-\infty}^{+\infty} \bar{E}_r \, d\bar{z}) + r^2 [\bar{E}_z]_{-\infty}^{+\infty} \right\} dr = 0 \end{aligned} \quad (2.147)$$

for the same symmetry reasons as those applying to the charge (2.145). It should be observed that, even if the net magnetic moment vanishes, the local magnetic field remains nonzero, as well as the local charge density.

An equivalent total mass can be defined from the electromagnetic field energy, due to the energy relation by Einstein. Consequently we write

$$m = \left(\frac{1}{c^2}\right) \int w_f dV \quad (2.148)$$

as given by the field energy density of Eq.(2.25). Applying the condition (2.131) and using Eqs.(2.134)–(2.139), we then have

$$m \cong 2\pi(\varepsilon_0/c^2) \int_{-\infty}^{+\infty} \int r |\bar{E}_r^2| dr d\bar{z} \quad (2.149)$$

When adopting the form (2.144) with an antisymmetric field $\bar{E}_r$, and introducing the integral

$$J_z = \int_{-\infty}^{+\infty} (\sin k_0 \bar{z})^2 e^{-2(\bar{z}/2z_0)^2} d\bar{z} \cong z_0(\pi/2)^{1/2} \quad (2.150)$$

in the limit $k_0 z_0 \gg 1$, the total mass finally becomes

$$m = a_0 W_m = h\nu_0/c^2 \quad (2.151)$$

$$W_m = \int \rho R_5^2 d\rho$$

where

$$a_0 = \varepsilon_0 \pi^{5/2} \sqrt{2} z_0 (g_0 / c k_0^2 z_0)^2 \equiv 2a_0^* g_0^2 \quad (2.152)$$

For this wave packet the energy relations by Planck and Einstein also combine to

$$mc^2 = h\nu_0 = (hc/\lambda_0)(\sin \alpha) \quad (2.153)$$

$$\lambda_0 = 2\pi/k_0$$

where use has been made of the dispersion relation (2.121) applied to the wave packet as a whole, i.e.

$$\omega_0 = 2\pi\nu_0 = k_0 c (\sin \alpha) \quad (2.154)$$

$$\nu_0 = (c/\lambda_0)(\sin \alpha)$$

Here the slightly reduced phase and group velocity of Eq.(2.121) can be considered as being associated with a very small nonzero rest mass

$$m_0 = m[1 - (v/c)^2]^{1/2} = m(\cos \alpha) \quad (2.155)$$

Turning to the momentum balance governed by Eqs.(2.20)–(2.22), all integrated components of the forces (2.21) in the r and φ directions will vanish on account of the axial symmetry, and since the φ component of $\bar{\mathbf{F}}_e + \bar{\mathbf{F}}_m$ will vanish on account of Eq.(2.137). The only components which remain are

$$\begin{aligned}\bar{F}_{ez} &= \int \bar{\rho} \bar{E}_z dV \\ \bar{F}_{mz} &= - \int \bar{\rho} c(\cos \alpha) \bar{B}_r dV\end{aligned}\tag{2.156}$$

where

$$\bar{\rho}/\varepsilon_0 = \frac{1}{r} \frac{\partial}{\partial r}(e \bar{E}_r) + \frac{\partial}{\partial z} \bar{E}_z\tag{2.157}$$

Here $\bar{\rho}$ becomes symmetric and $(\bar{E}_z, \bar{B}_r)$ antisymmetric with respect to $\bar{z}$. The integrated forces (2.156) will for this reason also vanish, and the balance equation then acquires its conventional form. Consequently, the conventional expressions[41]

$$\mathbf{s} = \mathbf{r} \times \mathbf{S}/c^2 \quad \mathbf{S} = \bar{\mathbf{E}} \times \bar{\mathbf{B}}/\mu_0\tag{2.158}$$

apply for the density $\mathbf{s}$ of the angular momentum, where $\mathbf{r}$ is the radius vector from the origin and $\mathbf{S}$ is the Poynting vector. The density of angular momentum now becomes

$$s_z = \varepsilon_0 r (\bar{E}_z \bar{B}_r - \bar{E}_r \bar{B}_z) \cong -\varepsilon_0 r \bar{E}_r \bar{B}_z\tag{2.159}$$

when condition (2.131) applies and use is made of Eqs.(2.134), (2.136), (2.137), and (2.139). The total angular momentum is given by

$$s = \int s_z dV = -2\pi\varepsilon_0 \int_{-\infty}^{+\infty} \int r^2 \bar{E}_r \bar{B}_z dr d\bar{z}\tag{2.160}$$

and it reduces to the final form

$$\begin{aligned}s &= a_0 r_0 c(\cos \alpha) W_s = h/2\pi \\ W_s &= - \int \rho^2 R_5 R_7 d\rho\end{aligned}\tag{2.161}$$

for the model of a photon having the angular momentum of a boson.

The disappearance of the total integrated volume forces is also supported by the fact that the solutions (2.125)–(2.130) and (2.134)–(2.139) result in field strengths $\mathbf{E} + \mathbf{C} \times \mathbf{B}$ and $\bar{\mathbf{E}} + \mathbf{C} \times \bar{\mathbf{B}}$ which are readily seen to vanish in lowest order when $|\cos \alpha| \ll 1$.

Here it is noticed that, with the inclusion of a dimensionless profile factor, relations (2.161) and (2.151) indicate that the form of the angular momentum agrees with the simple picture of a mass which rotates at the velocity $c(\cos \alpha)$ around the axis at the distance r_0 .

The result (2.161) and Eq.(2.155) show that a nonzero angular momentum requires a nonzero rest mass to exist, as represented by $\cos \alpha \neq 0$. Moreover, the factor $\cos \alpha$ in the expression (2.161) and Eq.(2.107) for the velocity vector $\mathbf{C}$ indicate that the angular momentum, as well as the nonzero rest mass, are associated with the component C_φ of circulation around the axis of symmetry.

First a generating function will now be considered which is finite at the axis $\rho = 0$, and which tends to zero at large ρ . These requirements are fulfilled by

$$R(\rho) = \rho^\gamma e^{-\rho} \quad (2.162)$$

In principle, the factor in front of the exponential part would in a general case have to be replaced by series of positive or negative powers of ρ , but since we will here proceed to the limit of large γ , only one term becomes sufficient. Moreover, the exponential factor in Eq.(2.162) has been included to secure the general convergence of any moment with R at large ρ . In any case, the forthcoming final results will be found not to depend explicitly on this exponential factor.

In the evaluation of expressions (2.151) and (2.161) for W_m and W_s the Euler integral

$$J_{2\gamma-2} = \int_0^\infty \rho^{2\gamma-2} e^{-\rho} d\rho = 2^{-(2\gamma+3)} \Gamma(2\gamma+1) \quad (2.163)$$

appears in terms of the gamma function $\Gamma(2\gamma+1)$. For $\gamma \gg 1$ only the dominant terms of the functions R_5 and R_7 of Eqs.(2.142) and (2.143) will prevail, and the result becomes

$$W_m = 2^{-(2\gamma+4)} \gamma^6 (2\gamma-1) \cdot \Gamma(2\gamma+1) = W_s / \gamma \quad (2.164)$$

The function (2.162) has a maximum at the radius

$$\hat{r} = \gamma r_0 \quad (2.165)$$

which becomes sharply defined at large γ , in analogy with an earlier obtained result[10]. Combination of Eqs.(2.151), (2.153), (2.161), (2.164), and (2.165) leads to an effective photon diameter

$$2\hat{r} = \frac{\lambda_0}{\pi(\cos \alpha)} \quad (2.166)$$

This result applies not only to an individual photon of the effective radius $\hat{r}$, but can also stand for the corresponding radius of a radially polarized dense multiphoton beam.

Turning then to the alternative of a radial part which *diverges* at the axis, the form

$$R(\rho) = \rho^{-\gamma} e^{-\rho} \quad \gamma > 0 \quad (2.167)$$

is taken into consideration, being identical with the form (2.60) for the electron model. When the radial variable increases monotonically, the function (2.167) decreases from large values, down to $R = e^{-1}$ at $\rho = 1$, and further to very small values when ρ becomes substantially larger than 1. Thus $r = r_0$ can be taken as a characteristic radial dimension of the configuration.

To obtain *finite* values of the integrated total mass m and angular momentum s of Eqs.(2.151) and (2.161), a special procedure similar to that applied to the electron model is being applied. The lower limits of the integrals (2.151) and (2.161) are specified by

$$\begin{aligned} W_m &= \int_{\rho_m}^{\infty} \rho R_5^2 d\rho \\ W_s &= - \int_{\rho_s}^{\infty} \rho^2 R_5 R_7 d\rho \end{aligned} \quad (2.168)$$

where $\rho_m \ll 1$ and $\rho_s \ll 1$. Making the choice $\gamma \gg 1$, these integrals reduce to

$$W_m = \frac{1}{2} \gamma^5 \rho_m^{-2\gamma} \quad W_s = \frac{1}{2} \gamma^5 \rho_s^{-2\gamma+1} \quad (2.169)$$

To secure finite values of m and s we must now permit the characteristic radius r_0 and the factor g_0 in Eqs.(2.132), (2.151), (2.152), and (2.161) to “shrink” to very small but nonzero values, as the lower limits ρ_m and ρ_s approach zero. This is attained by introducing the relations

$$r_0 = c_r \cdot \varepsilon \quad c_r > 0 \quad (2.170)$$

$$g_0 = c_g \cdot \varepsilon^\beta \quad c_g > 0 \quad (2.171)$$

where $0 < \varepsilon \ll 1$, $\beta > 0$, c_r is a constant with the dimension of length, and c_g one with the dimension of electrostatic potential. Eqs.(2.151), (2.152), (2.153), (2.161), and (2.169) now combine to

$$m = a_0^* \gamma^5 c_g^2 (\varepsilon^{2\beta} / \rho_m^{2\gamma}) \cong h / \lambda_0 c \quad (2.172)$$

$$s = a_0^* \gamma^5 c_g^2 c_r c (\cos \alpha) \cdot (\varepsilon^{2\beta+1} / \rho_s^{2\gamma-1}) = h / 2\pi \quad (2.173)$$

To obtain finite values of both m and s , it is then necessary to satisfy the conditions

$$\rho_m = \varepsilon^{\beta/\gamma} \quad \rho_s = \varepsilon^{(2\beta+1)/(2\gamma-1)} \quad (2.174)$$

We are here free to choose $\beta = \gamma \gg 1$ by which

$$\rho_s \cong \rho_m = \varepsilon \quad (2.175)$$

with good approximation. The lower limits of the integrals (2.168) and (2.169) then decrease linearly with ε and the radius r_0 . This forms a “similar” set of geometrical configurations which thus have a shape being independent of ρ_m , ρ_s , and ε in the range of small ε .

The ratio of expressions (2.172) and (2.173) finally yields an effective photon diameter

$$2r_0 = \frac{\varepsilon \lambda_0}{\pi(\cos \alpha)} \quad (2.176)$$

It should be observed that the wave packet diameters of Eqs.(2.176) and (2.166) both become independent of the particular values of the parameters γ and β . As compared to the relatively large photon diameter (2.166) obtained for a convergent generating function, the diameter (2.176) based on a divergent such function can shrink to very small dimensions. This is the case even when $\cos \alpha \ll 1$ provided that $\varepsilon \ll \cos \alpha$. Then the photon model becomes strongly “needle-shaped” in its transverse directions.

4. *Features of Present Photon Model*

The possible existence of a nonzero photon rest mass was first called attention to by Einstein[52], Bass and Schrödinger[53], and de Broglie and Vigier[54]. It includes such fundamental points as its relation to the Michelson-Morley experiment, and its so far undetermined absolute value.

The phase and group velocities of the present non-dispersive wave packets become slightly smaller than the velocity of light, as expressed by condition (2.131) and Eq.(2.121). Thereby the velocity constant c can be regarded as an asymptotic limit at infinite photon energy. A small deviation from this limit still permits the theory to be compatible with the Michelson-Morley experiments. With condition (2.131) and Eq.(2.121) the deviation of the phase and group velocity from c becomes

$$v/c \cong 1 - \frac{1}{2}(\cos \alpha)^2 \quad (2.177)$$

For $\cos \alpha \leq 10^{-4}$ corresponding to a photon rest mass $m_0 < 0.74 \times 10^{-39}$ kg, a change in the eight decimal of the recorded velocity would hardly become detectable.

The Lorentz invariance is satisfied even if there is an axial velocity of propagation being slightly smaller than c . This is due to the velocity vector $\mathbf{C}$ which forms helical orbits, also having a small component which circulates around the axis.

The Poynting vector (2.158) is also found from Eqs.(2.134)–(2.139), combined with the conditions $\gamma \gg 1$ and (2.131), to become parallel with the velocity vector $\mathbf{C}$ in the first approximation.

The previous analysis shows that the physics in presence even of a very small rest mass becomes fundamentally different from that being based on a rest mass which is exactly equal to zero. In the latter case we are back from an EMS mode to a conventional axisymmetric EM mode with its divergent and physically unacceptable properties. Since the results of the EMS mode hold only for a nonzero rest mass, the quantum conditions (2.153) for the total energy and (2.161) for the angular momentum become satisfied by a whole class of small values of $\cos \alpha$ and m_0 .

As being pointed out by de Broglie and Vigier, such an indeterminateness of the photon rest mass appears at a first sight to be a serious objection to the underlying theory. The problem is that the derivations depend simply on the existence of the nonzero rest mass, but not on its magnitude. To this concept de Broglie and Vigier add other analogous examples that have been considered in theoretical physics. Thus, the uncertainty in the absolute value of the nonzero photon rest mass does not necessarily imply that the corresponding theory is questionable. On the contrary, as has been seen from the present analysis, such an unspecified rest mass rather becomes a strength of the theory. It namely makes variable ranges possible of the effective diameters for both axisymmetric photon models and corresponding light beams.

As shown by Eq.(2.161) a nonzero angular momentum only becomes possible when there is a nonzero rest mass (2.155) with an associated factor $\cos \alpha$. In its turn, this factor is related to a nonzero axial magnetic field component of the helical field configuration. For a vanishing rest mass we would thus be back to the conventional case where there is no axial magnetic field.

Turning to the effective photon diameter of these wave packet models, we first notice that a convergent generating function leads to a diameter (2.166) which in some cases becomes

rather limited, but still does not become small enough to match atomic dimensions. As an example, an average wave length $\lambda_0 = 3 \times 10^{-7}\text{m}$ and a factor $\cos \alpha = 10^{-4}$ which makes the velocity v deviate from c by 5×10^{-9} only, results in a diameter $2\hat{r} \cong 10^{-3}\text{m}$.

A divergent generating function can on the other hand result in a very small photon diameter (2.176) for a sufficiently small ε , then having the size of atomic dimensions such as the Bohr radius. This becomes reconcilable with the ability of the photon to knock out an electron from an atom in the photoelectric effect. This is consistent with the proposed “needle radiation” of energy quanta with a directed momentum as obtained by Einstein[55].

The obtained wave packet solutions are in some respects similar to the earlier wave-particle duality by de Broglie. Thus the total energy $h\nu = mc^2$ in the laboratory frame could be regarded to consist of the fraction $(m - m_0)c^2$ of a “free” pilot wave propagating along the axis, plus the fraction m_0c^2 of a “bound” particle-like state of radiation which circulates around the same axis. The rest mass then merely represents an integrating part of the total energy. Such a subdivision into a particle and an associated pilot wave is therefore not necessary in the present case where the wave packet behaves as an entity, having both particle and wave properties at the same time.

In the two-slit experiments[6] the present needle-shaped wave packet model has an advantage over a plane wave model. The particle feature in the form of a very narrow needle-shaped transverse diameter makes the wave packet model reconcilable with the observed dot-shaped marks. At the same time the wave nature of the axisymmetric packet makes it possible for interference phenomena to occur, in the same way as for a plane wave. This dualism and ability of interference becomes particularly obvious in the zero line width limit $1/k_0 z_0 \longrightarrow 0$ where the spectral distribution of Eq.(2.133) reduces to that of the elementary normal mode of Section E.1.

The analysis performed so far may debouch into the idea that there exist different quantum states (modes) of the photon representing different solutions of the same basic equations. In a way this would become analogous to but not identical with the Copenhagen interpretation. Almost instantaneous transitions between these states also become imaginable. A similar situation has already been observed for neutrino oscillations which are associated with a nonzero neutrino rest mass. Consequently, it is here proposed that analogous “photon oscillations” can exist, by which rapid transitions between the various plane and axisymmetric photon modes take place, under the constraints of total energy and an-

gular momentum conservation. In this way the photon could behave differently in different physical situations. Such a preliminary proposal has, however, to be further investigated.

The photon is the field quantum being responsible for the electromagnetic interaction. Likewise the weak field interaction acquires quanta, in the form of the W^+ , W^- , and Z^0 bosons[3]. This raises the question whether an analogous Proca-type equation applied to the weak field case could result in axisymmetric solutions being similar to those deduced here for the electromagnetic field. This may provide the weak-field bosons with a nonzero rest mass, thereby arriving at a possible alternative to the Higgs particle concept.

Finally it should be observed that the results of Section E.3 for an individual radially polarized photon will also apply to a corresponding dense beam of N photons per unit length and circular cross-section, provided that the individual photon fields overlap each other. Thereby the right-hand members of Eqs.(2.151) and (2.161) for the total mass and angular momentum will include the same factor N . As a result, Eqs.(2.166) and (2.176) then apply also to the beam diameters.

F. Summary of Obtained Main Results in Section II

The main results which are specific to the present theoretical approach can now be summarized. These are expected to contribute to an increasing understanding of a number of fundamental physical phenomena, and they also predict new features of the electromagnetic field to exist. It should be noticed here that a recently presented theory by Tauber[56] also has some parts in common with these features.

Maxwell's equations in the vacuum have been used as a guideline and basis in the development of quantum electrodynamics (QED), which is therefore expected to become subject to the typical shortcomings of conventional electromagnetic theory. The present revised electromagnetics, with its nonzero electric field divergence, provides a way of eliminating these shortcomings, in a first step of extended quantum electrodynamics ("EQED").

The present theory leads to some results of a general character, of which the following should be mentioned:

- The extended equations make it possible for electromagnetic steady states to exist in the vacuum.

- New types of wave modes arise, such as a longitudinal purely electric space-charge (S) wave, and a nontransverse electromagnetic space-charge (EMS) wave.

The electrically charged steady states result in a model of charged leptons such as the electron:

- For a particle-shaped configuration to possess a nonzero net electric charge, its characteristic radius has to shrink to that of a point-charge-like geometry, in agreement with experiments.
- Despite of the success of the conventional renormalization procedure, a more satisfactory way from the physical point of view is needed in respect to the infinite self-energy problem of a point charge. This problem is removed by the present theory which provides a more surveyable alternative, by compensating the divergence of the generating function through a shrinking characteristic radius and thereby leading to finite integrated field quantities.
- The Lorentz invariance of the electron radius is formally satisfied in the present theory, by allowing the same radius to shrink to that of a point charge. The obtained solutions can on the other hand also be applied to the physically relevant case of a very small but nonzero radius of a configuration having an internal structure.
- A steady equilibrium of the average radial forces can under certain conditions be established, such as to prevent the electron from “exploding” under the action of its electric self-charge. This becomes possible for the point-charge-like model, at parameter values of the plateau region which correspond to $q^* \cong 0.99$. The remaining degrees of freedom being available in the parameter ranges of the plateau have then been used up by the condition of a radially balanced equilibrium.
- That there is only a small deviation of q^* from unity can, in itself, be interpreted as an experimental support of the present theory. This is particularly the case as the deduced result has been obtained from two independent aspects, namely the minimization of the charge by a variational analysis, and the determination of the charge from the requirement of a radial balance of forces.

- The deduced charge thus deviates only by about one percent from the measured elementary charge. To remove this small deviation, some additional quantum mechanical corrections have been proposed. Provided that such corrections become relevant, the elementary charge would no longer remain as an independent constant of nature, but can be derived from the present theory in terms of the velocity of light, Planck's constant, and the permittivity of vacuum.

Conventional theory includes a vanishing electric field divergence, and it has no electric and magnetic field components in the direction of propagation, i.e. in the case of plane as well as of axisymmetric and screw-shaped wave modes [11]. The Poynting vector then possesses a component only in the direction of propagation, and there is no angular momentum (spin) with respect to the same direction. This feature also prevails in a fully quantized analysis. The electromagnetic field strengths appearing in the Poynting vector can then be expressed in terms of corresponding quantum mechanical plane wave expressions. The deduced axisymmetric wave modes lead on the other hand to models of the individual photon, in the form of EMS wave-packets with a narrow line width and having the following features:

- For the photon to possess a nonzero angular momentum (spin) in the capacity of a boson particle, the axisymmetric field geometry has to become helical, as well as the Poynting vector.
- This helical structure is associated with a nonzero rest mass, and it has field components also in the longitudinal (axial) direction of propagation.
- The nonzero rest mass can become small enough not to get into conflict with experiments of the Michelson-Morley type, thereby still preserving the conditions for a helical field structure.
- There are wave-packet solutions with a very small characteristic transverse radius. These have the form of “needle radiation” which becomes a necessary concept for the explanation of the photoelectric effect where a photon interacts with a single electron in an atom, and of the dot-shaped marks observed at a screen in double-slit experiments.
- The axisymmetric wave-packet models behave as one single entity, having particle and wave properties at the same time.

III. Unified Theory of Bivacuum, Particles Duality, Fields & Time

by A. Kaivarainen

III. A. The new concept of Bivacuum

A.1. Properties of Bivacuum dipoles - Bivacuum fermions and Bivacuum bosons

Our Unified Theory (UT) is a result of long-term efforts for unification of vacuum, matter and fields from few ground postulates [21-23; 37, 57]. The Bivacuum concept is a result of new interpretation and development of Dirac theory [16], pointing to equal probability of positive and negative energy in Nature.

The Bivacuum is introduced, as a dynamic superfluid matrix of the Universe, composed from non-mixing *subquantum particles* of opposite polarization, separated by an energy gap. The hypothetical *microscopic* subquantum particles and antiparticles have a dimensions less than the Plank length (10^{-33} cm), zero mass and charge. They form the infinite number of *mesoscopic* paired vortices - Bivacuum dipoles of three generations with Compton radii, corresponding to electrons (e), muons (μ) and tauons (τ). Only such *mesoscopic* collective excitations of subquantum particles in form of pairs of rotating fast *torus and antitorus* are quantized. In turn, these Bivacuum 'molecules' compose the *macroscopic* superfluid ideal liquid, representing the infinitive Bivacuum matrix.

Each of two strongly correlated 'donuts' of Bivacuum dipoles acquire the opposite mass charge and magnetic moments, compensating each other in the absence of symmetry shift between them. The latter condition is valid only for symmetric *primordial* Bivacuum, where the influence of matter and fields on Bivacuum is negligible. The sub-elementary fermion and antifermion origination is a result of the Bivacuum dipole symmetry shift toward the torus or antitorus, correspondingly. The correlation between paired vortical structures in a liquid medium was theoretically proved by Kiehn [58].

The infinite number of paired vortical structures: [torus ($\mathbf{V}^+$) + antitorus ($\mathbf{V}^-$)] with the in-phase clockwise or anticlockwise rotation are named Bivacuum fermions ($\mathbf{BVF}^\uparrow = \mathbf{V}^+ \uparrow\uparrow \mathbf{V}^-$)^{*i*} and Bivacuum antifermions ($\mathbf{BVF}^\downarrow = \mathbf{V}^+ \downarrow\downarrow \mathbf{V}^-$)^{*i*}, correspondingly. Their intermediate - transition states are named Bivacuum bosons of two possible polarizations: ($\mathbf{BVB}^+ = \mathbf{V}^+ \uparrow\downarrow \mathbf{V}^-$)^{*i*} and ($\mathbf{BVB}^- = \mathbf{V}^+ \downarrow\uparrow \mathbf{V}^-$)^{*i*} The *positive and negative energies*

of torus and antitorus ($\pm \mathbf{E}_{\mathbf{V}\pm}$) of three lepton generations ($i = e, \mu, \tau$), interrelated with their radiuses ($\mathbf{L}_{\mathbf{V}\pm}^n$), are quantized as quantum harmonic oscillators of opposite energies:

$$[\mathbf{E}_{\mathbf{V}\pm}^n = \pm \mathbf{m}_0 \mathbf{c}^2 (\frac{1}{2} + \mathbf{n}) = \pm \hbar \omega_0 (\frac{1}{2} + \mathbf{n})]^i \quad \mathbf{n} = 0, 1, 2, 3...^i \quad (3.1)$$

$$or : \left[\mathbf{E}_{\mathbf{V}\pm}^n = \frac{\pm \hbar \mathbf{c}}{\mathbf{L}_{\mathbf{V}\pm}^n} \right]^i \quad (3.1a)$$

$$where : \left[\mathbf{L}_{\mathbf{V}\pm}^n = \frac{\pm \hbar}{\pm \mathbf{m}_0 \mathbf{c} (\frac{1}{2} + \mathbf{n})} = \frac{\mathbf{L}_0}{\frac{1}{2} + \mathbf{n}} \right]$$

where: $[\mathbf{L}_0 = \hbar / \mathbf{m}_0 \mathbf{c}]^{e, \mu, \tau}$ is a Compton radii of the electron of corresponding lepton generation ($i = e, \mu, \tau$) and $\mathbf{L}_0^e \gg \mathbf{L}_0^\mu > \mathbf{L}_0^\tau$. The Bivacuum fermions $(\mathbf{BVF}^\dagger)^{\mu, \tau}$ with smaller Compton radiuses can be located inside the bigger ones $(\mathbf{BVF}^\dagger)^e$.

The absolute values of increments of torus and antitorus energies ($\Delta \mathbf{E}_{\mathbf{V}\pm}^i$), interrelated with increments of their radii ($\Delta \mathbf{L}_{\mathbf{V}\pm}^i$) in primordial Bivacuum (i.e. in the absence of matter and field influence), resulting from in-phase symmetric fluctuations are equal:

$$\Delta \mathbf{E}_{\mathbf{V}\pm}^i = - \frac{\hbar \mathbf{c}}{(\mathbf{L}_{\mathbf{V}\pm}^i)^2} \Delta \mathbf{L}_{\mathbf{V}\pm}^i = - \mathbf{E}_{\mathbf{V}\pm}^i \frac{\Delta \mathbf{L}_{\mathbf{V}\pm}^i}{\mathbf{L}_{\mathbf{V}\pm}^i} \quad or : \quad (3.2)$$

$$- \Delta \mathbf{L}_{\mathbf{V}\pm}^i = \frac{\pi (\mathbf{L}_{\mathbf{V}\pm}^i)^2}{\pi \hbar \mathbf{c}} \Delta \mathbf{E}_{\mathbf{V}\pm}^i = \frac{\mathbf{S}_{\mathbf{BVF}\pm}^i}{2 \hbar \mathbf{c}} \Delta \mathbf{E}_{\mathbf{V}\pm}^i = \mathbf{L}_{\mathbf{V}\pm}^i \frac{\Delta \mathbf{E}_{\mathbf{V}\pm}^i}{\mathbf{E}_{\mathbf{V}\pm}^i} \quad (3.2a)$$

where: $\mathbf{S}_{\mathbf{BVF}\pm}^i = \pi (\mathbf{L}_{\mathbf{V}\pm}^i)^2$ is a square of the cross-section of torus and antitorus, forming Bivacuum fermions $(\mathbf{BVF}^\dagger)$ and Bivacuum bosons $(\mathbf{BVB}^\pm)$.

The virtual *mass*, *charge* and *magnetic moments* of torus and antitorus of $\mathbf{BVF}^\dagger$ and $\mathbf{BVB}^\pm$ are opposite and in symmetric *primordial* Bivacuum compensate each other in their basic ($\mathbf{n} = 0$) and excited ($\mathbf{n} = 1, 2, 3...$) states.

The Bivacuum 'atoms': $\mathbf{BVF}^\dagger = [\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-]^i$ and $\mathbf{BVB}^\pm = [\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-]^i$ represent dipoles of three different poles - the mass ($\mathbf{m}_V^+ = |\mathbf{m}_V^-| = \mathbf{m}_0$)ⁱ, electric (e_+ and e_-) and magnetic (μ_+ and μ_-) dipoles.

The torus and antitorus $(\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-)^i$ of Bivacuum fermions of opposite spins $\mathbf{BVF}^\dagger$ and $\mathbf{BVF}^\dagger$ are both rotating in the same direction: clockwise or anticlockwise. This determines the positive and negative spins ($\mathbf{S} = \pm 1/2 \hbar$) of Bivacuum fermions. Their opposite spins may compensate each other, forming virtual Cooper pairs: $[\mathbf{BVF}^\dagger \propto \mathbf{BVF}^\dagger]$ with neutral boson properties. The rotation of adjacent $\mathbf{BVF}^\dagger$ and $\mathbf{BVF}^\dagger$ in Cooper pairs is *side-by-*

side in opposite directions, providing zero resulting spin of such pairs and ability to virtual Bose condensation. The torus and antitorus of Bivacuum bosons $\mathbf{BVB}^\pm = [\mathbf{V}^+\uparrow\downarrow\mathbf{V}^-]^i$ with spin, equal to zero, are rotating in opposite directions.

The *energy gap* between the torus and antitorus of symmetric $(\mathbf{BVF}^\uparrow)^i$ or $(\mathbf{BVB}^\pm)^i$ is:

$$[\mathbf{A}_{BVF} = \mathbf{E}_{\mathbf{V}^+} - (-\mathbf{E}_{\mathbf{V}^-}) = \hbar\omega_0(1 + 2\mathbf{n})]^i = \mathbf{m}_0^i \mathbf{c}^2(1 + 2\mathbf{n}) = \frac{\hbar \mathbf{c}}{[\mathbf{d}_{\mathbf{V}^+\uparrow\mathbf{V}^-}]_n^i} \quad (3.3)$$

where the characteristic distance between torus $(\mathbf{V}^+)^i$ and antitorus $(\mathbf{V}^-)^i$ of Bivacuum dipoles (*gap dimension*) is a quantized parameter:

$$[\mathbf{d}_{\mathbf{V}^+\uparrow\mathbf{V}^-}]_n^i = \frac{\hbar}{\mathbf{m}_0^i \mathbf{c}(1 + 2\mathbf{n})} \quad (3.4)$$

From (3.2) and (3.2a) we can see, that at $\mathbf{n} \rightarrow \mathbf{0}$, the energy gap $\mathbf{A}_{BVF}^i$ is decreasing till $\hbar\omega_0 = \mathbf{m}_0^i \mathbf{c}^2$ and the spatial gap dimension $[\mathbf{d}_{\mathbf{V}^+\uparrow\mathbf{V}^-}]_n^i$ is increasing up to the Compton length $\lambda_0^i = \hbar/\mathbf{m}_0^i \mathbf{c}$. On the contrary, the infinitive symmetric excitation of torus and antitorus is followed by tending the spatial gap between them to zero: $[\mathbf{d}_{\mathbf{V}^+\uparrow\mathbf{V}^-}]_n^i \rightarrow 0$ at $\mathbf{n} \rightarrow \infty$. This means that the quantization of space and energy of Bivacuum are interrelated and discreet.

The Planck aether hypothesis of Winterberg [59], like Quon vacuum model of Aspden [24] has some similarity with our Bivacuum model. For example, Winterberg supposed that space is filled by an equal number of positive and negative Planck mass particles, each occupied the Planck volume and forming superfluid medium. By the analogy with conventional theory of superfluid helium theory, Winterberg conjectured the phonon-roton like spectrum of Planck aether. The rotons are considered as a small vortex rings of Planck radius. However, this model is too 'rigid'. It do not take into account the relativistic effects and can not explain the mass and charge origination of elementary particles, their duality, fields and a lot of other important phenomena, following from Bivacuum model, as it will be demonstrated in this section III of our paper.

A.2 The basic (carrying) Virtual Pressure Waves (VPW $^\pm$) and Virtual spin waves (VirSW $^{\pm 1/2}$) of Bivacuum

The emission and absorption of Virtual clouds $(\mathbf{VC}_{j,k}^+)^i$ and anticlouds $(\mathbf{VC}_{j,k}^-)^i$ in primordial Bivacuum, i.e. in the absence of matter and fields or where their influence on

symmetry of Bivacuum is negligible, are the result of correlated transitions between different excitation states (j, k) of torus $(\mathbf{V}_{j,k}^+)^i$ and antitoruses $(\mathbf{V}_{j,k}^-)^i$, forming symmetric $[\mathbf{BVF}^\dagger]^i$ and $[\mathbf{BVB}^\pm]^i$, corresponding to three lepton generations ($i = e, \mu, \tau$):

$$(\mathbf{VC}_{j,k}^+)^i \equiv [\mathbf{V}_j^+ - \mathbf{V}_k^+]^i - \text{virtual cloud} \quad (3.5)$$

$$(\mathbf{VC}_{j,k}^-)^i \equiv [\mathbf{V}_j^- - \mathbf{V}_k^-]^i - \text{virtual anticloud} \quad (3.5a)$$

where: $j > k$ are the integer quantum numbers of torus and antitorus excitation states.

The virtual clouds: $(\mathbf{VC}_{j,k}^+)^i$ and $(\mathbf{VC}_{j,k}^-)^i$ exist in form of collective excitation of *subquantum* particles and antiparticles of opposite energies, correspondingly. They can be considered as 'drops' of virtual Bose condensation of subquantum particles of positive and negative energy.

The process of [*emission* $\rightleftharpoons$ *absorption*] of virtual clouds should be accompanied by oscillation of *virtual pressure* $(\mathbf{VirP}^\pm)$ and *excitation of positive and negative virtual pressure waves* $(\mathbf{VPW}^+ \text{ and } \mathbf{VPW}^-)_{j,k}$. In primordial Bivacuum the virtual pressure waves of opposite energies totally compensate each other. However, in asymmetric secondary Bivacuum, in presence of matter and fields, the total compensation is absent and the resulting virtual pressure is nonzero [22, 23]: $(\Delta \mathbf{VirP}^\pm = |\mathbf{VirP}^+| - |\mathbf{VirP}^-|) > 0$.

In accordance with our model of Bivacuum, virtual particles and antiparticles represent the asymmetric Bivacuum dipoles $(\mathbf{BVF}^\dagger)^{as}$ and $(\mathbf{BVB}^\pm)^{as}$ of three electron generations ($i = e, \mu, \tau$) in unstable state, not corresponding to Golden mean conditions (see subsection III.C). Virtual particles and antiparticles are the result of correlated and opposite Bivacuum dipole symmetry fluctuations. Virtual particles, like the real sub-elementary particles, may exist in Corpuscular and Wave phases (see subsection III.E). The Corpuscular [C]- phase, represents strongly correlated pairs of asymmetric torus (V^+) and antitorus (V^-) of two different by absolute values energies. The Wave [W]- phase, results from quantum beats between these states, which are accompanied by emission or absorption of Cumulative Virtual Cloud ($\mathbf{CVC}^+$ or $\mathbf{CVC}^-$), formed by subquantum particles.

Virtual particles have a mass, charge, spin, etc., but they differs from real sub-elementary ones by their lower stability (short life-time) and inability for fusion to stable triplets (see section 3). They are a singlets or very unstable triplets or other clusters of Bivacuum dipoles $(\mathbf{BVF}^\dagger)^{as}$ in contrast to real fermions-triplets.

For Virtual Clouds $(\mathbf{VC}^\pm)$ and virtual pressure waves $(\mathbf{VPW}^\pm)$ excited by them, the

relativistic mechanics is not valid. *Consequently, the causality principle also does not work in a system (interference pattern) of $\mathbf{VPW}^\pm$.*

The quantized energies of positive and negative $\mathbf{VPW}^+$ and $\mathbf{VPW}^-$ and corresponding virtual clouds and anticlouds, emitted $\rightleftharpoons$ absorbed by $(\mathbf{BVF})^i$ and $(\mathbf{BVB})^i$, as a result of their transitions between $\mathbf{j}$ and $\mathbf{k}$ states can be presented as:

$$\mathbf{E}_{\mathbf{VPW}_{j,k}^+}^i = \hbar\omega_0^i(\mathbf{j} - \mathbf{k}) = \mathbf{m}_0^i \mathbf{c}^2(\mathbf{j} - \mathbf{k}) \quad (3.6)$$

$$\mathbf{E}_{\mathbf{VPW}_{j,k}^-}^i = -\hbar\omega_0^i(\mathbf{j} - \mathbf{k}) = -\mathbf{m}_0^i \mathbf{c}^2(\mathbf{j} - \mathbf{k}) \quad (3.6a)$$

The quantized fundamental Compton frequency of $\mathbf{VPW}^\pm$:

$$\mathbf{q}\omega_0^i = \mathbf{q}\mathbf{m}_0^i \mathbf{c}^2 / \hbar \quad (3.7)$$

where: $\mathbf{q} = (\mathbf{j} - \mathbf{k}) = 1, 2, 3..$ is the quantization number of $\mathbf{VPW}_{j,k}^\pm$ energy.

In symmetric primordial Bivacuum the total compensation of positive and negative Virtual Pressure Waves is possible:

$$\mathbf{q}\mathbf{E}_{\mathbf{VPW}_{j,k}^+}^i = -\mathbf{q}\mathbf{E}_{\mathbf{VPW}_{j,k}^-}^i = \mathbf{q}\hbar\omega_0^i \quad (3.8)$$

The density oscillation of $\mathbf{VC}_{j,k}^+$ and $\mathbf{VC}_{j,k}^-$ and virtual particles and antiparticles represent *positive and negative basic virtual pressure waves* ($\mathbf{VPW}_{j,k}^+$ and $\mathbf{VPW}_{j,k}^-$).

The correlated *virtual Cooper pairs* of adjacent Bivacuum fermions ($\mathbf{BVF}_{S=\pm 1/2}^\uparrow$), rotating in opposite direction with resulting spin, equal to zero and Boson properties, can be presented as:

$$[\mathbf{BVF}_{S=1/2}^\uparrow \bowtie \mathbf{BVF}_{S=-1/2}^\downarrow]_{S=0} \equiv [(\mathbf{V}^+ \uparrow\uparrow \mathbf{V}^-) \bowtie (\mathbf{V}^+ \downarrow\downarrow \mathbf{V}^-)]_{S=0} \quad (3.9)$$

Such a pairs, as well as Bivacuum bosons ($\mathbf{BVB}^\pm$) in conditions of ideal equilibrium, like the *Goldstone bosons*, have zero mass and spin: $S = 0$. The virtual clouds ($\mathbf{VC}_{j,k}^\pm$), emitted and absorbed in a course of correlated transitions of $[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow]_{S=0}^{j,k}$ between (j) and (k) sublevels, excite the virtual pressure waves $\mathbf{VPW}^+$ and $\mathbf{VPW}^-$. They compensate the energy of each other totally in primordial Bivacuum and partly in *secondary Bivacuum* - in presence of matter and fields.

The nonlocal virtual spin waves ($\mathbf{VirSW}_{j,k}^{\pm 1/2}$), with properties of massless collective Nambu-Goldstone modes, represent oscillation of equilibrium of Bivacuum fermions with opposite spins, accompanied by origination of intermediate states - Bivacuum bosons ($\mathbf{BVB}^\pm$):

$$\mathbf{VirSW}_{j,k}^{\pm 1/2} \sim \left[\mathbf{BVF}^\uparrow (\mathbf{V}^+ \uparrow\uparrow \mathbf{V}^-) \rightleftharpoons \mathbf{BVB}^\pm (\mathbf{V}^+ \uparrow\downarrow \mathbf{V}^-) \rightleftharpoons \mathbf{BVF}^\downarrow (\mathbf{V}^+ \downarrow\downarrow \mathbf{V}^-) \right] \quad (3.10)$$

The $\mathbf{VirSW}_{j,k}^{+1/2}$ and $\mathbf{VirSW}_{j,k}^{-1/2}$ are excited by $(\mathbf{VC}_{j,k}^{\pm})_{S=1/2}^{\odot}$ and $(\mathbf{VC}_{j,k}^{\pm})_{S=-1/2}^{\odot}$ of opposite angular momentums: $S_{\pm 1/2} = \pm \frac{1}{2}\hbar = \pm \frac{1}{2}\mathbf{L}_0\mathbf{m}_0\mathbf{c}$ and frequency, equal to $\mathbf{VPW}_{j,k}^{\pm}$ (3.7):

$$\mathbf{q}\omega_{\mathbf{VirSW}_{j,k}^{\pm 1/2}}^i = \mathbf{q}\omega_{\mathbf{VPW}_{j,k}^{\pm}}^i = \mathbf{q}\mathbf{m}_0^i\mathbf{c}^2/\hbar = \mathbf{q}\omega_0^i \quad (3.10a)$$

The most probable basic virtual pressure waves $\mathbf{VPW}_0^{\pm}$ and virtual spin waves $\mathbf{VirSW}_0^{\pm 1/2}$ correspond to minimum quantum number $\mathbf{q} = (\mathbf{j} - \mathbf{k}) = \mathbf{1}$.

The $\mathbf{VirSW}_{j,k}^{\pm 1/2}$, like so-called torsion field, can serve as a carrier of the phase/spin (angular momentum) and information - *qubits*, but not the energy.

The Bivacuum bosons $(\mathbf{BVB}^{\pm})$, may have two polarizations $(\pm)$, determined by spin state of their actual torus $(\mathbf{V}^+)$:

$$\mathbf{BVB}^+ = (\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-), \quad \text{when } \mathbf{BVF}^{\uparrow} \rightarrow \mathbf{BVF}^{\downarrow} \quad (3.11)$$

$$\mathbf{BVB}^- = (\mathbf{V}^+ \downarrow \uparrow \mathbf{V}^-), \quad \text{when } \mathbf{BVF}^{\downarrow} \rightarrow \mathbf{BVF}^{\uparrow} \quad (3.11a)$$

The Bose-Einstein statistics of energy distribution, valid for system of weakly interacting bosons (ideal gas), do not work for Bivacuum due to strong coupling of pairs $[\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}]_{S=0}$ and $(\mathbf{BVB}^{\pm})$, forming virtual Bose condensate ($\mathbf{VirBC}$) with non-local properties. The Bivacuum nonlocal properties can be proved, using the Virial theorem [22,23].

A.3 Virtual Bose condensation (*VirBC*), as a base of Bivacuum nonlocality

It follows from our model of Bivacuum, that the infinite number of Cooper pairs of Bivacuum fermions $[\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}]_{S=0}^i$ and their intermediate states - Bivacuum bosons $(\mathbf{BVB}^{\pm})^i$, as elements of Bivacuum, have zero or very small (in presence of fields and matter) translational momentum: $\mathbf{p}_{\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}}^i = \mathbf{p}_{\mathbf{BVB}}^i \rightarrow 0$ and corresponding de Broglie wave length tending to infinity: $\lambda_{\mathbf{VirBC}}^i = \mathbf{h}/\mathbf{p}_{\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}, \mathbf{BVB}}^i \rightarrow \infty$. It leads to origination of 3D net of virtual adjacent pairs of virtual microtubules from Cooper pairs $[\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}]_{S=0}$, rotating in opposite direction and $(\mathbf{BVB}^{\pm})_{S=0}$, which may form single microtubules, with resulting angular momentum, equal to zero. These twin and single microtubules, termed Virtual Guides ($\mathbf{VirG}^{\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}}$ and $\mathbf{VirG}^{\mathbf{BVB}^{\pm}}$), represent a quasi one-dimensional Bose condensate with nonlocal properties close to that of 'wormholes' (see section 9). Their radiuses are determined by the Compton radiuses of the electrons, muons and tauons. Their

length is limited only by decoherence effects. In symmetric Bivacuum, unperturbed by matter and fields, the length of **VirG** may have the order of stars and galactics separation.

Nonlocality, as the independence of potential on the distance from its source in the volume or filaments of virtual or real Bose condensate, follows from application of the Virial theorem to systems of Cooper pairs of Bivacuum fermions $[\mathbf{BVF}^\dagger \bowtie \mathbf{BVF}^\dagger]_{S=0}$ and Bivacuum bosons $(\mathbf{BVB}^\pm)$ [22,23].

The Virial theorem in general form is correct not only for classical, but also for quantum systems. It relates the averaged kinetic $\overline{\mathbf{T}}_k(\tilde{\mathbf{v}}) = \sum_i \overline{\mathbf{m}_i \mathbf{v}_i^2}/2$ and potential $\overline{\mathbf{V}}(\mathbf{r})$ energies of particles, composing these systems:

$$2\overline{\mathbf{T}}_k(\tilde{\mathbf{v}}) = \sum_i \overline{\mathbf{m}_i \mathbf{v}_i^2} = \sum_i \tilde{\mathbf{r}}_i \partial \overline{\mathbf{V}} / \partial \tilde{\mathbf{r}}_i \quad (3.12)$$

If the potential energy $\overline{\mathbf{V}}(\mathbf{r})$ is a homogeneous $\varkappa$ – order function like:

$$\overline{\mathbf{V}}(\mathbf{r}) \sim \mathbf{r}^\varkappa, \text{ then } \mathbf{n} = \frac{2\overline{\mathbf{T}}_k}{\overline{\mathbf{V}}(\mathbf{r})} \quad (3.12a)$$

For example, for a harmonic oscillator, when $\overline{\mathbf{T}}_k = \overline{\mathbf{V}}$, we have $\varkappa = 2$. For Coulomb interaction: $\varkappa = -1$ and $\overline{\mathbf{T}} = -\overline{\mathbf{V}}/2$.

The important consequence of the Virial theorem is that, if the average kinetic energy and momentum ($\overline{\mathbf{p}}$) of particles in a certain volume of a Bose condensate (BC) tends to zero:

$$\overline{\mathbf{T}}_k = \overline{\mathbf{p}}^2/2\mathbf{m} \rightarrow 0 \quad (3.13)$$

the interaction between particles in the volume of BC, characterized by the radius: $\mathbf{L}_{BC} = (\hbar/\overline{\mathbf{p}}) \rightarrow 0$, becomes nonlocal, as independent on distance between them:

$$\begin{aligned} \overline{\mathbf{V}}(\mathbf{r}) \sim \mathbf{r}^\varkappa = 1 = \text{const} \\ \text{at } \varkappa = 2\overline{\mathbf{T}}_k/\overline{\mathbf{V}}(\mathbf{r}) = 0 \end{aligned} \quad (3.14)$$

Consequently, it is shown, that nonlocality, as independence of potential on the distance from potential source, is the inherent property of macroscopic Bose condensate. The individual particles (real, virtual or subquantum) in a state of Bose condensation are spatially indistinguishable due to the uncertainty principle. The Bivacuum dipoles $[\mathbf{BVF}^\dagger \bowtie \mathbf{BVF}^\dagger]_{S=0}$ and $(\mathbf{BVB}^\pm)_{S=0}$ due to quasi one-dimensional Bose condensation are

tending to self-assembly by 'head-to-tail' principle. They compose very long virtual microtubules - Virtual Guides with wormhole properties. The 3D net of these two kind of Virtual Guides ($\mathbf{VirG}^{\mathbf{BVF}^\dagger \bowtie \mathbf{BVF}^\downarrow}$ and $\mathbf{VirG}^{\mathbf{BVB}^\pm}$) represent the nonlocal fraction of superfluid Bivacuum.

III.B Basic postulates of Unified Theory and their consequences

There are three basic postulates in our theory, interrelated with each other [23]:

I. The absolute values of internal rotational kinetic energies of torus and antitorus are equal to each other and to the half of the rest mass energy of the electrons of corresponding lepton generation, independently on the external group velocity ($\mathbf{v}$), turning the symmetric Bivacuum fermions ($\mathbf{BVF}^\dagger$) to asymmetric ones:

$$[\mathbf{I}] : \left(\frac{1}{2} \mathbf{m}_V^+ (\mathbf{v}_{gr}^{in})^2 = \frac{1}{2} |-\mathbf{m}_V^-| (\mathbf{v}_{ph}^{in})^2 = \frac{1}{2} \mathbf{m}_0 \mathbf{c}^2 = \mathbf{const} \right)_i \quad (3.15)$$

where: $\mathbf{m}_V^+$ and $\mathbf{m}_V^-$ are the 'actual' - inertial and 'complementary' - inertialess masses of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$); the $\mathbf{v}_{gr}^{in}$ and $\mathbf{v}_{ph}^{in}$ are the *internal* angular group and phase velocities of subquantum particles and antiparticles, forming torus and antitorus, correspondingly. In symmetric conditions of *primordial* Bivacuum and its virtual dipoles, when the influence of matter and fields is absent: $\mathbf{v}_{gr}^{in} = \mathbf{v}_{ph}^{in} = \mathbf{c}$ and $\mathbf{m}_V^+ = \mathbf{m}_V^- = \mathbf{m}_0$.

It will be proved in section (G.1) of this section of paper, that above condition means the infinitive life-time of torus and antitorus of $\mathbf{BVF}^\dagger$ and $\mathbf{BVB}^\pm$.

II. The internal magnetic moments of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of asymmetric Bivacuum fermions $\mathbf{BVF}_{as}^\dagger = [\mathbf{V}^+ \uparrow \uparrow \mathbf{V}^-]$ and antifermions: $\mathbf{BVF}_{as}^\downarrow = [\mathbf{V}^+ \downarrow \downarrow \mathbf{V}^-]$, when $\mathbf{v}_{gr}^{in} \neq \mathbf{v}_{ph}^{in}$, $\mathbf{m}_V^+ \neq |-\mathbf{m}_V^-|$ and $|\mathbf{e}_+| \neq |\mathbf{e}_-|$, are equal to each other and to that of Bohr magneton: $\left[\mu_B = \mu_0 \equiv \frac{1}{2} |\mathbf{e}_0| \frac{\hbar}{\mathbf{m}_0 \mathbf{c}} \right]$, independently on their external translational velocity ($\mathbf{v} > \mathbf{0}$) and symmetry shift. In contrast to permanent magnetic moments of $\mathbf{V}^+$ and $\mathbf{V}^-$, their actual and complementary masses $\mathbf{m}_V^+$ and $|-\mathbf{m}_V^-|$, internal angular velocities ($\mathbf{v}_{gr}^{in}$ and $\mathbf{v}_{ph}^{in}$) and electric charges $|\mathbf{e}_+|$ and $|\mathbf{e}_-|$, are dependent on ($\mathbf{v}$), however, they compensate each other variations:

$$[\mathbf{II}] : \left(|\pm \mu_{\pm}| \equiv \frac{1}{2} |\mathbf{e}_{\pm}| \frac{|\pm \hbar|}{|\mathbf{m}_V^{\pm}| (\mathbf{v}_{gr}^{in})_{rot}} = |\pm \mu_{\pm}| \equiv \frac{1}{2} |-\mathbf{e}_{\pm}| \frac{|\pm \hbar|}{|-\mathbf{m}_V^{\mp}| (\mathbf{v}_{ph}^{in})_{rot}} = \right. \\ \left. = \mu_0 \equiv \frac{1}{2} |\mathbf{e}_0| \frac{\hbar}{\mathbf{m}_0 \mathbf{c}} = \mathbf{const} \right)_i \quad (3.16)$$

This postulate reflects the condition of the invariance of the spin value, with respect to the external velocity of Bivacuum fermions.

III. The equality of Coulomb interaction between torus and antitorus $\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-$ of primordial Bivacuum dipoles of all three generations $i = e, \mu, \tau$ (electrons, muons and taus), providing uniform electric energy density distribution in Bivacuum:

$$[\text{III}] : \mathbf{F}_0^i = \left(\frac{\mathbf{e}_0^2}{[\mathbf{d}_{\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-}]_n} \right)^e = \left(\frac{\mathbf{e}_0^2}{[\mathbf{d}_{\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-}]_n} \right)^\mu = \left(\frac{\mathbf{e}_0^2}{[\mathbf{d}_{\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-}]_n} \right)^\tau \quad (3.17)$$

where: $[\mathbf{d}_{\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-}]_n^i = \frac{\hbar}{\mathbf{m}_0^i \mathbf{c} (1+2n)}$ is the separation between torus and antitorus of Bivacuum three pole dipoles (3.4) at the same state of excitation (n). A similar condition is valid as well for opposite magnetic poles interaction; $|\mathbf{e}_+| |\mathbf{e}_-| = \mathbf{e}_0^2$.

The important consequences of postulate **III** are the following equalities:

$$(\mathbf{e}_0 \mathbf{m}_0)^e = (\mathbf{e}_0 \mathbf{m}_0)^\mu = (\mathbf{e}_0 \mathbf{m}_0)^\tau = \sqrt{|\mathbf{e}_+ \mathbf{e}_-| |\mathbf{m}_V^+ \mathbf{m}_V^-|} = \text{const} \quad (3.18)$$

It means that the toruses and antitoruses of symmetric Bivacuum dipoles of generations with bigger mass: $\mathbf{m}_0^\mu = 206,7 \mathbf{m}_0^e$; $\mathbf{m}_0^\tau = 3487,28 \mathbf{m}_0^e$ have correspondingly smaller charges:

$$\mathbf{e}_0^\mu = \mathbf{e}_0^e (\mathbf{m}_0^e / \mathbf{m}_0^\mu); \quad \mathbf{e}_0^\tau = \mathbf{e}_0^e (\mathbf{m}_0^e / \mathbf{m}_0^\tau) \quad (3.19)$$

As is shown in the next section, just these conditions provide *the same charge symmetry shift* of Bivacuum fermions of three generations ($i = e, \mu$) at the different mass symmetry shift between corresponding torus and antitorus, determined by Golden mean.

It follows from second postulate, that the resulting magnetic moment of sub-elementary fermion or antifermion ($\mu^\pm$), equal to the Bohr's magneton, is interrelated with the actual spin of Bivacuum fermion or antifermion as:

$$\mu^\pm = (|\pm \mu_+| |\pm \mu_-|)^{1/2} = \mu_B = \pm \frac{1}{2} \hbar \frac{\mathbf{e}_0}{\mathbf{m}_0 \mathbf{c}} = \mathbf{S} \frac{\mathbf{e}_0}{\mathbf{m}_0 \mathbf{c}} \quad (3.20)$$

where: $\mathbf{e}_0 / \mathbf{m}_0 \mathbf{c}$ is gyromagnetic ratio of Bivacuum fermion, equal to that of the electron.

One may see from (3.20), that the spin of the actual torus, equal to that of the resulting spin of Bivacuum fermion (symmetric or asymmetric), is:

$$\mathbf{S} = \pm \frac{1}{2} \hbar \quad (3.21)$$

Consequently, the permanent absolute value of spin of torus and antitorus is a consequence of 2nd postulate.

The dependence of the *actual inertial* mass ($\mathbf{m}_V^+$) of torus $\mathbf{V}^+$ of asymmetric Bivacuum fermions ($\mathbf{BVF}_{as}^\uparrow = \mathbf{V}^+ \uparrow \uparrow \mathbf{V}^-$) on the external translational group velocity ($\mathbf{v}$) follows relativistic mechanics:

$$\mathbf{m}_V^+ = \mathbf{m}_0 / \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} = \mathbf{m} \quad (3.22)$$

while the *complementary inertialess* mass ($\mathbf{m}_V^-$) of antitorus $\mathbf{V}^-$ has the reverse velocity dependence:

$$-\mathbf{m}_V^- = i^2 \mathbf{m}_V^- = -\mathbf{m}_0 \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \quad (3.23)$$

where $i^2 = -1$; $i = \sqrt{-1}$ and complementary mass in terms of mathematics is the imaginary parameter.

For Bivacuum antifermions $\mathbf{BVF}_{as}^\downarrow = \mathbf{V}^+ \downarrow \downarrow \mathbf{V}^-$ the relativistic dependences of positive and negative mass are opposite to those described by (3.22) and (3.23) and the notions of actual and complementary parameters change place.

The product of actual (inertial) and complementary (inertialess) mass is a constant, equal to the rest mass of particle squared and reflect the *mass compensation principle*. It means, that increasing of mass/energy of the torus is compensated by in-phase decreasing of absolute values of these parameters for antitorus and vice versa:

$$|\mathbf{m}_V^+| \quad |-\mathbf{m}_V^-| = \mathbf{m}_0^2 \quad (3.24)$$

Taking (3.24) and (3.15) into account, we get for the product of the *internal* group and phase velocities of positive and negative subquantum particles, forming torus and antitorus, correspondingly:

$$\mathbf{v}_{gr}^{in} \mathbf{v}_{ph}^{in} = \mathbf{c}^2 \quad (3.24a)$$

A similar symmetric relation is reflecting the *charge compensation principle*:

$$|\mathbf{e}_+| \quad |\mathbf{e}_-| = \mathbf{e}_0^2 \quad (3.24b)$$

For Bivacuum antifermions $\mathbf{BVF}_{as}^\downarrow = \mathbf{V}^+ \downarrow \downarrow \mathbf{V}^-$ the relativistic dependences of positive and negative charge, like the positive and negative masses of torus and antitorus are opposite to that of Bivacuum fermions $\mathbf{BVF}_{as}^\uparrow = \mathbf{V}^+ \uparrow \uparrow \mathbf{V}^-$ (see eqs.3.29 and 3.29a). The symmetry of Bivacuum bosons ($\mathbf{BVB}^\pm = \mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-$)^{*i*} of each electron's generation (*i* = *e*, *μ*, *τ*) can be

ideal and independent on external velocity, due to opposite relativistic effects of their torus and antitorus, compensating each other.

The product of actual (inertial) and complementary (inertialess) mass is a constant, equal to the rest mass of particle squared and reflect the *mass compensation principle*. It means, that increasing of mass/energy of the torus is compensated by in-phase decreasing of absolute values of these parameters for antitorus and vice versa.

The difference between absolute values of total actual and the complementary energies (3.22 and 3.23) is equal to doubled kinetic energy of asymmetric $\mathbf{BVF}_{as}^\dagger$:

$$|\mathbf{m}_V^+| \mathbf{c}^2 - |-\mathbf{m}_V^-| \mathbf{c}^2 = (\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2 = \mathbf{m}_V^+ \mathbf{v}^2 = 2\mathbf{T}_k = \frac{\mathbf{m}_0 \mathbf{v}^2}{\sqrt{1 - (\mathbf{v}/\mathbf{c})^2}} \quad (3.25)$$

The fundamental Einstein equation for total energy of particle can be reformed and extended, using eqs. 3.25 and 3.24:

$$\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{m} \mathbf{c}^2 = \mathbf{m}_V^- \mathbf{c}^2 + \mathbf{m}_V^+ \mathbf{v}^2 \quad (3.26)$$

$$or : \mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \frac{\mathbf{m}_0^2}{\mathbf{m}_V^+} \mathbf{c}^2 + \mathbf{m}_V^+ \mathbf{v}^2 \quad (3.26a)$$

$$or : \mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \mathbf{m}_0 \mathbf{c}^2 + 2\mathbf{T}_k \quad (3.26b)$$

The ratio of (3.23) to (3.22), taking into account (3.24), is:

$$\frac{|-\mathbf{m}_V^-|}{\mathbf{m}_V^+} = \frac{\mathbf{m}_0^2}{(\mathbf{m}_V^+)^2} = 1 - \left(\frac{\mathbf{v}}{\mathbf{c}}\right)^2 \quad (3.27)$$

It can easily be transformed to the important formula (3.25).

B.1 The relation between the external and internal parameters

of Bivacuum fermions & quantum roots of Golden mean.

The rest mass and charge origination

The important formula, unifying a lot of internal and external (translational-rotational) parameters of $\mathbf{BVF}_{as}^\dagger$, taking into account that the product of internal and external phase and group velocities is equal to light velocity squared:

$$\mathbf{v}_{ph}^{in} \mathbf{v}_{gr}^{in} = \mathbf{v}_{ph}^{ext} \mathbf{v}_{gr}^{ext} = \mathbf{c}^2 \quad (3.28)$$

can be derived from eqs. (3.15 - 3.27):

$$\left(\frac{\mathbf{m}_V^+}{\mathbf{m}_V^-}\right)^{1/2} = \frac{\mathbf{m}_V^+}{\mathbf{m}_0} = \frac{\mathbf{v}_{ph}^{in}}{\mathbf{v}_{gr}^{in}} = \left(\frac{\mathbf{c}}{\mathbf{v}_{gr}^{in}}\right)^2 = \quad (3.29)$$

$$= \frac{\mathbf{L}^-}{\mathbf{L}^+} = \frac{|\mathbf{e}_+|}{|\mathbf{e}_-|} = \left(\frac{\mathbf{e}_+}{\mathbf{e}_0}\right)^2 = \frac{1}{[1 - (\mathbf{v}^2/\mathbf{c}^2)^{ext}]^{1/2}} \quad (3.29a)$$

where:

$$\mathbf{L}_V^+ = \hbar/(\mathbf{m}_V^+ \mathbf{v}_{gr}^{in}) \quad and \quad \mathbf{L}_V^- = \hbar/(\mathbf{m}_V^- \mathbf{v}_{ph}^{in}) \quad (3.30)$$

$$\mathbf{L}_0 = (\mathbf{L}_V^+ \mathbf{L}_V^-)^{1/2} = \hbar/\mathbf{m}_0 \mathbf{c} - \text{Compton radius} \quad (3.30a)$$

are the radii of torus ($\mathbf{V}^+$), antitorus ($\mathbf{V}^-$) and the resulting radius of $\mathbf{BVF}_{as}^\dagger = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]$, equal to Compton radius, correspondingly.

The relativistic dependence of the actual charge $\mathbf{e}_+$ on velocity of Bivacuum dipole, following from (3.29a), is:

$$\mathbf{e}_+ = \frac{\mathbf{e}_0}{[1 - (\mathbf{v}^2/\mathbf{c}^2)^{ext}]^{1/4}} \quad (3.31)$$

The influence of relativistic dependence of *real* particles charge on the resulting charge and electric field density of Bivacuum, which is known to be electrically quasi neutral vacuum/bivacuum, is negligible for two reasons:

1. Densities of positive and negative real charges (i.e. particles and antiparticles) are very small and approximately equal, as it follows from Bo Lehnert approach (see Chapter II). This quasi-equilibrium of opposite charges is Lorentz invariant;
2. The remnant uncompensated by real antiparticles charges density at any velocities can be compensated totally by virtual antiparticles and asymmetric Bivacuum fermions (BVF) of opposite charges.

The ratio of the actual charge to the actual inertial mass, as it follows from (3.29 and 3.29a), has also the relativistic dependence:

$$\frac{\mathbf{e}_+}{\mathbf{m}_V^+} = \frac{\mathbf{e}_0}{\mathbf{m}_0} [1 - (\mathbf{v}^2/\mathbf{c}^2)^{ext}]^{1/4} \quad (3.32)$$

The decreasing of this ratio with velocity increasing is weaker, than what follows from the generally accepted statement, that charge has no relativistic dependence in contrast to the actual mass $\mathbf{m}_V^+$. The direct experimental study of relativistic dependence of this ratio on the external velocity ($\mathbf{v}$) may confirm the validity of our formula (3.32) and general approach.

From eqs. (3.25; 3.29 and 3.29a) we find for mass and charge symmetry shift:

$$\Delta \mathbf{m}_{\pm} = \mathbf{m}_V^+ - \mathbf{m}_V^- = \mathbf{m}_V^+ \left(\frac{\mathbf{v}}{\mathbf{c}} \right)^2 \quad (3.33)$$

$$\Delta \mathbf{e}_{\pm} = \mathbf{e}_+ - \mathbf{e}_- = \frac{\mathbf{e}_+^2}{\mathbf{e}_+ + \mathbf{e}_-} \left(\frac{\mathbf{v}}{\mathbf{c}} \right)^2 \quad (3.33a)$$

These mass and charge symmetry shifts determines the relativistic dependence of the *effective* mass and charge of the fermions. In direct experiments only the actual mass ($\mathbf{m}_V^+$) and charge ($\mathbf{e}_{\pm}$) can be registered. It means that the complementary mass ($\mathbf{m}_V^-$) and charge are *hidden* for observation.

The ratio of charge to mass symmetry shifts is:

$$\frac{\Delta \mathbf{e}_{\pm}}{\Delta \mathbf{m}_{\pm}} = \frac{\mathbf{e}_+^2}{\mathbf{m}_V^+ (\mathbf{e}_+ + \mathbf{e}_-)} \quad (3.34)$$

The mass symmetry shift can be expressed via the squared charges symmetry shift also in the following manner:

$$\Delta \mathbf{m}_{\pm} = \mathbf{m}_V^+ - \mathbf{m}_V^- = \mathbf{m}_V^+ \frac{\mathbf{e}_+^2 - \mathbf{e}_-^2}{\mathbf{e}_+^2} \quad (3.35)$$

or using (3.27) this formula turns to:

$$\frac{\mathbf{e}_+^2 - \mathbf{e}_-^2}{\mathbf{e}_+^2} = \frac{\mathbf{v}^2}{\mathbf{c}^2} \quad (3.36)$$

When the mass and charge symmetry shifts of Bivacuum dipoles are small and $|\mathbf{e}_+| + |\mathbf{e}_-| \simeq 2\mathbf{e}_+ \simeq 2\mathbf{e}_0$, we get from 3.33a for variation of charge shift:

$$\Delta \mathbf{e}_{\pm} = \mathbf{e}_+ - \mathbf{e}_- = \frac{1}{2} \mathbf{e}_0 \frac{\mathbf{v}^2}{\mathbf{c}^2} \quad (3.37)$$

The formula, unifying the *internal* and *external* group and phase velocities of asymmetric Bivacuum fermions ($\mathbf{BVF}_{as}^{\uparrow}$), derived from (3.29) and (3.29a), is:

$$\left(\frac{\mathbf{v}_{gr}^{in}}{\mathbf{c}} \right)^4 = 1 - \left(\frac{\mathbf{v}}{\mathbf{c}} \right)^2 \quad (3.38)$$

where: $(\mathbf{v}_{gr}^{ext}) \equiv \mathbf{v}$ is the external translational-rotational group velocity of $\mathbf{BVF}_{as}^{\uparrow}$.

At the conditions of "Hidden Harmony", meaning the equality of the internal and external rotational group and phase velocities of asymmetric Bivacuum fermions $\mathbf{BVF}_{as}^{\uparrow}$:

$$(\mathbf{v}_{gr}^{in})_{\mathbf{v}^+}^{rot} = (\mathbf{v}_{gr}^{ext})^{tr} \equiv \mathbf{v} \quad (3.39)$$

$$(\mathbf{v}_{ph}^{in})_{\mathbf{v}^-}^{rot} = (\mathbf{v}_{ph}^{ext})^{tr} \quad (3.39a)$$

and introducing the notation:

$$\left(\frac{\mathbf{v}_{gr}^{in}}{\mathbf{c}}\right)^2 = \left(\frac{\mathbf{v}}{\mathbf{c}}\right)^2 = \left(\frac{\mathbf{v}_{gr}^{in}}{\mathbf{v}_{ph}^{in}}\right) = \left(\frac{\mathbf{v}_{gr}^{ext}}{\mathbf{v}_{ph}^{ext}}\right) \equiv \phi \quad (3.40)$$

formula (3.38) turns to a simple quadratic equation:

$$\phi^2 + \phi - 1 = 0, \quad (3.41)$$

which has a few modes :

$$\phi = \frac{1}{\phi} - 1 \quad or : \quad \frac{\phi}{(1-\phi)^{1/2}} = 1 \quad (3.41a)$$

$$or : \quad \frac{1}{(1-\phi)^{1/2}} = \frac{1}{\phi} \quad (3.41b)$$

The solution of (3.41), is equal to **Golden mean**: $(\mathbf{v}/\mathbf{c})^2 = \phi = 0.618$. *It is remarkable, that the Golden Mean, which plays so important role on different Hierarchic levels of matter organization: from elementary particles to galactics and even in our perception of beauty (i.e. our mentality), has so deep physical roots, corresponding to Hidden Harmony conditions (3.39 and 3.39a). Our theory is the first one, elucidating these roots [21-23]. This important fact points, that we are on the right track.*

The overall shape of asymmetric $\left(\mathbf{BVF}_{as}^\uparrow = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]\right)^i$ is a *truncated cone* (Fig.3.1) with plane, parallel to the base with radiuses of torus (L^+) and antitorus (L^-), defined by eq. (3.30).

Using Golden Mean equation in the form (2.22b), we can see, that all the ratios (3.29 and 3.29a) at Golden Mean conditions turn to:

$$\left[\left(\frac{\mathbf{m}_V^+}{\mathbf{m}_V^-} \right)^{1/2} = \frac{\mathbf{m}_V^+}{\mathbf{m}_0} = \frac{\mathbf{v}_{ph}^{in}}{\mathbf{v}_{gr}^{in}} = \frac{\mathbf{L}^-}{\mathbf{L}^+} = \frac{|\mathbf{e}_+|}{|\mathbf{e}_-|} = \left(\frac{\mathbf{e}_+}{\mathbf{e}_0} \right)^2 \right]^\phi = \frac{1}{\phi} \quad (3.42)$$

where the actual (e_+) and complementary (e_-) charges and corresponding mass at GM conditions are:

$$\mathbf{e}_+^\phi = \mathbf{e}_0 / \phi^{1/2}; \quad \mathbf{e}_-^\phi = \mathbf{e}_0 \phi^{1/2} \quad (3.43)$$

$$(\mathbf{m}_V^+)^\phi = \mathbf{m}_0 / \phi; \quad (\mathbf{m}_V^-)^\phi = \mathbf{m}_0 \phi \quad (3.44)$$

using (3.44 and 3.41a) it is easy to see, that the difference between the actual and complementary mass at GM conditions is equal to the rest mass:

$$[|\Delta \mathbf{m}_V|^\phi = \mathbf{m}_V^+ - \mathbf{m}_V^- = \mathbf{m}_0(1/\phi - \phi) = \mathbf{m}_0]^{e,\mu,\tau} \quad (3.45)$$

This is an important result, pointing that just a symmetry shift, determined by the Golden mean conditions, is responsible for origination of the rest mass of sub-elementary particles of each of three generation ($i = e, \mu, \tau$).

The same is true for charge origination. The GM difference between actual and complementary charges, using relation $\phi = (1/\phi - 1)$, determines corresponding minimum charge of sub-elementary fermions or antifermions (at $\mathbf{v}_{tr}^{ext} \rightarrow \mathbf{0}$):

$$\phi^{3/2}\mathbf{e}_0 = |\Delta\mathbf{e}_{\pm}|^{\phi} = |\mathbf{e}_+ - \mathbf{e}_-|^{\phi} \equiv |\mathbf{e}|^{\phi} \quad (3.46)$$

$$\text{where: } (|\mathbf{e}_+| |\mathbf{e}_-|) = \mathbf{e}_0^2 \quad (3.46a)$$

The absolute values of charge symmetry shifts for electron, muon and tauon at GM conditions are the same. This result determines the equal absolute values of empirical rest charges of the electron, positron, proton and antiproton. However, the mass symmetry shifts at GM conditions, equal to the rest mass of electrons, muons and tauons are very different.

The ratio of charge to mass symmetry shifts at Golden mean (GM) conditions ($\mathbf{v}_{tr}^{ext} = 0$) is a permanent value for all three electron generations (e, μ, τ). The different values of their rest mass are taken into account by postulate III and its consequences of their rest mass and charge relations: $\mathbf{e}_0^{\mu} = \mathbf{e}_0^e(\mathbf{m}_0^e/\mathbf{m}_0^{\mu})$; $\mathbf{e}_0^{\tau} = \mathbf{e}_0^e(\mathbf{m}_0^e/\mathbf{m}_0^{\tau})$ (see 3.19) :

$$\left[\frac{|\Delta\mathbf{e}_{\pm}|^{\phi}}{|\Delta\mathbf{m}_V|^{\phi}} = \frac{|\mathbf{e}|^{\phi}}{\mathbf{m}_0^e} = \frac{|\mathbf{e}_+|^{\phi}\phi}{|\mathbf{m}_V^+|^{\phi}} = \frac{\mathbf{e}_0\phi^{3/2}}{\mathbf{m}_0} = \frac{\mathbf{e}_0\phi^{3/2}}{\mathbf{m}_0^{\mu,\tau}} \right]^{e,\mu,\tau} \quad (3.47)$$

where: $(\mathbf{m}_V^+)^{\phi} = \mathbf{m}_0/\phi$ is the actual mass of unpaired sub-elementary fermion in [C] phase at Golden mean conditions (see next section); $\mathbf{e}_0 \equiv \mathbf{e}_0^e$; $\mathbf{m}_0^e \equiv \mathbf{m}_0$.

Formula (3.47) can be considered as a background of permanent value of gyromagnetic ratio, equal to ratio of magnetic moment of particle to its angular momentum (spin). For the electron it is:

$$\Gamma = \frac{\mathbf{e}_0}{2\mathbf{m}_e\mathbf{c}} \quad (3.48)$$

When searching for elementary magnetic charges (g^- and g^+) being symmetric in respect to the electric ones (e^- and e^+) and named *monopoles*, the Dirac theory leads to the relation $\mathbf{g}\mathbf{e} = \frac{1}{2}\mathbf{n}\mathbf{h}\mathbf{c}$ between the magnetic monopole and the electric charge of the same signs, where $\mathbf{n} = 1, 2, 3$ is an integer. It follows from this definition that the minimal magnetic charge (at $\mathbf{n} = 1$) is as big as $g = 67.7e$. The mass of the monopole should then be huge, about 10^{16} GeV. All numerous attempts to reveal such particles experimentally have failed. But

the postulate II of our theory (eq.3.16) explains the absence of magnetic monopoles, through the absence of a symmetry shift between the internal opposite actual quantities $|\mu_V^+|$ and the complementary $|\mu_V^-|$ magnetic moments of the virtual torus and antitorus, which form Bivacuum fermions. This is in contrast to the mass and charge symmetry shifts.

The absence of magnetic monopole - spatially localized magnetic charge, is one of the important consequences of our model of elementary particles, as far:

$$\Delta\mu^\pm = \mu_V^+ - \mu_V^- = 0 \quad (3.49)$$


i.e. the magnetic moments of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) symmetry shift are always zero, independently on the external group velocity of elementary particles. This consequence of our theory is in line with a lot of unsuccessful attempts to reveal monopole experimentally.

III. C. Fusion of triplets of elementary particles from sub-elementary fermions and antifermions at Golden mean conditions

The fusion of asymmetric sub-elementary fermions and antifermions of μ and τ generations to triplets $\langle [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_{x,y} + \mathbf{F}_\downarrow^\pm \rangle_z^{e,p}$, corresponding to electrons/positrons, protons/antiprotons and neutrons/antineutrons becomes possible also at the Golden mean (GM) conditions (Fig.3.1). It is accompanied by energy release and electronic and hadronic e, h -gluons origination, equal in sum to the mass defect. It was demonstrated theoretically, that the vortical structures at certain conditions self-organizes into vortex crystals [60].

Model of the electron, as a triplet of rotating sub-elementary fermions:

$$< [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\uparrow}^-] + \mathbf{F}_{\downarrow}^- >$$



The total energy of each sub-elementary fermion:

$$\mathbf{E}_{tot} = mc^2 = \sqrt{1 - (\mathbf{v}/c)^2} (m_0 \omega_0^2 L^2)_{rot}^{in} + \left(\frac{h^2}{m \lambda_B^2} \right)_{tr}^{ext}$$

$$or : \quad \mathbf{E}_{tot} = \sqrt{1 - (\mathbf{v}/c)^2} \hbar \omega_0^{in} + \hbar \omega^{ext}; \quad \lambda_B = h/m\mathbf{v}_{tr}^{ext}$$

1. **Fig.3.1** Model of the electron, as a triplets $< [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\uparrow}^-] + \mathbf{F}_{\downarrow}^- >^e$, resulting from fusion of unpaired sub-elementary antifermion $\mathbf{F}_{\downarrow}^- >$ and rotating pair $[\mathbf{F}_{\downarrow}^+ \bowtie \mathbf{F}_{\uparrow}^-]$ of μ -generation. The velocity of rotation of unpaired $\mathbf{F}_{\downarrow}^- >$ around the same axis of common rotation axis of pair provide the similar mass and charge symmetry shifts: $|\Delta \mathbf{m}_V|^\phi = \mathbf{m}_0$ and $|\Delta \mathbf{e}^\pm|^\phi / \phi^{3/2} = \mathbf{e}_0$, as have the paired sub-elementary fermions $[\mathbf{F}_{\uparrow}^+$ and $\mathbf{F}_{\downarrow}^-]$. Three effective anchor $\left(\mathbf{BVF}^\dagger = [\mathbf{V}^+ \updownarrow \mathbf{V}^-] \right)_{anc}$ in the vicinity of the sub-elementary particle base, participate in recoil effects, accompanied $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation and modulation of basic Bivacuum pressure waves ($\mathbf{VPW}_0^\pm$). The relativistic mass

change of triplets is determined only by symmetry shift of the anchor $(\mathbf{BVF}^\dagger)_{anc}$ of the unpaired sub-elementary fermion $\mathbf{F}_\uparrow^\pm >$ (see section 3.5).

The *asymmetry of rotation velocity of torus and antitorus* of $(\mathbf{BVF}_{as}^\dagger = \mathbf{V}^+ \Downarrow \mathbf{V}^-)$, is a result of participation of pairs of $\mathbf{BVF}_{as}^\dagger$ of opposite spins $[\mathbf{BVF}_a^\dagger \bowtie \mathbf{BVF}_a^\downarrow]_{S=0}^i$ in Bivacuum vorticity. This motion can be described, as a rolling of pairs of $\mathbf{BVF}_{as}^\dagger = [\mathbf{V}^+ \Uparrow \mathbf{V}^-]$ and $\mathbf{BVF}_{as}^\downarrow = [\mathbf{V}^+ \Downarrow \mathbf{V}^-]$ with their *internal* radiuses:

$$\mathbf{L}_{\mathbf{BVF}^\dagger}^{in} = \hbar / |\mathbf{m}_V^+ + \mathbf{m}_V^-|_{\mathbf{BVF}^\dagger} \mathbf{c} \quad (3.50)$$

around the inside of a larger *external* circle with radius of vorticity:

$$\mathbf{L}_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow}^{ext} = \frac{\hbar}{|\mathbf{m}_V^+ - \mathbf{m}_V^-|_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow} \cdot \mathbf{c}} = \frac{\hbar \mathbf{c}}{\mathbf{m}_V^+ \mathbf{v}^2_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow}} \quad (3.50a)$$

The increasing of velocity of vorticity $\mathbf{v}_{vor}$ decreases both dimensions: $\mathbf{L}_{in}$ and $\mathbf{L}_{ext}$ till minimum vorticity radius, including pair of adjacent Bivacuum fermions $[\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_a^\downarrow]_{S=0}^i$ with shape of *two identical truncated cones* of the opposite orientation of planes with common rotation axis (see Fig.3.1). The corresponding asymmetry of torus $\mathbf{V}^+$ and $\mathbf{V}^-$ is responsible for resulting mass and charge of $\mathbf{BVF}_{as}^\dagger$. The trajectory of a fixed point on each of two $\mathbf{BVF}_{as}^\dagger$, participating in such dual rotation, is **hypocycloid**.

The ratio of internal and external radii: (3.1) and (3.1a) is equal to the ratio of potential $\mathbf{V}_p$ and kinetic $\mathbf{T}_k$ energy of the de Broglie wave, related with its phase $\mathbf{v}_{ph}$ and group $\mathbf{v}_{gr}$ velocities, as demonstrated in next section 3.4:

$$\frac{\mathbf{L}_{\mathbf{BVF}^\dagger}^{in}}{\mathbf{L}_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow}^{ext}} = \frac{|\mathbf{m}_V^+ + \mathbf{m}_V^-|_{\mathbf{BVF}^\dagger}}{|\mathbf{m}_V^+ - \mathbf{m}_V^-|_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow}} = \frac{\mathbf{V}_p}{\mathbf{T}_k} = 2 \frac{\mathbf{v}_{ph}}{\mathbf{v}_{gr}} - 1 \quad (3.51)$$

At Golden mean conditions, when $\mathbf{v} \rightarrow \phi^{1/2}$, $|\mathbf{m}_V^+ - \mathbf{m}_V^-|_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow} \rightarrow \mathbf{m}_0$; $(\mathbf{m}_V^+)^{\phi} = \mathbf{m}_0/\phi$ and $(\mathbf{m}_V^-)^{\phi} = \mathbf{m}_0\phi$, we get for internal (3.50) and external (3.50a) radiuses:

$$(\mathbf{L}_{\mathbf{BVF}^\dagger}^{in})^{\phi} = \frac{\hbar}{\mathbf{m}_0(1 + 2\phi)\mathbf{c}} = \frac{\mathbf{L}_0}{2.23} \quad (3.52)$$

$$\mathbf{L}_{\mathbf{BVF}_{as}^\dagger \bowtie \mathbf{BVF}_{as}^\downarrow}^{ext} = \frac{\hbar}{\mathbf{m}_0\mathbf{c}} = \mathbf{L}_0 \quad (3.52a)$$

The performed analysis can be considered, as a strong evidence, pointing to important role of Golden mean in the process of elementary particle fusion from sub-elementary constituents.

The model of a photon (boson), resulting from fusion (annihilation) of pairs of fermions triplets (electron + positron), are presented by Fig.3.2. Such a structure can originate also, as a result of six Bivacuum dipoles opposite symmetry shift, accompanied by the transitions of the excited atoms and molecules, i.e. systems: [electrons + nuclears] to the ground state.

There are *two possible ways* to make the rotation of adjacent sub-elementary fermion and sub-elementary antifermion compatible.

One of them is interaction 'side-by-side', like in the 1st and 3d pairs on Fig.3.2. In such a case they are rotating in opposite directions and their angular momenta (spins) compensate each other and the resulting spin of such a pair is zero. The resulting energy of such a pair of sub-elementary particle and antiparticle is also zero, because their asymmetry with respect to Bivacuum is exactly opposite, compensating each other.

The other way of compatibility is interaction 'head-to-tail', like in a central pair of sub-elementary fermions on Fig.3.2. In this configuration they rotate in the *same direction* and the sum of their spins is: $\mathbf{s} = \pm 1\hbar$. The energy of this pair is a sum of the *absolute values* of the energies of sub-elementary fermion and antifermion, as far their resulting symmetry shift is a sum of the symmetry shifts of each of them. From formula (3.70) of the next section we can see, that if the velocity of de Broglie wave is equal to the light velocity or very close to it, the contribution of the rest mass to its total energy is zero or almost zero. In such a case, pertinent for photon, its total energy is interrelated with photon frequency (ν_{ph}) and length like:

$$\mathbf{E}_{ph} = \mathbf{m}_{ph}\mathbf{c}^2 = \mathbf{h}\nu_{ph} = \mathbf{h}^2/(\mathbf{m}_{ph}\lambda_{ph}^2) \quad (3.53)$$

where: $\mathbf{m}_{ph}$ is the effective mass of photon and $\lambda_{ph} = \mathbf{c}/\nu_{ph}$ is the photon wave length.

It follows from our model, that the minimum value of the photon effective mass and energy is equal to the sum of *absolute values* of minimum mass/energy of central sub-elementary fermion and antifermion: $\mathbf{E}_{ph}^{\min} = \mathbf{m}_{ph}^{\min}\mathbf{c}^2 = 2\mathbf{m}_0\mathbf{c}^2$, i.e. the sum of the rest energy of an electron - positron pair. This consequence of our model is in accordance with available experimental data.

The law of energy conservation for elementary particles can be reformulated, as a *law of resulting Bivacuum symmetry conservation*. The additivity of different forms of energy, as a consequence of the energy conservation law, means the additivity of Bivacuum dipoles torus and antitorus energy difference (i.e. forms of kinetic energy) and sum of their absolute

values (forms of potential energy). These energy conservation roots are illustrated for one particle case by eqs.(3.60 and 3.60a).

**Model of photon, as a double
[electron + positron] rotating structure:**

$$< 2[\mathbf{F}_{\downarrow}^+ \bowtie \mathbf{F}_{\uparrow}^-] + (\mathbf{F}_{\downarrow}^- + \mathbf{F}_{\downarrow}^+) >_{S=\pm 1}$$




Fig.3.2 Model of photon $< 2[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_{S=0} + (\mathbf{F}_{\downarrow}^- + \mathbf{F}_{\downarrow}^+)_{S=\pm 1} >$, as result of fusion of electron and positron-like triplets $< [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-] + \mathbf{F}_{\downarrow}^{\pm} >$ of sub-elementary fermions, presented on Fig.3.1. The resulting symmetry shift of such structure is equal to zero, providing the absence or very close to zero rest mass of photon and its propagation in primordial Bivacuum with light velocity or very close to it in the asymmetric secondary Bivacuum.

We may see, that it has axially symmetric configurations in respect to the directions of rotation and propagation, which are normal to each other. These configurations periodically change in the process of sub-elementary fermions and antifermions correlated

[*Corpuscle* $\rightleftharpoons$ *Wave*] pulsations in composition of photon (Fig.3.2). The clockwise and counter clockwise rotation of photons around the z-axes stands for two possible polarizations of photon.

C.1 Correlation between new model of hadrons and conventional quark model of

protons, neutrons and mesons

The *proton* ($Z = +1$; $S = \pm 1/2$) is constructed by the same principle as the electron (Fig.3.1). It is a result of fusion of pair of sub-elementary fermion and antifermion $< [\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]_{S=0}^{\tau}$ and one unpaired $[\tau]$ sub-elementary fermion $(\mathbf{F}_{\downarrow}^{+})_{S=\pm 1/2}^{\tau}$ - *tauons*. These three components of proton have some similarity with quarks: $(\mathbf{F}_{\downarrow}^{+})_{S=\pm 1/2}^{\tau} \sim \mathbf{q}^{+} \sim \tau^{+}$ and antiquarks $(\mathbf{F}_{\uparrow}^{-})_{S=\pm 1/2}^{\tau} \sim \mathbf{q}^{-} \sim \tau^{-}$.

The difference with conventional quark model of protons and neutrons is that we do not need to use the notion of fractional charge in our proton model:

$$\mathbf{p} \equiv < [\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]_{S=0} + (\mathbf{F}_{\downarrow}^{+})_{S=\pm 1/2}^{\tau} >^{\tau} \quad (3.54)$$

$$or : \mathbf{p} \sim \left\langle [\mathbf{q}^{-} \bowtie \mathbf{q}^{+}]_{S=0} + (\mathbf{q}^{+})_{S=\pm 1/2} \right\rangle \quad (3.54a)$$

$$or : \mathbf{p} \sim \left\langle [\tau^{-} \bowtie \tau^{+}]_{S=0} + (\tau^{+})_{S=\pm 1/2} \right\rangle \quad (3.54b)$$

The charges, spins and mass/energy of sub-elementary particles and antiparticles in pairs $[\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]^{\tau}$ compensate each other. The resulting properties of protons ($\mathbf{p}$) are determined by unpaired/uncompensated sub-elementary particle $\mathbf{F}_{\downarrow}^{+} >_{S=\pm 1/2}^{\tau}$ of heavy τ -electrons generation (tauons).

The *neutron* ($Z = 0$; $S = \pm 1/2$) can be presented as:

$$\mathbf{n} \equiv < [\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]_{S=0}^{\tau} + [(\mathbf{F}_{\uparrow}^{+})^{\tau} \bowtie (\mathbf{F}_{\downarrow}^{-})^e]_{S=\pm 1/2} > \quad (3.55)$$

$$or : \mathbf{n} \sim [\mathbf{q}^{+} \bowtie \mathbf{q}^{-}]_{S=0}^{\tau} + (\mathbf{q}_{\uparrow}^0)_{S=\pm 1/2}^{\tau e} \quad (3.55a)$$

$$or : \mathbf{n} \sim [\tau^{+} \bowtie \tau^{-}]_{S=0}^{\tau} + ([\tau_{\uparrow}^{+}]^{\tau} \bowtie [\mathbf{F}_{\downarrow}^{-}]^e) \quad (3.55b)$$

where: the neutral quark $(\mathbf{q}_{\uparrow}^0)_{S=\pm 1/2}^{\tau e}$ is introduced, as a metastable complex of positive sub-elementary τ -fermion $(\mathbf{F}_{\downarrow}^{+})^{\tau}$ with negative sub-elementary fermion of opposite charge $[\mathbf{F}_{\uparrow}^{-}]^e$:

$$(\mathbf{q}_{\uparrow}^0)_{S=\pm 1/2}^{\tau e} = ([\mathbf{q}_{\uparrow}^{+}] \bowtie [\mathbf{F}_{\downarrow}^{-}]^e) \quad (3.56)$$

This means that the positive charge of unpaired heavy sub-elementary particle $(\mathbf{F}_{\uparrow}^+)^{\tau}$ in neutron ($\mathbf{n}$) is compensated by the charge of the light sub-elementary fermion $(\mathbf{F}_{\downarrow}^-)^e$. In contrast to charge, the spin of unpaired $(\mathbf{F}_{\uparrow}^+)^{\tau}$ is not compensated (totally) by spin of $(\mathbf{F}_{\downarrow}^-)^e$ in neutrons, because of strong mass and angular momentum difference in conditions of the $(\mathbf{F}_{\downarrow}^-)^e$ confinement. The mass of τ - electron, equal to that of τ -positron is: $\mathbf{m}_{\tau\pm} = 1782(3)$ MeV, the mass of the regular electron is: $\mathbf{m}_{e\pm} = 0,511003(1)$ MeV and the mass of μ -electron is: $\mathbf{m}_{\mu\pm} = 105,6595(2)$ MeV.

On the other hand, the mass of proton and neutron are correspondingly: $\mathbf{m}_{\mathbf{p}} = 938,280(3)$ MeV and $\mathbf{m}_{\mathbf{n}} = 939,573(3)$ MeV. They are about two times less than the mass of τ - electron and equal in accordance to our model to the mass of its unpaired sub-elementary fermion $(\mathbf{F}_{\uparrow}^+)^{\tau}$. This difference characterizes the energy of neutral massless *gluons* (exchange bosons), stabilizing the triplets of protons and neutrons. In the case of neutrons this difference is a bit less (taking into account the mass of $[\mathbf{F}_{\downarrow}^-]^e$), providing much shorter life-time of isolated neutrons (918 sec.) than that of protons ($>10^{31}$ years).

In accordance with our *hadron (baryon)* models, each of three quarks (sub-elementary fermions of τ - generation) in **protons** and **neutrons** can exist in 3 states (*red*, *green* and *blue*), but not simultaneously:

1. The *red* state of **quark/antiquark** means that it is in corpuscular [C] phase;
2. The *green* state of **quark/antiquark** means that it is in wave [W] phase;
3. The *blue* state means that **quark/antiquark** $(\mathbf{F}_{\uparrow}^{\pm})^{\tau}$ is in the transition $[C] \rightleftharpoons [W]$ state.

The 8 different combinations of the above defined states of 3 quarks of protons and neutrons correspond to 8 *gluons colors*, stabilizing these *hadrons* [61]. The gluons with boson properties are represented by pairs of Cumulative Virtual Clouds ($\mathbf{CVC}^+ \bowtie \mathbf{CVC}^-$), emitted $\rightleftharpoons$ absorbed in the process of the in-phase and counterphase $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of paired quarks + antiquark or tauon + antitauon. These 8 gluons, responsible for strong interaction, can be presented as a different combinations $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ transition states of $\mathbf{q}^-$ and $\mathbf{q}^+$, corresponding to two spin states of proton ($S = \pm 1/2 \hbar$), equal to that of unpaired quark - tauon [23].

The known experimental values of life-times of μ and τ electrons, corresponding in accordance to our model, to monomeric asymmetric sub-elementary fermions $(\mathbf{BVF}_{as}^{\downarrow})^{\mu,\tau}$, are equal only to $2.19 \times 10^{-6}s$ and $3.4 \times 10^{-13}s$, respectively. We assume here, that stabil-

ity of monomeric sub-elementary particles/antiparticles of $\mathbf{e}$, μ and τ generations, strongly increases as a result of their fusion in triplets, which became possible at Golden mean conditions.

The well known example of weak interaction, like β – *decay* of the neutron to proton, electron and $\mathbf{e}$ –antineutrino:

$$\mathbf{n} \rightarrow \mathbf{p} + \mathbf{e}^- + \tilde{\nu}_e \quad (3.57)$$

$$or : \left([\mathbf{q}^+ \bowtie \tilde{\mathbf{q}}^-] + (\mathbf{q}_{\uparrow}^0)^{\tau e}_{S=\pm 1/2} \right) \rightarrow ([\mathbf{q}^+ \bowtie \tilde{\mathbf{q}}^-] + \mathbf{q}^+) + \mathbf{e}^- + \tilde{\nu}_e \quad (3.57a)$$

is in accordance with our model of elementary particles, neutrino and antineutrino, are a packet of pairs of interrelated positive and negative virtual pressure waves $[\mathbf{VPW}^+ \bowtie \mathbf{VPW}^-]$ of two possible polarization (see section F3).

The reduced sub-elementary fermion of τ – generation in composition of a proton or neutron can be considered, as a quark and the sub-elementary antifermion, as an antiquark:

$$(\mathbf{F}_{\uparrow}^+)^{\tau} \sim \mathbf{q}^+ \quad and \quad (\mathbf{F}_{\downarrow}^-)^{\tau} \sim \tilde{\mathbf{q}}^- \quad (3.58)$$

In the process of β –decay of the neutron (3.57; 3.57a) the unpaired negative sub-elementary fermion $[\mathbf{F}_{\uparrow}^-]^e$ in the neutral quark $(\mathbf{q}_{\uparrow}^0)^{\tau e}_{S=\pm 1/2}$ (see 3.56) fuses with asymmetric virtual pair $[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]^e_{S=0}$, which emerged from the vicinal to neutron polarized Cooper pair of Bivacuum fermion and antifermion. The result of this fusion is a release of the real electron and electronic antineutrino:

$$[\mathbf{F}_{\uparrow}^-]^e + [\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]^e_{S=0} \rightarrow \mathbf{e}^- + \tilde{\nu}_e \quad (3.58a)$$

The antineutrino $\tilde{\nu}_e$ is a consequence of inelastic recoil effect in Bivacuum matrix, accompanied the electron fusion and its separation from hadron. The neutrino and antineutrino of three generation can be considered as the asymmetric superposition of positive and negative virtual pressure waves: $[\mathbf{VPW}^+ \bowtie \mathbf{VPW}^-]_{e,\mu,\tau}$ with clockwise or counterclockwise rotation of their plane, which determines spirality/spin of neutrino ($\pm 1/2\hbar$).

The energy of 8 gluons, corresponding to different superposition of $[\mathbf{CVC}^+ \bowtie \mathbf{CVC}^-]_{S=0,1}$, emitted and absorbed with the in-phase $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of pair [quark + antiquark] in the baryons triplets:

$$[\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-]_{S=0,1}^{\tau} = [\mathbf{q}^+ + \tilde{\mathbf{q}}^-]_{S=0,1} \quad (3.59)$$

is about 50% of energy/mass of quarks. This explains, why the mass of isolated unstable tauon and antitauon is about 2 times bigger, than their mass in composition of proton or neutron.

It looks, that our model of elementary particles is compatible with existing data and avoid the strong assumption of fractional charge. We anticipate that future experiments, like deep inelastic scattering inside the hadrons, will be able to choose between models of fractal and integer charge of the quarks.

One of the versions of elementary particle fusion have some similarity with thermonuclear *fusion* and can be as follows. The rest mass of *isolated* sub-elementary fermion/antifermion *before* fusion of the electron or proton, is equal to the rest mass of unstable muon or tauon, correspondingly. The 200 times decrease of muons mass is a result of mass defect, equal to the binding energy of triplets: electrons or positrons. It is provided by origination of electronic *e-gluons* and release of the huge amount of excessive kinetic (thermal) energy, for example in form of high energy photons or *e-neutrino* beams.

In protons, as a result of fusion of three τ —electrons/positrons, the contribution of hadron *h-gluon* energy to mass defect is only about 50% of their mass. However, the absolute hadron fusion energy yield is higher, than that of the electrons and positrons.

Our hypothesis of stable electron/positron and hadron fusion from short-living μ and τ - electrons, as a precursor of electronic and hadronic quarks, correspondingly, can be verified using special collider [22, 23].

In accordance to our Unified Theory, there are two different mechanisms of stabilization of the electron and proton structures in form of triplets of sub-elementary fermions/antifermions of the reduced μ and τ generations, correspondingly, preventing them from exploding under the action of self-charge:

1. Each of sub-elementary fermion/antifermion, representing asymmetric pair of torus (V^+) and antitorus (V^-), as a charge, magnetic and mass dipole, is stabilized by the Coulomb, magnetic and gravitational attraction between torus and antitorus;
2. The stability of triplet, as a whole, is provided by the exchange of Cumulative Virtual Clouds (CVC^+ and CVC^-) between three sub-elementary fermions/antifermions in the process of their $[C \rightleftharpoons W]$ pulsation. In the case of proton and neutron, the 8 transition states corresponds to 8 *h-gluons* of hadrons, responsible for strong interaction. In the case of the electron or positron, the stabilization of triplets is realized by 8 lighter *e-gluons* [23]. The

process of internal exchange of pairs $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0,1}^{e,p}$ with unpaired sub-elementary fermion in triplets is accompanied by the external energy exchange with Bivacuum dipoles, modulating positive and negative virtual pressure waves $[\mathbf{VPW}^+ \bowtie \mathbf{VPW}^-]$. The feedback reaction between Bivacuum dipoles and elementary particles is also existing [23].

D. Total, potential and kinetic energies of elementary de Broglie waves

The total energy of sub-elementary particles of triplets of the electrons or protons $< [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0} + (\mathbf{F}_\uparrow^\pm)_{S=\pm 1/2} >^{e,p}$ we can present in three modes, as a sum of total potential $\mathbf{V}_{tot}$ and kinetic $\mathbf{T}_{tot}$ energies, including the internal and external contributions:

$$\mathbf{E}_{tot} = \mathbf{V}_{tot} + \mathbf{T}_{tot} = \frac{1}{2}(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2 + \frac{1}{2}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 \quad (3.60)$$

$$\mathbf{E}_{tot} = \mathbf{m}_V^+\mathbf{c}^2 = \frac{1}{2}\mathbf{m}_V^+(2\mathbf{c}^2 - \mathbf{v}^2) + \frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2 \quad (3.60a)$$

$$\mathbf{E}_{tot} = 2\mathbf{T}_k(\mathbf{v}/\mathbf{c})^2 = \frac{1}{2}\mathbf{m}_V^+\mathbf{c}^2[1 + \mathbf{R}^2] + \frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2 \quad (3.60b)$$

where: $\mathbf{R} = \mathbf{m}_0/\mathbf{m}_V^+ = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ is the dimensionless relativistic factor; $\mathbf{v}$ is external translational - rotational velocity of particle. One may see, that $\mathbf{E}_{tot} \rightarrow \mathbf{m}_0\mathbf{c}^2$ at $\mathbf{v} \rightarrow \mathbf{0}$ and $\mathbf{m}_V^+ \rightarrow \mathbf{m}_0$.

Taking into account that $\frac{1}{2}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 = \frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2$ and $\mathbf{c}^2 = \mathbf{v}_{gr}\mathbf{v}_{ph}$, where $\mathbf{v}_{gr} \equiv \mathbf{v}$, then dividing the left and right parts of (3.60 and 3.60a) by $\frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2$, we get:

$$2\frac{\mathbf{c}^2}{\mathbf{v}^2} - 1 = 2\frac{\mathbf{v}_{ph}}{\mathbf{v}_{gr}} - 1 = \frac{(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2}{\mathbf{m}_V^+\mathbf{v}^2} = \frac{\mathbf{m}_V^+ + \mathbf{m}_V^-}{\mathbf{m}_V^+ - \mathbf{m}_V^-} \quad (3.61)$$

Comparing formula (3.61) with known relation for relativistic de Broglie wave for ratio of its potential and kinetic energy (Grawford, 1973), we get the confirmation of our definitions of potential and kinetic energies of elementary particle in (3.60):

$$2\frac{\mathbf{v}_{ph}}{\mathbf{v}_{gr}} - 1 = \frac{\mathbf{V}_{tot}}{\mathbf{T}_{tot}} = \frac{\mathbf{m}_V^+ + \mathbf{m}_V^-}{\mathbf{m}_V^+ - \mathbf{m}_V^-} \quad (3.62)$$

In Golden mean conditions, necessary for triplet fusion, the ratio $(\mathbf{V}_{tot}/\mathbf{T}_{tot})^\phi = (1/\phi + \phi) = 2.236$.

Consequently, the total potential ($\mathbf{V}_{tot}$) and kinetic ($\mathbf{T}_{tot}$) energies of sub-elementary fermions and their increments can be presented as:

$$\mathbf{V}_{tot} = \frac{1}{2}(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2 = \frac{1}{2}\mathbf{m}_V^+ (2\mathbf{c}^2 - \mathbf{v}^2) = \frac{1}{2}\frac{\hbar\mathbf{c}}{\mathbf{L}_{\mathbf{V}_{tot}}} \gg \mathbf{V}_{tot}^\phi; \quad (3.63)$$

$$\Delta\mathbf{V}_{tot} = \Delta\mathbf{m}_V^+\mathbf{c}^2 - \Delta\mathbf{T}_{tot} = -\frac{1}{2}\frac{\hbar\mathbf{c}}{\mathbf{L}_{\mathbf{V}_{tot}}}\frac{\Delta\mathbf{L}_{\mathbf{V}_{tot}}}{\mathbf{L}_{\mathbf{V}_{tot}}} = -\mathbf{V}_p\frac{\Delta\mathbf{L}_{\mathbf{V}_{tot}}}{\mathbf{L}_{\mathbf{V}_{tot}}} \quad (3.63a)$$

where: the characteristic velocity of potential energy, squared, is related to the group velocity of particle ($\mathbf{v}$), as $\mathbf{v}_p^2 = \mathbf{c}^2(2 - \mathbf{v}^2/\mathbf{c}^2)$ and the characteristic *curvature of potential energy* of elementary particles is:

$$\mathbf{L}_{\mathbf{V}_{tot}} = \frac{\hbar}{(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}} \ll \mathbf{L}_0^\phi \quad \text{at} \quad \left(\frac{\mathbf{v}_{tot}}{\mathbf{c}}\right)^2 \gg \phi \quad (3.64)$$

The total kinetic energy of unpaired sub-elementary fermion of triplets includes the internal vortical dynamics and external translational one, which determines their de Broglie wave length, ($\lambda_B = 2\pi\mathbf{L}_{\mathbf{T}_{ext}}$) :

$$\mathbf{T}_{tot} = \frac{1}{2}|\mathbf{m}_V^+ - \mathbf{m}_V^-|\mathbf{c}^2 = \frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2 = \frac{1}{2}\frac{\hbar\mathbf{c}}{\mathbf{L}_{\mathbf{T}_{tot}}} \gg \mathbf{T}_{tot}^\phi; \quad (3.65)$$

$$\Delta\mathbf{T}_{tot} = \mathbf{T}_{tot}\frac{1 + \mathbf{R}^2}{\mathbf{R}^2}\frac{\Delta\mathbf{v}}{\mathbf{v}} = -\mathbf{T}_k\frac{\Delta\mathbf{L}_{\mathbf{T}_{tot}}}{\mathbf{L}_{\mathbf{T}_{tot}}} \quad (3.65a)$$

where the characteristic *curvature of kinetic energy* of sub-elementary particles in triplets is:

$$\mathbf{L}_{\mathbf{T}_{tot}} = \frac{\hbar}{(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}} \ll \mathbf{L}_0^\phi \quad \text{at} \quad \left(\frac{\mathbf{v}_{tot}}{\mathbf{c}}\right)^2 \gg \phi \quad (3.65b)$$

It is important to note, that in compositions of triplets $< [\mathbf{F}_\uparrow^- \rtimes \mathbf{F}_\downarrow^+]_{S=0} + \left(\mathbf{F}_\uparrow^\pm\right)_{S=\pm 1/2} >^{e,p}$ the *minimum* values of *total* potential and kinetic energies and the *maximum* values of their characteristic curvatures correspond to that, determined by Golden mean conditions (see eqs. 3.52 and 3.52a). In our formulas above it is reflected by corresponding inequalities. In accordance to our theory, the Golden mean conditions determine a threshold for triplets fusion from sub-elementary fermions.

The increment of total energy of elementary particle is a sum of total potential and kinetic energies increments:

$$\Delta\mathbf{E}_{tot} = \Delta\mathbf{V}_{tot} + \Delta\mathbf{T}_{tot} = -\mathbf{V}_{tot}\frac{\Delta\mathbf{L}_{\mathbf{V}_{tot}}}{\mathbf{L}_{\mathbf{V}_{tot}}} - \mathbf{T}_{tot}\frac{\Delta\mathbf{L}_{\mathbf{T}_{tot}}}{\mathbf{L}_{\mathbf{T}_{tot}}} \quad (3.66)$$

The well known Dirac equation for energy of a free relativistic particle, following also from Einstein relativistic formula (3.22), can be easily derived from (3.60a), multiplying its

left and right part on $\mathbf{m}_V^+ \mathbf{c}^2$ and using introduced mass compensation principle (3.24):

$$\mathbf{E}_{tot}^2 = (\mathbf{m}_V^+ \mathbf{c}^2)^2 = (\mathbf{m}_0 \mathbf{c}^2)^2 + (\mathbf{m}_V^+)^2 \mathbf{v}^2 \mathbf{c}^2 \quad (3.67)$$

where: $\mathbf{m}_0^2 = |\mathbf{m}_V^+ \mathbf{m}_V^-|$ and the actual inertial mass of torus of unpaired sub-elementary fermion in triplets is equal to regular mass of particle: $\mathbf{m}_V^+ = \mathbf{m}_0$.

Dividing the left and right parts of (3.67) to $\mathbf{m}_V^+ \mathbf{c}^2$, we may present the total energy of an elementary de Broglie wave, as a sum of *internal and external* energy contributions, in contrast to previous sum of *total potential and kinetic* energies (3.60):

$$\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{E}^{in} + \mathbf{E}^{ext} = \quad (3.68)$$

$$= \frac{\mathbf{m}_0}{\mathbf{m}_V^+} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + (\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} \quad (3.68a)$$

$$\mathbf{E}_{tot} = \mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + (\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} \quad (3.69)$$

$$or : \mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = (\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^4 / \mathbf{v}^2 \quad (3.69a)$$

$$\mathbf{E}_{tot} = \hbar \nu_{C \rightleftharpoons W} = \mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + \left[\frac{\hbar^2}{\mathbf{m}_V^+ \lambda_B^2} \right]_{tr}^{ext} \quad (3.70)$$

where: $\mathbf{R} \equiv \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ is relativistic factor, dependent on the *external* translational velocity ($\mathbf{v}$) of particle; $\mathbf{m}_V^+ = \mathbf{m}_0/\mathbf{R} = \mathbf{m}$ is the actual inertial mass of sub-elementary fermion; the external translational de Broglie wave length is: $\lambda_B = \frac{\hbar}{\mathbf{m}_V^+ \mathbf{v}}$ and $\nu_{C \rightleftharpoons W}$ is the resulting frequency of corpuscle - wave pulsation (see next section).

We can easily transform formula (3.69) to a mode, including the internal rotational parameters of sub-elementary fermion, necessary for the rest mass and charge origination:

$$\mathbf{E}_{tot} = \mathbf{R} (\mathbf{m}_0 \omega_0^2 \mathbf{L}_0^2)_{rot}^{in} + [(\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2]_{tr}^{ext} \quad (3.71)$$

where: $\mathbf{L}_0 = \hbar/\mathbf{m}_0 \mathbf{c}$ is the Compton radius of sub-elementary particle; $\omega_0 = \mathbf{m}_0 \mathbf{c}^2/\hbar$ is the angular Compton frequency of sub-elementary fermion rotation around the common axis in a triplet (Fig.3.1).

For potential energy of a sub-elementary fermion, we get from (3.69), assuming, that $(\mathbf{m}_V^+ \mathbf{v}^2) = 2\mathbf{T}_{tot}$ and $\mathbf{E}_{tot} = \mathbf{V}_{tot} + \mathbf{T}_{tot}$:

$$\mathbf{V}_{tot} = \mathbf{E}_{tot} - \frac{1}{2} (\mathbf{m}_V^+ \mathbf{v}^2) = \mathbf{R}^{tot} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + \frac{1}{2} (\mathbf{m}_V^+ \mathbf{v}^2) \quad (3.72)$$

The difference between potential and kinetic energies, as analog of Lagrangian, from (3.71) is:

$$\mathcal{L} = \mathbf{V}_p - \frac{1}{2} (\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} = \mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} \quad (3.73)$$

It follows from (3.71 - 3.73), that at $\mathbf{v} \rightarrow \mathbf{c}$, the *total* relativistic factor, involving both the external and internal translational - rotational dynamics of sub-elementary fermions in triplets: $\mathbf{R}^{tot} \equiv \sqrt{1 - (\mathbf{v}_{tot}/\mathbf{c})^2} \rightarrow 0$ and the rest mass contribution to total energy of sub-elementary particle also tends to zero: $\mathbf{R}^{tot} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} \rightarrow 0$. Consequently, the total potential and kinetic energies tend to equality $\mathbf{V}_{tot} \rightarrow \mathbf{T}_{tot}$, and the Lagrangian to zero. *This is a conditions for harmonic oscillations of the photon, propagating in unperturbed Bivacuum.*

The important formula for doubled external kinetic energy can be derived from (3.69), taking into account that the relativistic relation between the actual and rest mass is $\mathbf{m}_V^+ = \mathbf{m}_0/\mathbf{R}$:

$$2\mathbf{T}_k = \mathbf{m}_V^+ \mathbf{v}^2 = \mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{R} \mathbf{m}_0 \mathbf{c}^2 = \frac{\mathbf{m}_0 \mathbf{c}^2}{\mathbf{R}} (1^2 - \mathbf{R}^2) \quad or : \quad (3.74)$$

$$2\mathbf{T}_k = \frac{\mathbf{m}_0 \mathbf{c}^2}{\mathbf{R}} (1 - \mathbf{R})(1 + \mathbf{R}) = (1 + \mathbf{R}) [\mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{m}_0 \mathbf{c}^2] \quad (3.74a)$$

This formula is a background of the introduced in section (III: 7) notion of *Tuning energy* of Bivacuum Virtual Pressure Waves ($\mathbf{VPW}^\pm$).

Our expressions (3.60 - 3.74a) are more general, than the well known (3.67), as far they take into account the properties of both poles (actual and complementary) of Bivacuum dipoles and subdivide the total energy of particle to the internal and external or to kinetic and potential ones.

E. The dynamic mechanism of corpuscle-wave duality

It is generally accepted, that the manifestation of corpuscle - wave duality of a particle is dependent on the way in which the observer interacts with a system. However, the mechanism of duality, as a background of quantum physics, is still obscure.

It follows from our theory, that the [corpuscle (C) $\rightleftharpoons$ wave (W)] duality represents modulation of the internal (hidden) quantum beats frequency between the asymmetric 'actual' (torus) and 'complementary' (antitorus) states of sub-elementary fermions or antifermions by the external - empirical de Broglie wave frequency of these particles [23]. The [C] phase

of each sub-elementary fermions of triplets $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\downarrow^\pm >^i$ of elementary particles, like electrons and protons, exists as a mass and an electric and magnetic asymmetric dipole.

The $[\mathbf{C} \rightarrow \mathbf{W}]$ transition is a result of two stages superposition:

The 1st stage of transition is a reversible dissociation of charged sub-elementary fermion in $[\mathbf{C}]$ phase $(\mathbf{F}_\uparrow^\pm)_{\mathbf{C}}^{e^\pm}$ to charged Cumulative Virtual Cloud $(\mathbf{CVC}^\pm)_{\mathbf{F}_\uparrow^\pm}^{e^\pm - e_{anc}^\pm}$ of subquantum particles and the 'anchor' Bivacuum fermion in $\mathbf{C}$ phase $(\mathbf{BVF}_{anc}^\uparrow)_{\mathbf{C}}^{e_{anc}^\pm}$:

$$(I): \left[(\mathbf{F}_\uparrow^\pm)_{\mathbf{C}}^{e^\pm} \xrightarrow{\text{Recoil/Antirecoil}} (\mathbf{BVF}_{anc}^\uparrow)_{\mathbf{C}}^{e_{anc}^\pm} + (\mathbf{CVC}^\pm)_{\mathbf{F}_\uparrow^\pm}^{e^\pm - e_{anc}^\pm} \right]^i \quad (3.75)$$

where notations $e^\pm$, $e_{anc}^\pm$ and $e^\pm - e_{anc}^\pm$ mean, correspondingly, the total charge, the anchor charge and their difference, pertinent to $\mathbf{CVC}^\pm$.

The 2nd stage of $[\mathbf{C} \rightarrow \mathbf{W}]$ transition is a reversible dissociation of the anchor Bivacuum fermion $(\mathbf{BVF}_{anc}^\uparrow)^i = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]_{anc}^i$ to symmetric $(\mathbf{BVF}^\uparrow)^i$ and the anchor cumulative virtual cloud $(\mathbf{CVC}^\pm)_{\mathbf{BVF}_{anc}^\uparrow}$, with frequency, equal to the empirical frequency of de Broglie wave of particle:

$$(II): (\mathbf{BVF}_{anc}^\uparrow)_{\mathbf{C}}^{e_{anc}^\pm} \xrightarrow{\text{Recoil/Antirecoil}} \left[(\mathbf{BVF}^\uparrow)^0 + (\mathbf{CVC}^\pm)_{\mathbf{BVF}_{anc}^\uparrow}^{e_{anc}^\pm} \right]_W^i \quad (3.76)$$

The 2nd stage takes a place if $(\mathbf{BVF}_{anc}^\uparrow)^i$ is asymmetric, i.e. in the case of nonzero external translational - rotational velocity of particle, when the magnetic field originates. The beat frequency of $(\mathbf{BVF}_{anc}^\uparrow)^{e,p}$ is equal to that of the empirical de Broglie wave frequency: $\omega_B = \hbar/(\mathbf{m}_V^+ \mathbf{L}_B^2)$. The higher is the external kinetic energy of fermion, the higher is frequency ω_B and stronger magnetic field, generated by the anchor Bivacuum fermion. The frequency of the stage (II) oscillations modulates the internal frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation: $[\mathbf{R} \ \omega_0 = \mathbf{R} \ \mathbf{m}_0 \mathbf{c}^2 / \hbar]^{e,p}$, related to contribution of the rest mass energy to the total energy of the de Broglie wave [23].

The $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations of unpaired sub-elementary fermion $\mathbf{F}_\uparrow^\pm$, of triplets of the electrons or protons $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\downarrow^\pm >^{e,p}$ are in counterphase with the in-phase pulsation of paired sub-elementary fermion and antifermion, modulating Bivacuum virtual pressure waves $(\mathbf{VPW}^\pm)$:

$$[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_W^{e,p} \xrightarrow{\text{CVC}^+ + \text{CVC}^-} [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_C^{e,p} \quad (3.77)$$

The basic frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation, corresponding to Golden mean conditions, $(\mathbf{v}/\mathbf{c})^2 = 0,618 = \phi$, is equal to that of the 1st stage frequency (5.1) at zero external trans-

lational velocity ($\mathbf{v}_{tr}^{ext} = 0$; $\mathbf{R} = \mathbf{1}$). This frequency is the same as the basic Bivacuum virtual pressure waves ($\mathbf{VPW}_{q=1}^{\pm}$) and virtual spin waves ($\mathbf{VirSW}_{q=1}^{S=\pm 1/2}$) frequency (3.10): $[\omega_{q=1} = \mathbf{m}_0 \mathbf{c}^2 / \hbar]^i$.

The empirical parameters of de Broglie wave of elementary particle are determined by asymmetry of the torus and antitorus of the *anchor* Bivacuum fermion ($\mathbf{BVF}_{anc}^{\uparrow}$) $^{e,p} = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]_{anc}^{e,p}$ (Fig.3.1) and the frequency of its reversible dissociation to symmetric ($\mathbf{BVF}^{\updownarrow}$) i and the anchor cumulative virtual cloud ($\mathbf{CVC}_{anc}^{\pm}$) – stage (II) of duality mechanism (3.76).

The total energy, charge and spin of triplets - fermions, moving in space with velocity ($\mathbf{v}$) is determined by the unpaired sub-elementary fermion ($\mathbf{F}_{\updownarrow}^{\pm}$) $_z$, as far the paired ones in $[\mathbf{F}_{\up}^+ \bowtie \mathbf{F}_{\downarrow}^-]_{x,y}$ of triplets compensate each other. From (3.69 and 3.70) it is easy to get:

$$\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \hbar \omega_{C \rightleftharpoons W} = \mathbf{R} (\hbar \omega_0)_{rot}^{in} + (\hbar \omega_B^{ext})_{tr} = \mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + (\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} \quad (3.78)$$

$$\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{E}_{rot}^{in} + \mathbf{E}_{tr}^{ext} = \mathbf{R} (\mathbf{m}_0 \omega_0^2 \mathbf{L}_0^2)_{rot}^{in} + \left(\frac{\hbar^2}{\mathbf{m}_V^+ \lambda_B^2} \right)_{tr}^{ext} \quad (3.78a)$$

where: $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ is the relativistic factor; $\mathbf{v} \equiv \mathbf{v}_{tr}^{ext}$ is the external translational group velocity; $\lambda_B = \hbar / \mathbf{m}_V^+ \mathbf{v} = 2\pi \mathbf{L}_B$ is the external translational de Broglie wave length; the actual inertial mass is $\mathbf{m}_V^+ = \mathbf{m} = \mathbf{m}_0 / \mathbf{R}$; $\mathbf{L}_0^i = \hbar / \mathbf{m}_0^i \mathbf{c}$ is a Compton radius of the elementary particle.

It follows from our approach, that the fundamental phenomenon of **corpuscle – wave duality** (Fig.3.3) is a result of modulation of the primary - carrying frequency of the internal $[\mathbf{C} \rightleftharpoons \mathbf{W}]^{in}$ pulsation of individual sub-elementary fermions (*1st stage*): $(\omega^{in})^i = \mathbf{R} \omega_0^i = \mathbf{R} \mathbf{m}_0^i \mathbf{c}^2 / \hbar = \mathbf{R} \mathbf{c} / \mathbf{L}_0^i$, by the frequency of the external empirical de Broglie wave of triplet: $\omega_B^{ext} = \mathbf{m}_V^+ \mathbf{v}_{ext}^2 / \hbar = 2\pi \mathbf{v}_{ext} / \mathbf{L}_B$, equal to angular frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]_{anc}$ pulsation of the anchor Bivacuum fermion ($\mathbf{BVF}_{anc}^{\updownarrow}$) i (*2nd stage*).

The contribution of this external dynamics modulation of the internal one, determined by asymmetry of the *anchor* ($\mathbf{BVF}_{anc}^{\updownarrow}$) $^i = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]_{anc}^i$ to the total energy of particle, is determined by second terms in (3.78) and (3.78a):

$$\mathbf{E}_B^{ext} = (\hbar \omega_B^{ext})_{tr} = \left(\frac{\hbar^2}{\mathbf{m}_V^+ \lambda_B^2} \right)_{tr} = [(\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2]_{tr}^{ext} \quad (3.79)$$

$$= (\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} = (\mathbf{m}_V^+ \omega_B^2 \mathbf{L}_B^2)_{rot}^{ext} \quad (3.79a)$$

This *external* contribution is increasing with particle acceleration and tending to light velocity. At $\mathbf{v} \rightarrow \mathbf{c}$, $(\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} \rightarrow \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{E}_{tot}$.

In contrast to *external* translational contribution of triplets, the *internal* rotational contribution of individual unpaired sub-elementary fermions *inside the triplets* is tending to zero at the same conditions:

$$\mathbf{E}_{rot}^{in} = \mathbf{R} (\hbar\omega_0)_{rot}^{in} = \mathbf{R} (\mathbf{m}_0\omega_0\mathbf{L}_0^2)_{rot}^{in} = \mathbf{R} (\mathbf{m}_0\mathbf{c}^2)_{rot}^{in} \rightarrow 0 \quad \text{at } \mathbf{v} \rightarrow \mathbf{c} \quad (3.80)$$

as far at $\mathbf{v} \rightarrow \mathbf{c}$, the $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \rightarrow 0$.

For a regular nonrelativistic electron the carrier frequency is $\omega^{in} = R\omega_0^e \sim 10^{21}s^{-1} \gg \omega_B^{ext}$. However, for relativistic case at $\mathbf{v} \rightarrow \mathbf{c}$, the situation is opposite: $\omega_B^{ext} \gg \omega^{in}$ at $\omega^{in} \rightarrow 0$.




Fig. 3.3. Dynamic model of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of triplets of sub-elementary fermions/antifermions (the reduced by fusion to triplets μ and τ

electrons) composing, correspondingly, electron and proton $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\uparrow^\pm >^{e,p}$. The pulsation of the pair $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$, modulating virtual pressure waves of Bivacuum (VPW^+ and VPW^-), is counterphase to pulsation of unpaired sub-elementary fermion/antifermion $\mathbf{F}_\uparrow^\pm$.

The properties of the *anchor* Bivacuum fermion $\mathbf{BVF}_{anc}^\uparrow$ where analyzed [22], at three conditions:

1. The external translational velocity ($\mathbf{v}$) is zero;
2. The external translational velocity corresponds to Golden mean ($\mathbf{v} = \mathbf{c}\phi^{1/2}$);
3. The relativistic case, when $\mathbf{v} \sim \mathbf{c}$.

Under nonrelativistic conditions ($\mathbf{v} \ll \mathbf{c}$), the de Broglie wave (modulation) frequency is low: $2\pi(\nu_B)_{tr} \ll (\omega^{in} = \mathbf{R}\omega_0)$. However, in relativistic case ($\mathbf{v} \sim \mathbf{c}$), the modulation frequency of the 'anchor' ($\mathbf{BVF}_{anc}^\uparrow$), equal to that of de Broglie wave, can be higher, than the internal one : $2\pi(\nu_B)_{tr} \geq \omega^{in}$.

The paired sub-elementary fermion and antifermion of $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0}$ also have the 'anchor' Bivacuum fermion and antifermion ($\mathbf{BVF}_{anc}^\uparrow$), similar to that of unpaired. However, the opposite energies of their $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation compensate each other in accordance with proposed model.

F. The nature of electrostatic, magnetic and gravitational potentials, based on Unified theory

F.1 Electromagnetic dipole radiation as a consequence of charge oscillation

The $[\text{emission} \rightleftharpoons \text{absorption}]$ of photons in a course of elementary fermions - triplets $< [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0} + (\mathbf{F}_\uparrow^+)_{S=\pm 1/2} >^{e,\tau}$ vibrations can be described by known mechanism of the electric dipole radiation (ε_{EH}), induced by charge acceleration (a), following from Maxwell equations [68]:

$$\varepsilon_{EH} = \frac{2e^2}{3c^3} a^2 \quad (3.81)$$

The resulting frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of each of three sub-elementary fermions in triplets is a sum of internal frequency contribution ($\mathbf{R}\omega_0^{in}$) and the external frequency

(ω_B) of de Broglie wave from (3.78):

$$[\omega_{C \rightleftharpoons W} = \mathbf{R} \omega_0^{in} + \omega_B]^i \quad (3.82)$$

where: $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ is relativistic factor.

The acceleration can be related only with external translational dynamics which determines the empirical de Broglie wave parameters of particles. Acceleration is a result of alternating change of the charge deviation from the position of equilibrium: $\Delta\lambda_B(\mathbf{t}) = (\lambda_B^t - \lambda_0) \sin \omega_B \mathbf{t}$ with de Broglie wave frequency of triplets: $\omega_B = \hbar/(m_V^+ L_B^2)$, where $L_B = \hbar/m_V^+ \mathbf{v}$. It is accompanied by oscillation of the instant de Broglie wave length (λ_B^t).

The acceleration of charge in the process of $\mathbf{C} \rightleftharpoons \mathbf{W}$ pulsation of the anchor $\mathbf{BVF}_{anc}^\dagger$ can be expressed as:

$$\mathbf{a} = \omega_B^2 \Delta\lambda_B(\mathbf{t}) \quad (3.83)$$

$$\mathbf{a} = \omega_B^2 (\lambda_B - \lambda_0) \sin \omega_B \mathbf{t} \quad (3.84)$$

where: $\lambda_B^t = 2\pi \mathbf{L}_B$ is the instant de Broglie wave length of the particle and $\lambda_0 = \mathbf{h}/\mathbf{m}_0 \mathbf{c}$ is the Compton length of triplet.

The intensity of dipole radiation of pulsing $\mathbf{BVF}_{anc}^\dagger$ from 3.82 and 3.83 is:

$$\varepsilon_{EM} = \frac{2}{3c^3} \omega_B^4 (\mathbf{d}_E^t)^2 \quad (3.85)$$

where the oscillating electric dipole moment is: $\mathbf{d}_E^t = \mathbf{e} (\lambda_B - \lambda_0)$.

Consequently, in accordance with our model of duality, the EM dipole radiation is due to modulation of the anchor Bivacuum fermion $\mathbf{BVF}_{anc}^\dagger$ frequency of $\mathbf{C} \rightleftharpoons \mathbf{W}$ pulsation of unpaired sub-elementary fermion by thermal vibrations of triplets. The electromagnetic field, is a result of Corpuscle - Wave pulsation of photon (Fig.3.2) and its fast rotation with angle velocity (ω_{rot}), equal to pulsation frequency. The clockwise or anticlockwise direction of photon rotation as respect to direction of its propagation, corresponds to its two polarization.

F.2 The different kind of Bivacuum dipoles perturbation, induced by dynamics of elementary particles

In the process of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of sub-elementary particles in triplets $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\uparrow^\pm >^{e,p}$ the [local (internal) $\Leftrightarrow$ distant (external)] compensation effects stand for the

energy conservation law. The *local* effects are pertinent for the [C] phase of particles. They are confined in the volume of sub-elementary fermions and are determined by gravitational interaction between opposite actual and complementary mass, Coulomb attraction between opposite electric charges and magnetic attraction between magnetic moments of asymmetric torus and antitorus of sub - elementary fermions. These potential attraction forces are balanced by the energy gap between torus (V^+) and antitorus (V^-) and centripetal force of the axial rotation of triplets. The axis of triplet rotation is strictly related, in accordance with our model, with the direction of its translational propagation. It is supposed, that like magnetic field force lines, this rotation follows the *right hand screw rule* and is responsible for *magnetic field* origination. The total energy of triplet, the angular frequency of its rotation and the velocity of its translational propagation are interrelated.

The *distant* (external) effects, related with certain polarization and alignment of Bivacuum dipoles (circular and linear) are related with $[C \rightarrow W]$ transition of unpaired $F_{\downarrow}^{\pm} >^{e,p}$. It is accompanied by the recoil effect and the distant elastic deformation of Bivacuum superfluid matrix, shifting correspondingly a symmetry of Bivacuum dipoles. The reverse $[W \rightarrow C]$ transition represents the *antirecoil effect*. The latter is accompanied by giving back the delocalized recoil energy and restoration of the unpaired sub-elementary fermion and the whole triplet properties.

The recoil - antirecoil and polarization effects on surrounding Bivacuum dipoles ($BVF^{\uparrow}$ and $BVF^{\downarrow}$) in form of spherical elastic waves, excited by $[C \rightleftharpoons W]$ pulsations of *unpaired* sub-elementary fermions $F_{\downarrow}^+$ and antifermions $F_{\downarrow}^-$ of triplets $< [F_{\uparrow}^+ \bowtie F_{\downarrow}^-] + F_{\downarrow}^{\pm} >^{e,p}$, are *opposite*. This determines the opposite effects of positive and negative charged $F_{\downarrow}^-$ on symmetry of Bivacuum torus (V^+) and antitorus (V^-) of Bivacuum dipoles $BVF^{\uparrow}$, accompanied by their mass and charge polarization and acquiring the external momentum. Corresponding energy shift between torus and antitorus is dependent on distance (R) from charge as $(\vec{r}/R)$. The induced by such mechanism linear alignment of Bivacuum bosons ($BVB^{\pm}$) or Cooper pairs $[BVF^{\uparrow} \bowtie BVF^{\downarrow}]$ with opposite direction of $BVF^{\uparrow}$ and $BVF^{\downarrow}$ rotation between remote $F_{\downarrow}^+$ and $F_{\downarrow}^-$ of different triplets, like bundles of virtual microtubules stands for *electrostatic field and its 'force lines* origination.

On the other hand, pulsations of each of paired sub-elementary fermions $[F_{\uparrow}^+ \bowtie F_{\downarrow}^-]$ have similar and symmetric effect on excitation of (V^+) and (V^-) of $BVF^{\uparrow} = [V^+ \updownarrow V^-]$, independently on the charge of unpaired fermion and direction of triplets motion. Corre-

sponding excitation of positive and negative virtual pressure waves (VPW⁺ and VPW⁻) is a background of *gravitational* attraction between triplets, like between pulsing spheres in liquid medium. The influence of the in-phase recoil effect of pulsing $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]$ on the external momentum and energy of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of Bivacuum dipoles $\mathbf{BVF}^\uparrow = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]$ should be symmetric and equal by absolute value to increment of energy of unpaired $\mathbf{F}_\uparrow^\pm$.

The fast rotation of pairs of sub-elementary fermion and antifermion $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]$ of triplets with opposite charges and actual magnetic moments should induce a polarization of Bivacuum dipoles around the vector of triplets propagation, shifting the spin equilibrium $[\mathbf{BVF}^\uparrow \rightleftharpoons \mathbf{BVB}^\pm \rightleftharpoons \mathbf{BVF}^\downarrow]$ to the left or right and stimulate their 'head-to-tail' circular structures assembly in form of closed virtual microtubules. *Our conjecture is that corresponding circular/axial 'polymerization' of polarized Bivacuum dipoles around direction of current, coinciding with that of triplets propagation, stands for curled magnetic field origination.* The fast chaotic thermal motion of conducting electrons in metals or charged particles in plasma became more ordered in electric current, increasing correspondingly the magnetic cumulative effects due to triplet fast rotation in the same plane.

It is possible to present the above explanation of field nature and origination in more formal way. The total energies of $[\mathbf{C} \rightarrow \mathbf{W}]$ and $[\mathbf{W} \rightarrow \mathbf{C}]$ transitions of particles we present using general formula (3.60). However, here we take into account the corresponding *recoil* $\rightleftharpoons$ *antirecoil* effects and the reversible conversion of the internal - local (Loc) gravitational, Coulomb and magnetic potentials to the external - distant (Dis) Bivacuum matrix perturbation, stimulated by these transitions. The $[\mathbf{C} \rightarrow \mathbf{W}]$ transition, accompanied by three kinds of *recoil* effects, can be described as:

$$\mathbf{E}^{C \rightarrow W} = \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{V}_{tot} + [(\mathbf{E}_G)_{Rec}^{Loc} - (\mathbf{E}_G)_{Rec}^{Dist}] + \quad (3.86)$$

$$+ \mathbf{T}_{tot} + [(\mathbf{E}_E)_{Rec}^{Loc} - (\mathbf{E}_E)_{Rec}^{Dist}]_{tr} + \quad (3.86a)$$

$$+ [(\mathbf{E}_H)_{Rec}^{Loc} - (\mathbf{E}_H)_{Rec}^{Dist}]_{rot} \quad (3.86b)$$

In the process of the reverse $[\mathbf{W} \rightarrow \mathbf{C}]$ transition the unpaired sub-elementary fermion $\mathbf{F}_\uparrow^\pm$ of triplet $< [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0} + (\mathbf{F}_\uparrow^\pm)_{S=\pm 1/2} >$ gets back the *antirecoil* (ARec) energy in form of relaxation of Bivacuum matrix, transforming back the Bivacuum matrix elastic deformation and VPW[±] excitation:

$$\mathbf{E}^{W \rightarrow C} = \mathbf{m}_V^+ \mathbf{c}^2 = \mathbf{V}_{tot} + [-(\mathbf{E}_G)_{ARec}^{Loc} + (\mathbf{E}_G)_{ARec}^{Dist}] + \quad (3.87)$$

$$+ \mathbf{T}_{tot} + [-(\mathbf{E}_E)_{ARec}^{Loc} + (\mathbf{E}_E)_{ARec}^{Dist}]_{tr} + \quad (3.87a)$$

$$+ [-(\mathbf{E}_H)_{ARec}^{Loc} + (\mathbf{E}_H)_{ARec}^{Dist}]_{rot} + \quad (3.87b)$$

where: $\mathbf{V}_{tot} = \frac{1}{2}(\mathbf{m}_V^+ + \mathbf{m}_V^-) \mathbf{c}^2 = \frac{1}{2} \mathbf{m}_V^+ (2\mathbf{c}^2 - \mathbf{v}^2)$ is a total potential energy of triplet (3.63), equal to that of unpaired sub-elementary fermion, and $\mathbf{T}_{tot} = \frac{1}{2}(\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2$ is its total kinetic energy (3.65).

The reversible conversions of the *local gravitational potential* $\pm(\mathbf{V}_G)_{Rec, ARec}^{Loc}$ to the distant one $\pm(\mathbf{V}_G)_{Rec, ARec}^{Dist}$, accompanied the $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of unpaired sub-elementary fermions of triplets, interrelated with paired ones $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]$. The latter excite the positive $\mathbf{VPW}^+$ and negative $\mathbf{VPW}^-$ spherical virtual pressure waves, propagating in space with light velocity. They are a consequence of transitions of torus $\mathbf{V}^+$ and antitorus $\mathbf{V}^-$ of surrounding Bivacuum dipoles $\mathbf{BVF}^\uparrow = [\mathbf{V}^+ \updownarrow \mathbf{V}^-]$ between the excited and ground states.

The $\pm(\mathbf{E}_E)_{Rec, ARec}^{Loc}$ and $\pm(\mathbf{E}_E)_{Rec, ARec}^{Dist}$ are the local and distant electrostatic potential oscillations; the $\pm(\mathbf{E}_H)_{Rec, ARec}^{Loc}$ and $\pm(\mathbf{E}_H)_{Rec, ARec}^{Dist}$ are the local and distant magnetic potentials oscillations, accompanied $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations and rotation of triplets.

Let us consider in more detail the interconversions of the *internal - local* and the *external - distant* gravitational, Coulomb and magnetic potentials of charged elementary particles, like electron or proton.

F3. The new approach to quantum gravity and antigravity.

*The mechanism of relativistic mass increasing and
a reason for gravitational and inertial mass equality.*

The unified right parts of eqs. (3.86) and (3.87), describing the excitation of *gravitational waves*, which represent superposition of positive and negative virtual pressure waves ($\mathbf{VPW}^+$ and $\mathbf{VPW}^-$), are:

$$\overline{\mathbf{V}}_{tot}^{C=W} = \mathbf{V}_{tot} \pm [(\mathbf{E}_G)_{Rec}^{Loc} - (\mathbf{E}_G)_{Rec}^{Dist}] \sim \quad (3.88)$$

$$\sim \mathbf{VPW}^\pm \pm [\Delta \mathbf{VPW}^+ + \Delta \mathbf{VPW}^-] \quad (3.88a)$$

This formula reflects the fluctuations of *potential energy* of triplets, induced by $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of unpaired sub-elementary fermions (3.75 and 3.76), interrelated with and paired ones (5.3), which are responsible for gravitational field.

The more detailed presentation of (3.88) is:

$$\begin{aligned} \overline{\mathbf{V}}_{tot}^{\mathbf{C}=\mathbf{W}} = \frac{1}{2}(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2 \pm \\ \pm \left\{ \left[\mathbf{G} \frac{(\mathbf{m}_V^+ \mathbf{m}_V^-)}{\mathbf{L}_G} \right]^{Loc} - \left[\left(\frac{\mathbf{m}_0}{\mathbf{M}_{Pl}} \right)^2 (\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2 \right]^{Dis} \right\} \end{aligned} \quad (3.89)$$

The local *internal* gravitational interaction between the opposite mass poles of the mass-dipoles of unpaired sub-elementary fermions (antifermions) $(\mathbf{F}_{\uparrow}^{\pm})_{S=\pm 1/2}$ turns reversibly to the *external* distant one:

$$\left[\mathbf{G} \frac{|\mathbf{m}_V^+ \mathbf{m}_V^-|}{\mathbf{L}_G} \right]^{Loc} \xrightleftharpoons[\mathbf{W} \rightarrow \mathbf{C}]{\mathbf{C} \rightarrow \mathbf{W}} \left[\beta^i (\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}^2 = \beta^i \mathbf{m}_V^+ (2\mathbf{c}^2 - \mathbf{v}^2) \right]^{Dis} \quad (3.90)$$

where: $\mathbf{L}_G = \hbar/(\mathbf{m}_V^+ + \mathbf{m}_V^-)\mathbf{c}$ is a characteristic curvature of potential energy (3.64); $\mathbf{M}_{Pl}^2 = \hbar\mathbf{c}/\mathbf{G}$ is a Plank mass; $\mathbf{m}_0^2 = |\mathbf{m}_V^+ \mathbf{m}_V^-|$ is a rest mass squared; $\beta^i = \left(\frac{\mathbf{m}_0^i}{\mathbf{M}_{Pl}} \right)^2$ is the introduced earlier dimensionless gravitational fine structure constant [21-23]. For the electron $\beta^e = 1.739 \times 10^{-45}$ and $\sqrt{\beta^e} = \frac{\mathbf{m}_0^e}{\mathbf{M}_{Pl}} = 0.41 \times 10^{-22}$

The excitation of the *external* - distant spherical virtual pressure waves of positive and negative energy: $\mathbf{VPW}^+$ and $\mathbf{VPW}^-$ is a result of torus and antitorus energy fluctuations, accompanied $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of paired sub-elementary fermions $[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_{S=0}$, strictly equal by absolute values to fluctuation energy of the unpaired $(\mathbf{F}_{\uparrow}^{\pm})_{S=\pm 1/2}$ one. It is important to note, that the introduced gravitational field does not depend on charge of triplet, determined by unpaired sub-elementary fermion $< [\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_{S=0} + (\mathbf{F}_{\uparrow}^{\pm})_{S=\pm 1/2} >$, in contrast to electrostatic and magnetic field.

From the proposed mechanism of gravitation and similar values of $(\mathbf{m}_V^+)$ in the left and right parts of eq. 3.90 it directly follows *the equality of gravitational and inertial mass*. The *inertia* itself is a consequence of particles system symmetry and *effective* mass relativistic increasing with acceleration. Consequently, inertia is proportional to the number of particles in this system and (i.e. total mass of the system) and a generalized Le Chatelier's Principle, as a reluctance/metastability of system to additional symmetry shift.

In accordance to our hypothesis [21-23], **the mechanism of gravitational attraction** is similar to Bjerknes attraction between pulsing spheres in liquid medium - Bivacuum. The

dependence of Bjerknes force on the distance between centers of pulsing objects is quadratic:
 $\mathbf{F}_{Bj} \sim 1/r^2$:

$$\mathbf{F}_G = \mathbf{F}_{Bj} = \frac{1}{r^2} \pi \rho_G \mathbf{R}_1^2 \mathbf{R}_2^2 \mathbf{v}^2 \cos \beta \quad (3.91)$$

where ρ_G is density of liquid, i.e. virtual density of secondary Bivacuum. It is determined by Bivacuum dipoles (BVF[†] and BVB[±]) symmetry shift; $\mathbf{R}_1$ and $\mathbf{R}_2$ radiuses of pulsing/gravitating spheres; $\mathbf{v}$ is velocity of spheres surface oscillation (i.e. velocity of $\mathbf{VPW}^\pm$, excited by $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of elementary particles, which can be assumed to be equal to light velocity: $\mathbf{v} = \mathbf{c}$); β is a phase shift between pulsation of spheres.

It is important to note, that on the big enough distances the *attraction* may turn to *repulsion*. The latter effect, depending on the phase shift of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of interacting remote triplets (β), can explain the revealed acceleration of the Universe expansion. The possibility of artificial phase shift of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of coherent elementary particles of any object may (for example by magnetic field) may change its gravitational attraction to repulsion. The volume and radius of pulsing spheres (R_1 and R_2) in such approach is determined by sum of volume of hadrons, composing gravitating systems in solid, liquid, gas or plasma state. The gravitational attraction or repulsion is a result of increasing or decreasing of virtual pressure of subquantum particles between interacting systems as respect to its value outside them. This model can serve as a background for new quantum gravity theory.

The effective radiuses of gravitating objects $\mathbf{R}_1$ and $\mathbf{R}_2$ can be calculated from the effective volumes of the objects:

$$\mathbf{V}_{1,2} = \frac{4}{3} \pi \mathbf{R}_{1,2}^3 = \mathbf{N}_{1,2} \frac{4}{3} \pi \mathbf{L}_{p,n}^3 \quad (3.91a)$$

where: $\mathbf{N}_{1,2} = \mathbf{M}_{1,2}/\mathbf{m}_{p,n}$ is the number of protons and neutrons in gravitating bodies with mass $\mathbf{M}_{1,2}$; $\mathbf{m}_{p,n}$ is the mass of proton and neutron; $\mathbf{L}_{p,n} = \hbar/\mathbf{m}_{p,n}\mathbf{c}$ is the Compton radius of proton and neutron.

From (3.91a) we get for effective radiuses:

$$\mathbf{R}_{1,2} = \left(\frac{\mathbf{M}_{1,2}}{\mathbf{m}_{p,n}} \right)^{1/3} \mathbf{L}_{p,n} = \left(\frac{\mathbf{M}_{1,2}}{\mathbf{m}_{p,n}} \right)^{1/3} \frac{\hbar}{\mathbf{m}_{p,n}\mathbf{c}}$$

Putting this to (3.91) we get for the force of gravitational interaction between two macroscopic objects, each of them formed by atoms with coherently pulsing protons and neutrons:

$$\mathbf{F}_G = \frac{1}{r^2} \pi \rho_{Bv} \frac{(\mathbf{M}_1 \mathbf{M}_2)^{2/3}}{\mathbf{m}_{p,n}^{4/3}} \left(\frac{\hbar}{\mathbf{m}_{p,n}} \right)^4 \frac{1}{\mathbf{c}^2} \quad (3.91b)$$

Equalizing this formula with Newton's one: $\mathbf{F}_G^N = \frac{1}{r^2} \mathbf{G} (\mathbf{M}_1 \mathbf{M}_2)$, we get the expression for gravitational constant:

$$\mathbf{G} = \pi \frac{\rho_G}{\sqrt[3]{\mathbf{M}_1 \mathbf{M}_2}} \frac{\hbar^2 / \mathbf{c}^2}{\sqrt[3]{\mathbf{m}_{p,n}^{16}}} \quad (3.91c)$$

The condition of gravitational constant permanency from (3.91c), is the anticipated from our theory interrelation between the mass of gravitating bodies $\sqrt[3]{\mathbf{M}_1 \mathbf{M}_2}$ and the virtual density ρ_G of secondary Bivacuum, determined by Bivacuum fermions symmetry shift and excitation in gravitational field:

$$\mathbf{G} = \text{const}, \quad \text{if} \quad \frac{\rho_G}{\sqrt[3]{\mathbf{M}_1 \mathbf{M}_2}} = \text{const} \quad (3.91d)$$

where, taking into account (3.90):

$$\sqrt[3]{\mathbf{M}_1 \mathbf{M}_2} \sim \rho_G = \frac{\frac{1}{2} \left(\frac{\mathbf{m}_0}{\mathbf{M}_{P1}} \right)^2 (\mathbf{m}_V^+ + \mathbf{m}_V^-)}{\frac{3}{4} \pi \mathbf{L}_G^3} = \frac{2}{3} \frac{\left(\frac{\mathbf{m}_0}{\mathbf{M}_{P1}} \right)^2 \mathbf{m}_V^+ (2 - \mathbf{v}^2 / \mathbf{c}^2)}{\pi \mathbf{L}_G^3}$$

assuming, that the radius of characteristic volume of asymmetry of Bivacuum fermion, responsible for gravitation is:

$$\mathbf{L}_G = \frac{\hbar}{(\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}}$$

we get for reduced gravitational density:

$$\rho_G = \frac{2}{3} \frac{1}{\pi \hbar^3} \left(\frac{\mathbf{m}_0}{\mathbf{M}_{P1}} \right)^2 \cdot (\mathbf{m}_V^+ + \mathbf{m}_V^-) (\mathbf{m}_V^+ - \mathbf{m}_V^-)^3 \mathbf{c}^3 \quad (3.91e)$$

we may see that at conditions of ideal symmetry of primordial Bivacuum, i.e. in the absence of matter and fields the virtual density of Bivacuum and gravitational interaction is equal to zero ($\rho_G = 0$), as far $(\mathbf{m}_V^+ - \mathbf{m}_V^-) = 0$.

F.4. Possible nature of neutrino and antineutrino

We put forward a conjecture, that neutrinos or antineutrinos represent a *stable non elastic* Bivacuum excitations in form of packet of positive and negative virtual pressure waves, slightly uncompensated each other:

$$\nu, \tilde{\nu} \sim [\mathbf{VPW}^+ \boxtimes \mathbf{VPW}^-] \sim \mathbf{VirP}^+ - \mathbf{VirP}^-$$

However, their nonlocal asymmetry is compensating the local asymmetry in Bivacuum, accompanied origination of positrons or electrons. This asymmetry can be described as a small

difference in energy of sub-elementary fermion and antifermion $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0}$, composing pair in triplets $< [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+] + (\mathbf{F}_\uparrow^\pm) >$. It can be positive for electrons and negative for positrons and expressed as energy increment of elementary de Broglie wave:

$$\mathbf{E}_\nu^i = \pm \left[(\mathbf{m}_V^+ \mathbf{c}^2)_{[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]}^{\mathbf{F}_\downarrow^+} - (\mathbf{m}_V^- \mathbf{c}^2)_{[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]}^{\mathbf{F}_\downarrow^-} \right]^i$$

$$\text{or : } [\Delta \mathbf{m}_V^\pm \mathbf{c}^2 = \Delta(\mathbf{Rm}_0 \mathbf{c}^2) + \Delta \mathbf{m}_V^\pm \mathbf{v}^2]^i$$

We suppose that this increment can be defined by gravitational fine structure constant $\beta^i = \left(\frac{\mathbf{m}_0^i}{\mathbf{M}_{Pl}} \right)^2$ different for each lepton generation:

$$\mathbf{E}_\nu^i = \beta^i \mathbf{m}_V^\pm \mathbf{c}^2 = \beta^i [(\mathbf{Rm}_0 \mathbf{c}^2) + \mathbf{m}_V^\pm \mathbf{v}^2]$$

The compensation of the local symmetry shift, related to particles/matter dynamics and transitions by spatially delocalized symmetry shifts in huge number of Bivacuum dipoles around these particles in form of different fields potentials reaction - looks to be a general phenomena. This idea of interconversion between local and non-local effects can be formulated as *symmetry shifts compensation principle* or:

The total sum of symmetry shifts, related to

Matter and Bivacuum transitions in the Universe is zero:

$$\frac{1}{Z} \left[\sum_k^\infty \mathbf{P}_k \Delta(\mathbf{m}_V^+ - \mathbf{m}_V^-)_k \mathbf{c}^2 + \sum_j^\infty \mathbf{P}_j \Delta(\mathbf{m}_V^+ \mathbf{v}^2)_j \right] = 0 \quad (3.91f)$$

where: $\mathbf{Z} = \sum_k^\infty \mathbf{P}_k + \sum_j^\infty \mathbf{P}_j$ is the total partition function, i.e. sum of probabilities of all transitions, conversions, fluctuations of energy in the Universe, including fields: $\Delta(\mathbf{m}_V^+ - \mathbf{m}_V^-)_k \mathbf{c}^2$ and matter: $\Delta(\mathbf{m}_V^+ \mathbf{v}^2)_j$; $\mathbf{m}_V^+$ and $\mathbf{m}_V^-$ are the actual and complementary mass of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of each Bivacuum dipoles and elementary particle in the Universe. The right part of 3.91f in the bracket reflects the changes of energy of matter and the left part - the interrelated changes of energy of secondary Bivacuum. This *Bivacuum symmetry shifts compensation principle* can be considered, as a background for the *energy conservation law*.

*F.5 The Bivacuum dipoles symmetry shift and linear and circular ordering,
as a background of electrostatic and magnetic fields origination*

Let us consider now the origination of electrostatic and magnetic fields, as a consequence of proposed models of elementary particles and their duality. The unified *right* parts of eqs. (3.86) and (3.86a) can be subdivided to translational (*electrostatic*) and rotational (*magnetic*) contributions, determined by corresponding degrees of freedom of Cumulative Virtual Cloud ($\mathbf{CVC}_{tr,rot}^\pm$):

$$\overline{\mathbf{T}}_{tot}^{\mathbf{C}=\mathbf{W}} = \mathbf{T}_{tot} \pm [(\mathbf{E}_E)_{Rec}^{Loc} - (\mathbf{E}_E)_{Rec}^{Dist}]_{tr} \pm \quad (3.92)$$

$$\pm [(\mathbf{E}_H)_{Rec}^{Loc} - (\mathbf{E}_H)_{Rec}^{Dist}]_{rot} \quad (3.92a)$$

where the most probable total kinetic energy of particle can be expressed only via its actual inertial mass ($\mathbf{m}_V^+$) and external velocity ($\mathbf{v}$):

$$\mathbf{T}_{tot} = \frac{1}{2}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 = \frac{1}{2}\mathbf{m}_V^+\mathbf{v}^2 \quad (3.93)$$

The comparison of (3.93) with (3.90) prove that the inertial and gravitational mass are equal.

Formula (3.92-3.92a) reflects the fluctuations of the most probable total kinetic energy, accompanied $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of unpaired sub-elementary fermion, responsible for linear - electrostatic and curled - magnetic fields origination. The more detailed form of (3.92) is:

$$\overline{\mathbf{T}}_{tot}^{\mathbf{C}=\mathbf{W}} = \frac{1}{2}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 \pm \left\{ \left[\frac{|\mathbf{e}_+\mathbf{e}_-|}{\mathbf{L}_T} \right]^{Loc} - \left[\frac{\mathbf{e}^2}{\hbar\mathbf{c}}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 \right]_{tr}^{Dis} \right\} \pm \quad (3.94)$$

$$\pm \left\{ \left[\mathbf{K}_{HE} \frac{|\mu_+\mu_-|}{\mathbf{L}_T} \right]^{Loc} - \left[\mathbf{K}_{HE} \frac{\mu_0^2}{\hbar\mathbf{c}}(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}^2 \right]_{rot}^{Dis} \right\} \quad (3.94a)$$

The oscillation of electrostatic translational contribution, taking into account the obtained relation between mass and charge symmetry shifts (3.35): $\mathbf{m}_V^+ - \mathbf{m}_V^- = \mathbf{m}_V^+ \frac{\mathbf{e}_+^2 - \mathbf{e}_-^2}{\mathbf{e}_+^2}$ can be expressed as:

$$\left[\frac{|\mathbf{e}_+\mathbf{e}_-|}{\mathbf{L}_T} \right]^{Loc} \xrightleftharpoons[\mathbf{W} \rightarrow \mathbf{C}]{\mathbf{C} \rightarrow \mathbf{W}} \left[\alpha \left(\mathbf{m}_V^+ \mathbf{c}^2 \frac{\mathbf{e}_+^2 - \mathbf{e}_-^2}{\mathbf{e}_+^2} \right) \right]_{tr}^{Dis} \quad (3.95)$$

where: $\mathbf{L}_T = \hbar/(\mathbf{m}_V^+ - \mathbf{m}_V^-)\mathbf{c}$ is a characteristic curvature of kinetic energy (3.65b); $|\mathbf{e}_+\mathbf{e}_-| = \mathbf{e}_0^2$ is a rest charge squared; $\alpha = \mathbf{e}^2/\hbar\mathbf{c}$ is the well known dimensionless electromagnetic fine structure constant.

The right part of (3.95) taking into account that: $\mathbf{e}_+^2 - \mathbf{e}_-^2 = (\mathbf{e}_+ - \mathbf{e}_-)(\mathbf{e}_+ + \mathbf{e}_-)$ characterizes the electric dipole moment of triplet, equal to that of unpaired sub-elementary fermion ($\mathbf{F}_{\uparrow}^\pm$).

Like in the case of gravitational potential, it is assumed, that the local *internal* Coulomb potential between opposite charges of torus and antitorus of unpaired sub-elementary fermions (antifermions) $\left(\mathbf{F}_{\downarrow}^{\pm}\right)_{S=\pm 1/2}$ turn reversibly to the *external* distant one due to elastic recoil $\rightleftharpoons$ antirecoil effects, induced by $\mathbf{C} \rightleftharpoons \mathbf{W}$ pulsation of $\left(\mathbf{F}_{\downarrow}^{\pm}\right)_{S=\pm 1/2}$.

The factors, responsible for Coulomb interaction between elementary particles

There are three factors, which determines the attraction or repulsion between opposite or similar elementary charges, correspondingly. They are provided by the recoil $\rightleftharpoons$ antirecoil effects, induced by $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation and emission $\rightleftharpoons$ absorption of positive or negative cumulative virtual clouds CVC^+ or CVC^- of the unpaired sub-elementary fermion $\left(\mathbf{F}_{\downarrow}^{\pm}\right)_{S=\pm 1/2}$ of triplets:

1. The opposite (attraction) or similar (repulsion) Bivacuum dipoles symmetry shift and polarization;
2. Polymerization or depolymerization of Bivacuum dipoles of opposite or similar symmetry/polarization shifts;
3. The different kind of interference of virtual pressure waves of the opposite or similar by sign energy:

$$[\mathbf{VPW}^{\pm} + \mathbf{VPW}^{\mp}] \text{ or } [\mathbf{VPW}^{\pm} + \mathbf{VPW}^{\pm}] \quad (3.96)$$

excited by corresponding cumulative virtual clouds - opposite or similar:

$$[\mathbf{CVC}^{\pm} + \mathbf{CVC}^{\mp}] \text{ or } [\mathbf{CVC}^{\pm} + \mathbf{CVC}^{\pm}] \quad (3.96a)$$

These virtual clouds from subquantum particles, in accordance to our model of duality, are emitted $\rightleftharpoons$ absorbed by the unpaired sub-elementary fermions or/and sub-elementary antifermions in triplets of elementary particles $< [\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]_{S=0} + \left(\mathbf{F}_{\downarrow}^{\pm}\right)_{S=\pm 1/2} >$.

The 1st factor is a basic one. The attraction between opposite charges is a consequence of tendency of Bivacuum to decrease the resulting symmetry shift, decreasing the separation between charges. The repulsion between similar charges also is a consequence of decreasing of resulting symmetry shift, being possible in this case only because of distance increasing between charges. Both of these processes are the consequence of energy conservation law, formulated as eq. 3.91f, i.e. tendency of the Bivacuum symmetry increments to zero.

The 2nd factor is a consequence of the 1st one. The electrostatic Coulomb attraction between opposite electric charges can be a result of Bivacuum bosons $\mathbf{BVB}^\pm = [\mathbf{V}^+ \uparrow\downarrow \mathbf{V}^-]$ or Cooper pairs of Bivacuum dipoles:

$$\{(\mathbf{BVF}^\uparrow = [\mathbf{V}^+ \uparrow\uparrow \mathbf{V}^-]) \bowtie (\mathbf{BVF}^\downarrow = [\mathbf{V}^+ \downarrow\downarrow \mathbf{V}^-])\} \quad (3.96b)$$

polarized by the unpaired sub-elementary fermions of triplets due to charge symmetry shift: This induces the linear assembly of Bivacuum dipoles, their "head to tail" polymerization between $\left(\mathbf{F}_{\uparrow}^\pm\right)_{S=\pm 1/2} >$ of two opposite charges (i.e. electron and proton).

These quasi one-dimensional single $\sum \mathbf{BVB}^\pm$ and twin $\sum(\mathbf{BVF}^\uparrow \bowtie (\mathbf{BVF}^\downarrow)$ virtual microtubules, like 'springs', are responsible for the 'force lines' origination, connecting the opposite charges. The energy of attraction has the following dependence on the length (R) of virtual 'springs': $\vec{r}/R$, where $\vec{r}$ is the unitary radius-vector. On the other hand, the repulsion between similar charges by sign is a result of the opposite phenomena - chaotization of Bivacuum dipoles orientation between such charges. So, the repulsion between similar charges can be explained also in terms of Bivacuum entropy increasing.

The 3d factor is determined by interaction of positive and negative subquantum particles density oscillation, representing virtual pressure waves: $\mathbf{VPW}^+$ and $\mathbf{VPW}^-$. Its effect on attraction or repulsion of charges also can be explained in terms of tending of system: [Charges + Bivacuum] to minimum symmetry shift in space between charges in accordance to energy conservation law in form of eq. 3.91f.

The oscillation of magnetic dipole radiation contribution in the process of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations of sub-elementary fermions between local and distant modes is not accompanied by magnetic moments symmetry shift, but only by the oscillation of separation between torus and antitorus of $\mathbf{BVF}^\uparrow$. It can be described as:

$$\left[\mathbf{K}_{HE}^i \frac{|\mu_+ \mu_-|}{\mathbf{L}_T} \right]^{Loc} \xrightleftharpoons[\mathbf{W} \rightarrow \mathbf{C}]{\mathbf{C} \rightarrow \mathbf{W}} \left[\mathbf{K}_{HE}^i \frac{\mu_0^2}{\hbar c} (\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2 \right]_{rot}^{Dis} \quad (3.97)$$

$$or : \left[\mathbf{K}_{HE}^i \frac{|\mu_+ \mu_-|}{\mathbf{L}_T} \right]^{Loc} \xrightleftharpoons[\mathbf{W} \rightarrow \mathbf{C}]{\mathbf{C} \rightarrow \mathbf{W}} \left[\mathbf{K}_{HE}^i \frac{\mu_0^2}{\hbar c} \mathbf{m}_V^+ \omega_T^2 \mathbf{L}_T^2 \right]_{rot}^{Dis} \quad (3.97a)$$

where: $\frac{\mu_0^2}{\hbar c} = \gamma$ is the magnetic fine structure constant, introduced in our theory. The magneto - electric conversion coefficient $\mathbf{K}_{HE}$ we find from the equality of the electrostatic and magnetic contributions:

$$\mathbf{E}_H = \mathbf{E}_E = \mathbf{T}_{rec} = \frac{1}{2} \frac{e^2}{\hbar c} (\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2 = \frac{1}{2} \mathbf{K}_{HE}^i \frac{\mu_0^2}{\hbar c} \mathbf{m}_V^+ \omega_T^2 \mathbf{L}_T^2 \quad (3.98)$$

These equality is a consequence of equiprobable energy distribution between translational (electrostatic) and rotational (magnetic) independent degrees of freedom of an unpaired sub-elementary fermion and its cumulative virtual cloud (CVC $^\pm$). This becomes evident for the limiting case of photon in vacuum. The sum of these two contributions is equal to

$$\mathbf{E}_H + \mathbf{E}_E = \alpha \mathbf{m}_V^+ \mathbf{v}_{\text{res}}^2 \quad (3.98a)$$

where $\mathbf{v}_{\text{res}}$ is a resulting translational - rotational recoil velocity.

From the above conditions it follows, that:

$$\mathbf{K}_{HE} \frac{\mu_0^2}{\hbar c} = \mathbf{K}_{HE} \frac{\hbar e_0^2}{4\mathbf{m}_0^2 c^3} = \frac{\mathbf{e}_0^2}{\hbar c} \quad (3.99)$$

where $\mu_0^2 = |\mu_+ \mu_-| = \left(\frac{1}{2} e_0 \frac{\hbar}{m_0 c}\right)^2$ is the Bohr magneton.

Consequently, the introduced magneto-electric conversion coefficient is

$$\mathbf{K}_{HE}^{e,p} = \left(\frac{\mathbf{m}_0^{e,p} c}{\hbar/2}\right)^2 = \left(\frac{2}{\mathbf{L}_0^{e,p}}\right)^2 \quad (3.100)$$

where $\mathbf{L}_0^{e,p} = \hbar/\mathbf{m}_0^{e,p} c$ is the Compton radius of the electron or proton.

Origination of magnetic field is assumed to be a result of dynamic equilibrium

$$[\mathbf{BVF}_{S=+1/2}^\uparrow \rightleftharpoons \mathbf{BVB}^\pm \rightleftharpoons \mathbf{BVF}_{S=-1/2}^\downarrow] \quad (3.100b)$$

shift between Bivacuum fermions and Bivacuum antifermions to the left or right, accompanied by corresponding shift of equilibrium between Bivacuum bosons of zero spin and magnetic moment of opposite polarization:

$$\langle \mathbf{BVB}^+ \equiv [\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-] \rangle \rightleftharpoons \langle [\mathbf{V}^+ \downarrow \uparrow \mathbf{V}^-] \equiv \mathbf{BVB}^- \rangle \quad (3.100c)$$

and circulation in the plane, normal to direction of charged particles propagation in the current. It should be related to asymmetric properties of the 'anchor' $\mathbf{BVF}_{anc}^\downarrow$ of unpaired sub-elementary fermion $\left(\mathbf{F}_{\uparrow}^\pm\right)_{S=\pm 1/2} >$ of triplets $< [\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_{S=0} + \left(\mathbf{F}_{\downarrow}^\pm\right)_{S=\pm 1/2} >$ and fast rotation of *pairs* of charge and magnetic dipoles $[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_{S=0}$ in plane, normal to *directed* motion of triplets, i.e. current. This statement is a consequence of the empirical fact, that the magnetic field can be excited only by the electric current: $\vec{\mathbf{j}} = \mathbf{n} e \vec{\mathbf{v}}_j$, i.e. *directed motion* of the charged particles. The cumulative effect of rotation of many of the electrons of current

in plane, normal to current and direction of rotation is determined by the *hand screw rule* and induce the circular structure formation around $\vec{\mathbf{j}}$ in Bivacuum. These axisymmetric structures are the result of 'head-to-tail' vortical assembly of Bivacuum dipoles, i.e. by the same principle as between opposite charges. The vortex - like motion of these axial circled structures along the force lines of the magnetic field occur due to a small symmetry shift between mass and charge of torus and antitorus of $\mathbf{BVF}^\uparrow$ in accordance with (2.11) and (2.13a). This asymmetry is produced by perturbation of Bivacuum dipoles, induced by the unpaired sub-elementary fermions of triplets. The unpaired sub-elementary fermions and antifermions have the opposite influence on symmetry shift between torus and antitorus, interrelated with their opposite influence on the direction of the $[\mathbf{BVF}_{S=+1/2}^\uparrow \rightleftharpoons \mathbf{BVB}^\pm \rightleftharpoons \mathbf{BVF}_{S=-1/2}^\downarrow]$ equilibrium shift.

The equilibrium constant between Bivacuum fermions of opposite spins, characterizing their uncompensated magnetic moment, we introduce, using (3.27), as function of the external translational velocity of $\mathbf{BVF}^\uparrow$:

$$\begin{aligned} \mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow} &= \frac{\mathbf{BVF}_{S=+1/2}^\uparrow}{\mathbf{BVF}_{S=-1/2}^\downarrow} = \exp \left[-\frac{\alpha(\mathbf{m}_V^+ - \mathbf{m}_V^-)}{\mathbf{m}_V^+} \right] = \\ &= \exp \left[-\alpha \frac{\mathbf{v}^2}{\mathbf{c}^2} \right] = \exp \left[-\frac{\omega_T^2 \mathbf{L}_T^2}{\mathbf{c}^2} \right] \quad (3.101) \end{aligned}$$

The similar values of $\mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow}$ have the axial distribution with respect to the current vector ($\mathbf{j}$) of charges. The conversion of Bivacuum fermions or Bivacuum antifermions to Bivacuum bosons ($\mathbf{BVB}^\pm = \mathbf{V}^+ \updownarrow \mathbf{V}^-$) with different probability, also may provide an increasing or decreasing of the equilibrium constant $\mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow}$. The corresponding probability difference is dependent on the direction of current related, in-turn, with direction of unpaired sub-elementary fermion circulation, affecting in opposite way on the angular momentum and stability of $\mathbf{BVF}_{S=+1/2}^\uparrow$ and $\mathbf{BVF}_{S=-1/2}^\downarrow$.

The magnetic field tension can be presented as a gradient of the constant of equilibrium:

$$\mathbf{H} = \mathbf{grad} (\mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow}) = (\vec{r}/R) \mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow} \quad (3.102)$$

The chaotic thermal velocity of the 'free' conductivity electrons in metals and ions at room temperature is very high even in the absence of current, as determined by Maxwell-Boltzmann distribution:

$$\mathbf{v}_T = \sqrt{\frac{\mathbf{kT}}{\mathbf{m}_V^+}} \sim 10^7 \text{ cm/s} \quad (3.103)$$

It proves, that not the acceleration, but the ordering of the electron translational and rotational dynamics in space, provided by current, is a main reason of the curled magnetic field excitation. In contrast to conventional view, the electric current itself is not a *primary*, but only a *secondary* reason of magnetic field origination, as the charges translational and rotational dynamics 'vectorization factor'.

*F6. Interpretation of the Maxwell displacement current,
based on Bivacuum model*

The magnetic field origination in Bivacuum can be analyzed also from more conventional point of view.

Let us analyze the 1st Maxwell equation, interrelating the circulation of vector of magnetic field tension $\mathbf{H}$ along the closed contour $\mathbf{L}$ with the conduction current ($\mathbf{j}$) and *displacement current* $\mathbf{j}_d = \frac{1}{4\pi} \frac{\partial \mathbf{E}}{\partial t}$ through the surface, limited by $\mathbf{L}$:

$$\oint_{\mathbf{L}} \mathbf{H} d\mathbf{l} = \frac{4\pi}{c} \int_{\mathbf{S}} \left(\mathbf{j} + \frac{1}{4\pi} \frac{\partial \mathbf{E}}{\partial t} \right) d\mathbf{s} \quad (3.104)$$

where ($\mathbf{s}$) is the element of surface, limited with contour ($\mathbf{l}$).

The existence of the displacement current: $\mathbf{j}_d = \frac{1}{4\pi} \frac{\partial \mathbf{E}}{\partial t}$ is in accordance with our model of Bivacuum, as the oscillating virtual dipoles ($\text{BVF}^\dagger$ and $\text{BVB}^\pm$) continuum.

In condition of *primordial* Bivacuum of the ideal virtual dipoles symmetry (i.e. in the absence of matter and fields) these charges totally compensate each other.

However, even in primordial symmetric Bivacuum the oscillations of distance and energy gap between torus and antitorus of Bivacuum dipoles is responsible for *displacement current*. This alternating current generates corresponding *displacement magnetic field*.

Corresponding virtual dipole oscillations are the consequence of the in-phase transitions of $\mathbf{V}^+$ and $\mathbf{V}^-$ between the excited and ground states, compensating each other. These transitions are accompanied by spontaneous emission and absorption of positive and negative virtual pressure waves: $\mathbf{VPW}^+$ and $\mathbf{VPW}^-$.

The primordial displacement current and corresponding displacement magnetic field can be enhanced by presence of matter. It is a consequence of fluctuations of potential energy of sub-elementary fermions of triplets, accompanied by excitation of positive and negative virtual pressure waves, which in accordance to our approach, is interrelated with gravitational field of particles (see 3.88 - 3.90).

F7. New kind of current in secondary Bivacuum, additional to displacement one

This additional current is a consequence of vibrations of $\mathbf{BVF}^\dagger$, induced by recoil-antirecoil effects, accompanied $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ transitions of unpaired sub-elementary fermion of triplets $\langle [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]_{S=0} + (\mathbf{F}_\uparrow^+)_{S=\pm 1/2} \rangle^{e,\tau}$.

The corresponding elastic deformations of Bivacuum fermions $(\mathbf{BVF}^\dagger) \equiv [\mathbf{V}^+ \updownarrow \mathbf{V}^-]$ are followed by small charge-dipole symmetry zero-point oscillations ($\mathbf{v}^{ext} = 0$) with amplitude, determined by the most probable resulting translational - rotational recoil velocity ($\mathbf{v}_{rec}$). At conditions $\mathbf{e}_+ \simeq \mathbf{e}_- \simeq \mathbf{e}_0$ and $|\mathbf{e}_+ - \mathbf{e}_-| \ll \mathbf{e}_0$, i.e. at small perturbations of torus and antitorus: $\mathbf{V}^+$ and $\mathbf{V}^-$ we have for the charge symmetry shift oscillation amplitude:

$$\Delta \mathbf{e}_\pm = \mathbf{e}_+ - \mathbf{e}_- = \frac{1}{2} \mathbf{e}_0 \frac{\mathbf{v}_{rec}^2}{\mathbf{c}^2} \quad (3.105)$$

The resulting most probable recoil velocity using (3.98) can be defined from two expressions of recoil momentum, standing for electromagnetism:

$$\mathbf{T}_{rec} = \frac{1}{2} \alpha (\mathbf{m}_\mathbf{V}^+ - \mathbf{m}_\mathbf{V}^-) \mathbf{c}^2 = \frac{1}{2} \alpha \mathbf{m}_\mathbf{V}^+ \mathbf{v}_{res}^2 \quad (3.106)$$

$$\mathbf{v}_{rec}^2 = \alpha \mathbf{v}_{res}^2 \quad (3.106a)$$

The minimum value of recoil velocity, corresponding to zero *external* translational velocity of triplets and internal velocity, determined by Golden mean conditions $(\mathbf{v}_{res}/\mathbf{c})^2 = \phi = 0.61803398$, can be considered as the *velocity of zero-point oscillations*:

$$(\mathbf{v}_{rec}^2)^{\min} \equiv (\mathbf{v}_0^2)_{HE}^{\min} = \alpha \phi \mathbf{c}^2 \quad (3.107)$$

where: $\alpha = e^2/\hbar c = 0,0072973506$; $\alpha \phi = (\mathbf{v}_{rec}^2)^{\min}/\mathbf{c}^2 = 4.51 \cdot 10^{-3}$.

The alternating *recoil current* (j_{rec}^{EH}) , additional to that of Maxwell *displacement current* (j_d) , existing in presence of charged particles even in the absence of conducting current ($\mathbf{j} = \mathbf{0}$) is equal to product of (3.105) and square root of (3.107). At Golden mean conditions $(\mathbf{v}/\mathbf{c})^2 = \phi$ this *recoil current* is:

$$(\mathbf{j}_{rec}^\phi)^{EH} = (\Delta \mathbf{e}_\pm)^\phi (\mathbf{v}_{rec})^{\min} = \frac{1}{2} \alpha^{1/2} \phi^{3/2} \mathbf{e}_0 \mathbf{c} \quad (3.108)$$

Corresponding gravitational contribution of recoil velocity, related to vibration of potential energy of particle (3.90) is much smaller, as far $\beta \ll \alpha$:

$$\mathbf{V}_{rec} = \frac{1}{2} \beta (\mathbf{m}_\mathbf{V}^+ + \mathbf{m}_\mathbf{V}^-) \mathbf{c}^2 = \frac{1}{2} \beta \mathbf{m}_\mathbf{V}^+ \mathbf{c}^2 (2 - \mathbf{v}^2/\mathbf{c}^2) \quad (3.109)$$

The zero-point characteristic potential recoil velocity squared at GM conditions $(\mathbf{v}^2/\mathbf{c}^2)^\phi = \phi$ is:

$$\begin{aligned} (\mathbf{v}_0^2)_G &= \beta \mathbf{c}^2 (2 - \phi); & (\mathbf{v}_0^2)_G / \mathbf{c}^2 &= \beta (2 - \phi) \\ (\mathbf{v}_0)_G &= \mathbf{c} \beta^{1/2} (2 - \phi)^{1/2} = 1,446 \cdot 10^{-12} \text{ cm/s} \end{aligned} \quad (3.109a)$$

Consequently, the Maxwell equation (3.104) can be modified, taking into account the EH recoil current, as

$$\oint_{\mathbf{L}} \mathbf{H} d\mathbf{l} = \frac{4\pi}{c} \int_{\mathbf{S}} \left(\mathbf{j} + \frac{1}{4\pi} \frac{\partial \mathbf{E}}{\partial t} + \mathbf{j}_{rec}^{EH} \right) d\mathbf{s} = \mathbf{I}_{tot} \quad (3.110)$$

where: $\mathbf{I}_{tot}$ is total current via surface ($\mathbf{S}$).

We have to note, that $\mathbf{j}_{rec}^{EH}$ is nonzero not only in the vicinity of particles, but as well in any remote space regions of Bivacuum, perturbed by electric and magnetic potentials. This consequence of our theory coincides with the extended electromagnetic theory of Bo Lehnert (section II of this paper), also considering current in vacuum, additional to displacement one.

In accordance with the known Helmholtz theorem, each kind of vector field ($\mathbf{F}$), tending to zero at infinity, can be presented, as a sum of the gradient of some scalar potential (ϕ) and a rotor of vector potential ($\mathbf{A}$):

$$\mathbf{F} = \mathbf{grad} \phi + \mathbf{rot} \mathbf{A} \quad (3.111)$$

The scalar and vector potentials are convenient to use for description of electromagnetic field, i.e. photon properties. They are characterized by the interrelated translational and rotational degrees of freedom, indeed (Fig.3.2).

To explain the *ability of secondary Bivacuum to keep the average (macroscopic) mass and charge equal to zero*, we have to postulate, that the mass and charge symmetry shifts oscillations of Bivacuum fermions and antifermions, forming virtual Cooper pairs:

$$(\mathbf{BVF}^\dagger)_{S=+1/2}^\pm \equiv [\mathbf{V}^+ \uparrow \uparrow \mathbf{V}^-]^\pm \propto [\mathbf{V}^+ \downarrow \downarrow \mathbf{V}^-]^\mp \equiv (\mathbf{BVF}^\dagger)_{S=-1/2}^\mp \quad (3.112)$$

are opposite by sign, but equal by the absolute value. Consequently, the polarized secondary Bivacuum (i.e. perturbed by matter and field) can be considered, as a *plasma of the in-phase oscillating virtual dipoles (BVF)* of opposite resulting charge and mass/energy.

F8. The mechanism, increasing the refraction index of Bivacuum

By definition, the *torus* is a figure, formed by rotation of a circle with maximum radius, corresponding to minimum quantum number $\mathbf{n} = \mathbf{0}$ (see 3.1a) $\mathbf{L}_{\mathbf{V}\pm}^i = \frac{2\hbar}{\mathbf{m}_0^i \mathbf{c}}$, around the axis, shifted from the center of the circle at the distance $\pm \Delta \mathbf{L}_{EH,G}$. The electromagnetic (*EH*) and gravitational (*G*) vibrations of positions $(\pm \Delta \mathbf{L}_{EH,G})_{V\pm}$ of the big number of recoiled $\mathbf{BVF}_{rec}$, induced by the elastic recoil $\Rightarrow$ antirecoil deformations of Bivacuum matrix, are accompanied by vibrations of square and volume of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of perturbed Bivacuum dipoles: $(\mathbf{BVF}_{rec}^\dagger)^i = [\mathbf{V}^+ \Updownarrow \mathbf{V}^-]_{rec}^i$. The electromagnetic and gravitational increments of square ($\Delta \mathbf{S}_{\mathbf{V}\pm}^{E,G}$) and volume ($\Delta \mathbf{V}_{\mathbf{V}\pm}^{E,G}$) of toruses and antitoruses of $(\mathbf{BVF}_{rec}^\dagger)^i$, as a consequence of their center vibrations can be presented, correspondingly, as:

$$\Delta \mathbf{S}_{\mathbf{V}\pm}^{E,G} = 4\pi^2 |\Delta \mathbf{L}_{EH,G}|_{V\pm}^{E,G} \cdot \mathbf{L}_{\mathbf{V}\pm} \quad (3.113)$$

$$\Delta \mathbf{V}_{\mathbf{V}\pm}^{E,G} = 4\pi^2 |\Delta \mathbf{L}_{EH,G}|_{V\pm}^{E,G} \cdot \mathbf{L}_{\mathbf{V}\pm}^2 \quad (3.113a)$$

At conditions of zero-point oscillations, corresponding to Golden Mean (GM), when the ratio $(\mathbf{v}_0/\mathbf{c})^2 = \phi$ and external translational velocity ($\mathbf{v}$) is zero, the maximum shifts of center of secondary Bivacuum dipoles *in vicinity of pulsing elementary particles* due to electromagnetic and gravitational recoil-antirecoil (zero-point) vibrations are, correspondingly:

$$(\Delta \mathbf{L}_{\mathbf{EH}}^i)_{V\pm}^\phi = \left(\tau_{C\Rightarrow W}^\phi \mathbf{v}_{EH}^\phi \right)^i = \frac{\hbar}{\mathbf{m}_0^i \mathbf{c}} (\alpha\phi)^{1/2} = 0,067 (\mathbf{L}_{\mathbf{V}\pm}^i) \quad (3.114)$$

$$(\Delta \mathbf{L}_{\mathbf{G}}^i)_{V\pm}^\phi = \left(\tau_{C\Rightarrow W}^\phi \mathbf{v}_G^\phi \right)^i = \frac{\hbar}{\mathbf{m}_0^i \mathbf{c}} \beta^{1/2} (2 - \phi)^{1/2} = 3,27 \cdot 10^{-23} (\mathbf{L}_{\mathbf{V}\pm}^i) \quad (3.114a)$$

where: the recoil $\Leftrightarrow$ antirecoil oscillation period is $\left[\tau_{C\Rightarrow W}^\phi = 1/\omega_{C\Rightarrow W}^\phi = \hbar/\mathbf{m}_0^i \mathbf{c}^2 \right]^i$; the recoil $\Rightarrow$ antirecoil most probable velocity of zero-point oscillations, which determines the electrostatic and magnetic fields is: $\mathbf{v}_{EH}^\phi = \mathbf{c}(\alpha\phi)^{1/2} = 0,201330447 \times 10^8 \text{ m s}^{-1}$ and $(\alpha\phi)^{1/2} = 0,067$ the corresponding zero-point velocity, which determines gravitational field is: $\mathbf{v}_G^\phi = \mathbf{c} \beta_e^{1/2} (2 - \phi)^{1/2} = 1,446 \cdot 10^{-12} \text{ m s}^{-1}$ and $\beta_e^{1/2} (2 - \phi)^{1/2} = 0,48 \cdot 10^{-22}$.

The dielectric permittivity of Bivacuum and corresponding refraction index, using our theory of refraction index of matter [63, 64], can be presented as a ratio of volume of Bivacuum fermions and bosons in symmetric *primordial* Bivacuum ($\mathbf{V}_{pr}$) to their volume in *secondary* Bivacuum: $\mathbf{V}_{sec} = \mathbf{V}_{BVF} - (\mathbf{r}/r) \Delta \mathbf{V}_{\mathbf{BVF}_{rec}}^{E,G}$, perturbed by matter and fields. The

secondary Bivacuum is optically more dense, if we assume that the volume, occupied by Bivacuum fermion torus and antitorus, is excluded for photons. The Coulomb and gravitational potentials and the related excluded volumes of perturbed Bivacuum fermions/antifermions decline with distance (r) as:

$$(\vec{\mathbf{r}}/r)\Delta\mathbf{V}_{\mathbf{BVF}_{rec}}^{EH} \quad \text{and} \quad (\vec{\mathbf{r}}/r)\Delta\mathbf{V}_{\mathbf{BVF}_{rec}}^G \quad (3.115)$$

where: (r) is a distance from the charged and/or gravitating particle and $\vec{\mathbf{r}}$ is the unitary radius vector. Taking all this into account, we get for permittivity of secondary Bivacuum:

$$[c]c\varepsilon = \mathbf{n}^2 = \left(\frac{\mathbf{c}}{\mathbf{v}_{EH,G}}\right)^2 = \frac{N \mathbf{V}_{\mathbf{pr}}}{N \mathbf{V}_{\text{sec}}} = \frac{\mathbf{V}_{BVF}}{\mathbf{V}_{BVF} - (\mathbf{r}/r)\Delta\mathbf{V}_{\mathbf{BVF}_{rec}}^{EH,G}} = \frac{1}{(1 - \mathbf{r}/r)\Delta\mathbf{V}_{\mathbf{BVF}_{rec}}^{EH,G}/\mathbf{V}_{BVF}} \quad (3.116)$$

$$\mathbf{n}^2 = \frac{1}{1 - (\mathbf{r}/r) \frac{3\pi |\Delta\mathbf{L}_{V\pm}^{EH,G} \cdot \mathbf{L}_{V\pm}}{V_{BVF}}} \quad (3.116a)$$

where: the velocity of light propagation in asymmetric secondary Bivacuum of higher virtual density, than in primordial one, is notated as: $\mathbf{v}_{EH,G} = \mathbf{c}_{EH,G}$; the volume of primordial Bivacuum fermion is $\mathbf{V}_{BVF} = (4/3)\pi\mathbf{L}_{V\pm}^3$ and its increment in secondary Bivacuum: $\Delta\mathbf{V}_{\mathbf{BVF}_{rec}}^{E,G} = \Delta\mathbf{V}_{\mathbf{V}\pm}^{E,G}$ (3.113a).

$(\mathbf{r}/r)$ is a ratio of unitary radius-vector to distance between the source of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations (elementary particle) and perturbed by the electrostatic, magnetic and gravitational potential $\mathbf{BVF}_{rec}^{EH,G}$.

Putting (3.114) into formula (3.116a) we get for the refraction index of Bivacuum and relativistic factor ($\mathbf{R}_E$) in the vicinity of charged elementary particle (electron, positron or proton, antiproton) the following expression:

$$\left[\varepsilon = \mathbf{n}^2 = \left(\frac{\mathbf{c}}{\mathbf{c}_{EH}}\right)^2\right]_E = \frac{1}{1 - (\mathbf{r}/r) \frac{3\pi(\alpha\phi)^{1/2}}{V_{BVF}}} \lesssim 2.71 \quad (3.117)$$

where: $1 \lesssim \mathbf{n}^2 \lesssim 2,71$ is tending to 1 at $r \rightarrow \infty$.

The Coulomb relativistic factor:

$$\mathbf{R}_{EH} = \sqrt{1 - \frac{(\mathbf{c}_{EH})^2}{\mathbf{c}^2}} = \sqrt{(\mathbf{r}/r) \frac{0,631}{V_{BVF}}} \lesssim (\mathbf{r}/r)^{1/2} \frac{0,794}{V_{BVF}} \quad (3.118)$$

$0 \lesssim \mathbf{R}_E \lesssim 0,794$ is tending to zero at $r \rightarrow \infty$.

In similar way, using (3.114a) and (3.116a), for the refraction index of Bivacuum and the corresponding relativistic factor ($\mathbf{R}_G$) of gravitational vibrations of Bivacuum fermions ($\mathbf{BVF}^\dagger$) in the vicinity of pulsing elementary particles at zero-point conditions, we get:

$$\left[\varepsilon = \mathbf{n}^2 = \left(\frac{\mathbf{c}_G}{\mathbf{c}} \right)^2 \right]_G = \frac{1}{1 - (\mathbf{r}/r) 3\pi(\beta^e)^{1/2}(2-\phi)^{1/2}} \approx 1 \quad (3.119)$$

where $(\beta^e)^{1/2}(2-\phi)^{1/2} = 0.48 \times 10^{-22}$.

The gravitational relativistic factor:

$$\mathbf{R}_G = \sqrt{1 - \left(\frac{\mathbf{c}_G}{\mathbf{c}} \right)^2} = \sqrt{(\mathbf{r}/r) 0,48 \cdot 10^{-22}} \lesssim (\mathbf{r}/r)^{1/2} 0,69 \cdot 10^{-11} \quad (3.120)$$

Like in previous case, the Bivacuum refraction index, increased by gravitational potential, is tending to its minimum value: $\mathbf{n}^2 \rightarrow 1$ at the increasing distance from the source: $r \rightarrow \infty$.

The charge - induced refraction index increasing of secondary Bivacuum, in contrast to the mass - induced one, is independent of lepton generations of Bivacuum dipoles (e, μ, τ).

The formulas (3.117) and (3.119) for Bivacuum dielectric permittivity and refraction index near elementary particles, perturbed by their Coulomb and gravitational potentials, point out that bending and scattering probability of photons on charged particles is much higher, than that on neutral particles with similar mass.

We have to point out, that the *light velocity* in conditions: $\left[\mathbf{n}_{EH,G}^2 = \mathbf{c}/\mathbf{v}_{EH,G} = \mathbf{c}/\mathbf{c}_{EH,G} \right] > 1$ is not longer a scalar, but a vector, determined by the gradient of Bivacuum fermion symmetry shift:

$$grad \Delta \left| \mathbf{m}_V^+ - \mathbf{m}_V^- \right|_{EH,G} \mathbf{c}^2 = grad \Delta \left(\mathbf{m}_V^+ \mathbf{v}^2 \right) \quad (3.121)$$

and corresponding gradient of torus and antitorus equilibrium constant increment:

$$\Delta \mathbf{K}_{\mathbf{V}+\uparrow \mathbf{V}-} = \mathbf{1} - \mathbf{m}_V^-/\mathbf{m}_V^+ = (\mathbf{c}_{EH,G}/\mathbf{v})^2:$$

$$grad[\Delta \mathbf{K}_{\mathbf{V}+\uparrow \mathbf{V}-} = \mathbf{1} - \mathbf{m}_V^-/\mathbf{m}_V^+] = \quad (3.122)$$

$$= grad \left(\frac{\mathbf{c}_{EH,G}}{\mathbf{c}} \right)^2 = grad \frac{1}{\mathbf{n}^2} \quad (3.122a)$$

The other important consequence of: $[\mathbf{n}^2]_{E,G} > 1$ is that *the contributions of the rest mass energy of photons and neutrino to their total energy is not zero*, as far the electromagnetic

and gravitational relativistic factors ($\mathbf{R}_{EH,G}$) are greater than zero. It follows from the basic formula for the total energy of de Broglie wave (the photon in our case):

$$\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \hbar \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} = \mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in} + (\hbar \omega_B^{ext})_{tr} \quad (3.123)$$

where the gravitational relativistic factor of electrically neutral objects: $\mathbf{R}_G = \sqrt{(\mathbf{r}/r) 3,08 \cdot 10^{-22}} \lesssim (\mathbf{r}/r)^{1/2} 1.75 \times 10^{-11}$.

This consequence is also consistent with the models of the photon and neutrino, developed by Bo Lehnert (see section II of this paper).

We can see, that in conditions of *primordial* Bivacuum, when $r \rightarrow \infty$, the $\mathbf{n}_{EH,G} \rightarrow 1$, $\mathbf{R}_{EH,G} \rightarrow 0$ and the contribution of the rest mass energy $\mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in}$ tends to zero. At these limiting conditions the frequency of photon [*Corpuscle* $\rightleftharpoons$ *Wave*] pulsation is equal to the frequency of the photon as a wave:

$$\mathbf{E}_{ph} = \hbar \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} = \hbar \omega_{ph} = h \frac{\mathbf{c}}{\lambda_{ph}} \quad (3.124)$$

The results of our analysis explain the bending of light beams, under the influence of strong gravitational potential in another way, than by Einstein's general theory of relativity. A similar idea of polarizable vacuum and its permittivity variations has been developed by Dicke [65], Fock [32] and Puthoff [31], as a background of 'vacuum engineering'.

For the spherically symmetric star or planet it was shown using Dicke model, that the dielectric constant $\mathbf{K}$ of polarizable vacuum is given by the exponential form:

$$\mathbf{K} = \exp(2\mathbf{G}\mathbf{M}/\mathbf{r}\mathbf{c}^2) \quad (3.125)$$

where $\mathbf{G}$ is the gravitational constant, $\mathbf{M}$ is the mass, and $\mathbf{r}$ is the distance from the mass center.

For comparison with expressions derived by conventional General Relativity techniques, the following approximation of the formula above is sufficient [31]:

$$\mathbf{K} \approx 1 + \frac{2\mathbf{G}\mathbf{M}}{\mathbf{r}\mathbf{c}^2} + \frac{1}{2} \left(\frac{2\mathbf{G}\mathbf{M}}{\mathbf{r}\mathbf{c}^2} \right)^2 \quad (3.126)$$

Our approach propose the concrete mechanism of Bivacuum optical density increasing near charged and gravitating particles, inducing light beams bending.

**G. The Principle of least action, as a consequence of Bivacuum
basic Virtual Pressure Waves (VPW_{q=1}[±]) resonance interaction with particles.
Possible source of energy for overunity devices.**

Let us analyze the Maupertuis-Lagrange formula for *action*:

$$\mathbf{S}_{ext} = \int_{t_0}^{t_1} 2\mathbf{T}_k^{ext} \cdot d\mathbf{t} \quad (3.127)$$

The *principle of Least action*, choosing one of few possible trajectories of system changes from one configuration to another at the permanent total energy of a system of elementary particles: $\mathbf{E}_{tot} = \mathbf{m}_V^+ \mathbf{c}^2 = \text{const}$ has a form:

$$\Delta \mathbf{S}_{ext} = 0 \quad (3.128)$$

This means, that the optimal trajectory of particle corresponds to minimum variations of the external energy of it wave B.

The time interval: $\mathbf{t} = \mathbf{t}_1 - \mathbf{t}_2 = \mathbf{n} \mathbf{t}_B$ is equal or bigger than the period of the external (modulation) frequency of wave B ($\mathbf{t}_B = 1/\nu_B$):

$$\mathbf{t} = \mathbf{t}_1 - \mathbf{t}_2 = \mathbf{n} \mathbf{t}_B = \mathbf{n} / \nu_B \quad (3.129)$$

Using eqs.(3.127 and 3.74; 3.74a), we get for the action:

$$\mathbf{S}_{ext} = 2\mathbf{T}_k^{ext} \cdot \mathbf{t} = \mathbf{m}_V^+ \mathbf{v}^2 \cdot \mathbf{t} = (\mathbf{1} + \mathbf{R})[\mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{m}_0 \mathbf{c}^2]^i \cdot \mathbf{t} = \quad (3.130)$$

$$\text{or : } \mathbf{S}_{ext} = \mathbf{m}_V^+ \mathbf{v}^2 \cdot \mathbf{t} = (\mathbf{1} + \mathbf{R}) \mathbf{TE} \cdot \mathbf{t} \quad (3.130a)$$

where relativistic factor: $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ and $(\mathbf{v})$ is the resulting external translational velocity.

We introduce here the important notions of *Tuning energy* of Bivacuum and corresponding frequency difference, as:

$$\mathbf{TE} = \mathbf{E}_{tot} - \mathbf{E}_{rot}^{in} = [\mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{q} \mathbf{m}_0 \mathbf{c}^2]^i = \hbar \omega_{\mathbf{TE}} = \hbar [\omega_{\mathbf{C} \Rightarrow \mathbf{W}} - \mathbf{q} \omega_0]^i = \frac{\mathbf{m}_V^+ \mathbf{v}^2}{\mathbf{1} + \mathbf{R}} \quad (3.131)$$

where: $\mathbf{q} = \mathbf{j} - \mathbf{k}$ is a quantum number characterizing excitation of Bivacuum virtual pressure (VPW_q[±]) and spin (VirSW_q) waves (see section 3.1.2).

The frequency (ω_{TE}^i) of Bivacuum Tuning energy is defined as:

$$\omega_{TE} = (\mathbf{m}_V^+ - \mathbf{q} \mathbf{m}_0)^i \mathbf{c}^2 / \hbar = [\omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} - \mathbf{q} \omega_0^i] \quad (3.132)$$

The influence of Tuning energy of Bivacuum on matter is a result of *forced combinational resonance* between virtual pressure waves ($\mathbf{VPW}_q^\pm$) of Bivacuum with quantized frequency (3.10a):

$$\omega_{\mathbf{VPW}_q^\pm}^i = \mathbf{q} \omega_0^i = \mathbf{q} \mathbf{m}_0^i \mathbf{c}^2 / \hbar \quad (3.133)$$

and $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of elementary particles $\omega_{\mathbf{C} \rightleftharpoons \mathbf{W}}$.

The condition of combinational resonance is:

$$\omega_{\mathbf{VPW}_q}^i = q \omega_0^i = \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} \quad (3.134)$$

$$\text{or : } \mathbf{E}_{VPW} = \mathbf{q} \mathbf{m}_0^i \mathbf{c}^2 = \mathbf{m}_V^+ \mathbf{c}^2 \quad (3.134a)$$

The energy exchange between $\mathbf{VPW}_q^+ + \mathbf{VPW}_q^-$ and real particles in the process of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of pairs $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_{x,y}$ of real fermions - triplets $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_{x,y} + \mathbf{F}_\uparrow^\pm >_z^i$ at *pull-in -range* state accelerate them, if $\mathbf{q} = \mathbf{2}, \mathbf{3}, \mathbf{4}$, i.e. bigger than $\mathbf{1}$, driving to resonant conditions (3.134). On the other hand, at the minimum energy of $\mathbf{VPW}_{q=1}^\pm$, when $\mathbf{q} = \mathbf{j} - \mathbf{k} = \mathbf{1}$, their interaction with triplets slow the translational velocity of particles down and the deceleration (cooling) effect takes a place.

In accordance to rules of combinational resonance of Bivacuum virtual pressure waves with elementary particles, we have the following relation between quantized energy and frequency of $\mathbf{VPW}_q^\pm$ and energy/frequency of triplets $\mathbf{C} \rightleftharpoons \mathbf{W}$ pulsation in resonance conditions:

$$\mathbf{E}_{\mathbf{VPW}^\pm} = \hbar \omega_{\mathbf{VPW}^\pm}^i = \mathbf{q} \hbar \omega_0^i = \hbar \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}}^i = \mathbf{R} \hbar \omega_0 + \hbar \omega_B \quad (3.135)$$

$$\mathbf{E}_{\mathbf{VPW}^\pm} = \mathbf{q} \mathbf{m}_0^i \mathbf{c}^2 = \mathbf{R} \mathbf{m}_0^i \mathbf{c}^2 + \mathbf{m}_V^+ \mathbf{v}^2$$

$$\text{or : } \mathbf{E}_{\mathbf{VPW}^\pm} = \mathbf{R} \mathbf{m}_0^i \mathbf{c}^2 + \frac{\mathbf{m}_0^i \mathbf{c}^2 (\mathbf{v}/\mathbf{c})^2}{\mathbf{R}} \quad (3.136)$$

$$\text{or : } \mathbf{q} = \mathbf{R} + \frac{(\mathbf{v}/\mathbf{c})^2}{\mathbf{R}} \quad (3.136a)$$

where the $\mathbf{VPW}_q^\pm$ quantum number: $\mathbf{q} = \mathbf{j} - \mathbf{k} = 1, 2, 3 \dots$ (integer numbers)

The angle frequency of de Broglie waves of particles $(\omega_B)_{1,2,3}$, is dependent on the external translational velocity $(\mathbf{v})_{1,2,3}$:

$$(\omega_B = \hbar/2\mathbf{m}_V^+ \mathbf{L}_B^2 = \mathbf{m}_V^+ \mathbf{v}^2/2\hbar)_{1,2,3} \quad (3.137)$$

The important relation between the translational most probable velocity of particle $(\mathbf{v})$ and quantization number $(\mathbf{q})$, corresponding to resonant interaction of Bivacuum $\mathbf{VPW}_q^\pm$ with pulsing particles, derived from (3.136) is:

$$\mathbf{v} = \mathbf{c} \left(\frac{\mathbf{q}^2 - 1}{\mathbf{q}^2} \right)^{1/2} \quad (3.138)$$

At the conditions of triplets fusion, when $\mathbf{q} = 1$, the resonant conditions correspond to *translational* velocity of particle equal to zero: $\mathbf{v}_{n=1} = \mathbf{0}$.

When the quantized energy of $\mathbf{E}_{\mathbf{VPW}_n^\pm} = \mathbf{q} \mathbf{m}_0^i \mathbf{c}^2$ corresponds to $\mathbf{q} = 2$, the resonant translational velocity of particle from (3.138) should be: $\mathbf{v}_{q=2} = \mathbf{c} \times 0.866 = 2,6 \times 10^{10} \text{cm/s}$.

At $\mathbf{q} = 3$, we have from (3.138) for resonant velocity: $\mathbf{v}_{q=3} = \mathbf{c} \times 0.942 = 2.83 \times 10^{10} \text{cm/s}$.

It is natural to assume, that if the velocity of particles $(\mathbf{v})$, corresponds to $\mathbf{q} < 1.5$, the interaction of these pulsating particles with basic $\mathbf{VPW}_{n=1}^\pm$ should slow down their velocity, driving it to resonant conditions: $\mathbf{q} = 1$, $\mathbf{v} \rightarrow \mathbf{0}$. *The 2nd and 3d laws of thermodynamics for the closed systems, reflecting the 'spontaneous' cooling of matter and tending the entropy to zero [66], can be a consequence of just this condition.*

On the other hand, if velocity of particles is high enough and corresponds to $\mathbf{q} > 1.5$ in (3.138), their pull-in range interaction with excited $\mathbf{VPW}_{n=2}^\pm$ can accelerate them up to conditions: $\mathbf{q} = 2$, $\mathbf{v} \rightarrow 2.6 \times 10^{10} \text{cm/s}$. If the starting particles velocity corresponds to $\mathbf{q} > 2.5$, their forced resonance with even more excited $\mathbf{VPW}_{n=3}^\pm$ should accelerate them up to conditions: $\mathbf{q} = 3$, corresponding to $\mathbf{v} = 2.83 \times 10^{10} \text{cm/s}$. The described mechanism of Bivacuum - Matter interaction, can be a general physical background of all kinds of *overunity devices* [21, 67]. The coherent electrons and protons of hot enough plasma in stars and in artificial conditions also may get the additional energy from high-frequency virtual pressure waves of Bivacuum $\mathbf{VPW}_{n=2,3,..}^\pm$, excited by strong gravitational and/or magnetic fields.

We can see from the formulas above, that the action of Bivacuum Tuning energy due to interaction of Bivacuum low frequency $\mathbf{VPW}_{n=1}^\pm$ with particles, is responsible for real-

ization of fundamental principle of Least action: $\Delta S_{ext} \rightarrow 0$ at $\mathbf{TE} \rightarrow 0$, corresponding to minimization of translational kinetic energy of particles $\mathbf{v}_{tr}^{ext} \rightarrow 0$ at $\mathbf{q} \rightarrow 1$.

The action of $\mathbf{TE}$ on virtual Cooper pairs of sub-elementary fermions is opposite to that on real particles. It increases the velocity and kinetic energy of virtual particles and finally may turn them to real ones. Such a mechanism of real particles (like photon) origination may work in the process of photons emission by excited atoms, molecules and accelerated charges.

The second law of thermodynamics, formulated as a spontaneous irreversible transferring of the heat energy from the warm body to the cooler body or surrounding medium, also means slowing down the kinetic energy of particles, composing this body. Consequently, the 2nd law of thermodynamics, as well as Principle of least action, can be a consequence of Tuning energy ($\mathbf{TE}$) minimization, slowing down particles thermal *translational* dynamics at pull-in range conditions:

$$\mathbf{TE} = \hbar(\omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} \rightarrow \omega_0)^{\mathbf{v} \rightarrow 0} \mathbf{0} \quad \text{at} \quad (\mathbf{1}, \mathbf{5} < \mathbf{q}) \xrightarrow{\mathbf{v} \rightarrow 0} (\mathbf{q} = \mathbf{1}) \quad (3.139)$$

The third law of thermodynamics states, that the entropy of equilibrium system is tending to zero at the absolute temperature close to zero. Again, this may be a consequence of forced combinational resonance between basic $\mathbf{VPW}_{n=1}^{\pm}$ and particles $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation, when *translational* velocity of particles $\mathbf{v} \rightarrow 0$ and $\mathbf{TE} = \hbar(\omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} \rightarrow \omega_0)^{\mathbf{v} \rightarrow 0} \mathbf{0}$ at $(\mathbf{q} < \mathbf{1}, \mathbf{5}) \xrightarrow{\mathbf{v} \rightarrow 0} (\mathbf{q} = \mathbf{1})$. At these conditions in accordance with our Hierarchic theory of condensed matter [63, 64] the de Broglie wave length of atoms is tending to infinity and state of macroscopic Bose condensation of ultimate coherence and order, i.e. minimum entropy.

This result of our Unified theory could explain the energy conservation law, independently of the Universe cooling. Decreasing of thermal kinetic energy in the process of cooling is compensated by increasing of potential energy of the matter particle-particle interaction, accompanied Bose condensation. The energy of resonant interaction of matter with Bivacuum virtual pressure waves of basic energy: $\mathbf{E}_{VPW} = n \hbar \omega_0 = n \mathbf{m}_0 \mathbf{c}^2$ at $n = 1$ is also increasing at $\mathbf{v} \rightarrow 0$, keeping the total energy of the Universe permanent.

G.1 The new approach to problem of Time

It follows from the section above, that the Principle of least action is a consequence of basic Tuning energy ($\mathbf{TE}_{q=1}$) of Bivacuum influence on particles, driving the properties of

matter on all hierarchical levels to Golden mean condition. It is shown, using the formula for action (3.130a) that the introduced dimensionless internal *pace of time* for any closed coherent system is determined by the pace of its kinetic energy change (anisotropic in general case), related to changes of Tuning energy (eq.3.131):

$$[\mathbf{dt}/\mathbf{t} = \mathbf{d} \ln \mathbf{t} = -\mathbf{d} \ln \mathbf{T}_k]_{x,y,z} = -\mathbf{d} \ln[(\mathbf{1} + \mathbf{R})\mathbf{TE}]_{x,y,z} \quad (3.140)$$

Using relations (3.140 and 3.131), the pace of the internal time and time itself for closed system of particles can be presented via their acceleration and velocity:

$$\left[\frac{\mathbf{dt}}{\mathbf{t}} = \mathbf{d} \ln \mathbf{t} = -\frac{\mathbf{d}\vec{\mathbf{v}}}{\vec{\mathbf{v}}} \frac{2 - (\mathbf{v}/\mathbf{c})^2}{1 - (\mathbf{v}/\mathbf{c})^2} \right]_{x,y,z} \quad (3.141)$$

$$\left[\mathbf{t} = -\frac{\vec{\mathbf{v}}}{\mathbf{d}\vec{\mathbf{v}}/\mathbf{dt}} \frac{1 - (\mathbf{v}/\mathbf{c})^2}{2 - (\mathbf{v}/\mathbf{c})^2} = -\frac{[(\mathbf{1} + \mathbf{R})\mathbf{TE}]}{\mathbf{d}[(\mathbf{1} + \mathbf{R})\mathbf{TE}]/\mathbf{dt}} \right]_{x,y,z} \quad (3.142)$$

The pace of time and time itself are positive ($\mathbf{t} > \mathbf{0}$), if the particle motion is slowing down ($\mathbf{d}\vec{\mathbf{v}}/\mathbf{dt} < \mathbf{0}$ and $\mathbf{d}\vec{\mathbf{v}} < \mathbf{0}$) and negative, if particles are accelerating. For example, at temperature decreasing the time and its pace are positive. At temperature increasing they are negative. Oscillations of atoms and molecules in condensed matter, like pendulums, are accompanied by alternation the sign of acceleration and, consequently, sign of time. In the absence of acceleration ($\mathbf{d}\vec{\mathbf{v}}/\mathbf{dt} = \mathbf{0}$ and $\mathbf{d}\mathbf{v} = \mathbf{0}$), the time is infinitive and its pace zero:

$$\begin{aligned} \mathbf{t} \rightarrow \infty \quad \text{and} \quad \frac{\mathbf{dt}}{\mathbf{t}} \rightarrow 0 \\ \text{at } \mathbf{d}\vec{\mathbf{v}}/\mathbf{dt} \rightarrow \mathbf{0} \quad \text{and} \quad \mathbf{v} = \text{const} \end{aligned}$$

The postulated principle [I] of conservation of internal kinetic energy of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of symmetric and asymmetric Bivacuum fermions/antifermions: $\left(\mathbf{BVF}_{as}^\uparrow \right)^\phi \equiv \mathbf{F}_\uparrow^\pm$ (3.15) in fact reflects the condition of infinitive life-time of Bivacuum dipoles in symmetric and asymmetric states. The latter means a stability of sub-elementary fermions and elementary particles, formed by them.

In scale of the Universe, the decreasing of all-pervading Tuning energy ($\mathbf{TE}$) in (3.142):

$$\{\mathbf{d}[(\mathbf{1} + \mathbf{R})\mathbf{TE}]/\mathbf{dt}\} < \mathbf{0} \quad (\mathbf{t} > \mathbf{0}) \quad (3.143)$$

means positive direction of "TIME ARROW" $\mathbf{t} > \mathbf{0}$. This corresponds to tending of the actual energy of particles to the energy of rest mass: $\mathbf{m}_v^+ \mathbf{c}^2 \rightarrow \mathbf{q} \mathbf{m}_0 \mathbf{c}^2$ at $\mathbf{v} \rightarrow \mathbf{0}$, mean-

ing cooling of the Universe, if quantization number of virtual pressure waves (VPW[±]) is minimum: $\mathbf{q} = \mathbf{1}$.

The presented approach to the TIME problem differs from the conventional one, following from relativistic theory, however, they do not contradict each other. For example, from (3.142) we can see, that in relativistic conditions, when as a result of acceleration $(d\vec{\mathbf{v}}/dt) > \mathbf{0}$ and $\mathbf{v} \rightarrow \mathbf{c}$ the negative internal time of closed system is tending to zero, like it follows from Lorentz relations for the 'own' time of system.

The internal time (3.142) turns to infinity and its pace (3.141) to zero in the absence of acceleration and deceleration in a closed system $(d\vec{\mathbf{v}}/dt) = \mathbf{0}$. The permanent ($\mathbf{t} \rightarrow \infty$) collective motion of the electrons and atoms of ⁴He in superconductors and superfluid liquids, correspondingly, with constant velocity ($\mathbf{v} = \mathbf{const}$) in the absence of collisions and accelerations are good examples.

In Bivacuum with superfluid properties the existence of stable excitations, like quantized toruses/antitoruses of Bivacuum dipoles and vortical filaments - virtual guides (VirG), described in section 3.9, are also in-line with our time theory.

Each closed real system of elementary particles and macroscopic objects, rotating around common center on stable orbits, like in atoms, planetary systems, galactics, etc. is characterized by different centripetal acceleration $(-d\vec{\mathbf{v}}/dt = \vec{\mathbf{v}}^2/\mathbf{R} = \omega^2\mathbf{R})$ and velocity ($\vec{\mathbf{v}}$). Corresponding characteristic times of such systems is equal to periods of their cycles: $\mathbf{t} = \mathbf{T} = 2\pi/\omega$. The closed systems of virtual particles and virtual waves in Bivacuum do not follow the causality principle and the notion of time for such a systems is uncertain. In conditions of primordial symmetric Bivacuum, when the real and virtual particles are absent, the time is absent also.

At conditions: $[\mathbf{v} = \mathbf{c} = \mathbf{const}]$, valid for the case of photons, we get from (3.142) the uncertainty for time, like: $\mathbf{t} = \mathbf{0}/\mathbf{0}$. The similar result we get for state of virtual Bose condensate (**VirBC**) in Bivacuum, when the *external translational* velocity of Bivacuum fermions (**BVF**[†]) and Bivacuum bosons (**BVB**[±]) is equal to zero ($\mathbf{v} = \mathbf{0} = \mathbf{const}$). The latter condition corresponds to totally symmetric **BVF**[†] and **BVB**[±], of primordial Bivacuum, when their torus (V⁺) and antitorus (V⁻) mass, charge and magnetic moments are equal to each other (see eqs.3.16 and 3.29). It follows from (3.142) that the internal time for each selected closed system of particles is a parameter, characterizing the average velocity and acceleration of these particles, i.e. this system internal dynamics.

H. Virtual Replicas (VR) of material objects in Bivacuum

The basically new concept of Virtual replica (**VR**) or virtual hologram of any material object in Bivacuum, is introduced in our Unified theory [21-23]. The **VR** is result of interference of *primary* all-pervading quantized Virtual Pressure Waves ($\mathbf{VPW}_q^+$ and $\mathbf{VPW}_q^-$) and Virtual Spin waves ($\mathbf{VirSW}_q^{S=\pm 1/2}$) of Bivacuum, working as the "reference waves" in hologram formation and the same waves, modulated by de Broglie waves of atoms and molecules, representing the "object waves" $\mathbf{VPW}_m^\pm$ and $\mathbf{VirSW}_m^{\pm 1/2}$. The frequencies of *basic reference* virtual pressure waves ($\mathbf{VPW}_{q=1}^\pm \equiv \mathbf{VPW}_0^\pm$) and virtual spin waves ($\mathbf{VirSW}_{q=1}^{\pm 1/2} \equiv \mathbf{VirSW}_0^{\pm 1/2}$) of Bivacuum are equal to Compton frequencies, like the carrying frequencies of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of sub-elementary fermions of triplets at zero external translational velocity of particles ($\mathbf{v} = \mathbf{0}$) (Fig.3.1):

$$[\omega_{VPW_0} = \omega_{VirSW_0} = \omega_0 = \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}}^{\mathbf{v}=\mathbf{0}} = \mathbf{q}\mathbf{m}_0\mathbf{c}^2/\hbar]^i$$

Two kinds of primary wave modulation ($\mathbf{VPW}_m^\pm$ and $\mathbf{VirSW}_m^{\pm 1/2}$) are realized by cumulative virtual clouds ($\mathbf{CVC}^\pm$), emitted/absorbed in the process of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of pairs: $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_C \rightleftharpoons [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]_W$ of elementary triplets (electrons, protons, neutrons) $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\downarrow^\pm >^i$, and the *recoil angular momentum*, generated by $\mathbf{CVC}^{\pm 1/2}$ of unpaired sub-elementary fermion $\mathbf{F}_\downarrow^\pm >^i$, correspondingly:

$$[(\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-)_C + (\mathbf{F}_\downarrow^\pm)_W] \overset{+\mathbf{CVC}^\pm - \text{Recoil}}{\underset{-\mathbf{CVC}^\pm + \text{Antirecoil}}{<====>}} [(\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-)_W + (\mathbf{F}_\downarrow^\pm)_C] \quad (3.144)$$

The in-phase $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of a sub-elementary fermion $\mathbf{F}_\downarrow^+$ and antifermion $\mathbf{F}_\uparrow^-$ of pair $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$, accompanied by reversible [emission $\rightleftharpoons$ absorption] of cumulative virtual clouds $\mathbf{CVC}^+$ and $\mathbf{CVC}^-$ and the 'object' virtual pressure wave ($\mathbf{VPW}_m^+$ and $\mathbf{VPW}_m^-$) excitation. However, the recoil energy and the angular momenta of $\mathbf{CVC}^+$ and $\mathbf{CVC}^-$ of $\mathbf{F}_\uparrow^-$ and $\mathbf{F}_\downarrow^+$ of pairs compensate each other and the resulting recoil energy of $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$ is zero.

Superposition of $\mathbf{VPW}_m^+$ and $\mathbf{VPW}_m^-$, excited by cumulative virtual clouds: $\mathbf{CVC}^+$ and $\mathbf{CVC}^-$, emitted and absorbed in the process of the in-phase $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of pairs $[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]$ of rotating triplets (Fig.3.1), activate *quantized whirls* in Bivacuum. The stability of VR of object, as a *hierarchical system of quantized metastable torus-like and vortex filaments structures* formed by $\mathbf{VPW}_m^\pm$ and by $\mathbf{VirSW}_m^{\pm 1/2}$ excited by paired and unpaired

sub-elementary fermions, correspondingly, in superfluid Bivacuum, could be responsible for so-called ”**phantom effect**” of object after its removing to another distant place.

H.1 Bivacuum perturbations, induced by the oscillation of the total energy of de Broglie waves, accompanied by their thermal vibrations and recoil $\rightleftharpoons$ antirecoil effects

In contrast to the situation with unpaired sub-elementary fermion $(\mathbf{F}_{\uparrow}^{\pm})$ in triplets, the recoil/antirecoil momenta and energy, accompanying the in-phase emission/absorption of $\mathbf{CVC}_{\mathbf{S}=+1/2}^+$ and $\mathbf{CVC}_{\mathbf{S}=-1/2}^-$ by $\mathbf{F}_{\uparrow}^+$ and $\mathbf{F}_{\downarrow}^-$ of pair $[\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-]$, totally compensate each other in the process of their $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation. Such pairs display the properties of neutral particles with zero spin and zero rest mass:

$$[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_C < \frac{[\mathbf{E}_{CVC^+} + \mathbf{E}_{CVC^-}] + \Delta \mathbf{v} \mathbf{P}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-}}{[\mathbf{E}_{CVC^+} + \mathbf{E}_{CVC^-}] - \Delta \mathbf{v} \mathbf{P}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-}} > [\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]_W \quad (3.145)$$

The total energy increment of elementary particle, equal to that of each of sub-elementary fermions of triplet, generated in nonequilibrium processes, accompanied by entropy change, like melting, boiling, etc., can be presented in a few manners:

$$\Delta \mathbf{E}_{tot} = \Delta(\mathbf{m}_V^+ \mathbf{c}^2) = \Delta \left(\frac{\mathbf{m}_0 \mathbf{c}^2}{[1 - (\mathbf{v}/\mathbf{c})^2]^{1/2}} \right) = \frac{\mathbf{m}_0 \mathbf{v}}{\mathbf{R}^3} \Delta \mathbf{v} = \frac{\mathbf{p}}{\mathbf{R}^2} \Delta \mathbf{v} = \frac{\mathbf{h}}{\lambda_B \mathbf{R}^2} \Delta \mathbf{v} \quad (3.146)$$

$$or : \Delta \mathbf{E}_{tot} = \Delta[(\mathbf{m}_V^+ - \mathbf{m}_V^-) \mathbf{c}^2 (\mathbf{c}/\mathbf{v})^2] = \frac{2\mathbf{T}_k}{\mathbf{R}^2} \frac{\Delta \mathbf{v}}{\mathbf{v}} \quad (3.147)$$

$$or : \Delta \mathbf{E}_{tot} = \frac{2\mathbf{T}_k}{\mathbf{R}^2} \frac{\Delta \mathbf{v}}{\mathbf{v}} = \Delta[\mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in}] + \Delta(\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} \quad (3.147a)$$

where: $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ is the relativistic factor; $\Delta \mathbf{v}$ is the increment of the external translational velocity of particle; the actual inertial mass of sub-elementary particle is: $\mathbf{m}_V^+ = \mathbf{m}_0/\mathbf{R}$; $\mathbf{p} = \mathbf{m}_V^+ \mathbf{v} = \mathbf{h}/\lambda_B$ is the external translational momentum of unpaired sub-elementary particle $\mathbf{F}_{\uparrow}^{\pm} >^i$, equal to that of whole triplet $< [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-] + \mathbf{F}_{\uparrow}^{\pm} >^i$; $\lambda_B = \mathbf{h}/\mathbf{p}$ is the de Broglie wave of particle; $2\mathbf{T}_k = \mathbf{m}_V^+ \mathbf{v}^2$ is a doubled kinetic energy; $\Delta \ln \mathbf{v} = \Delta \mathbf{v}/\mathbf{v}$.

The increments of *internal* rotational and *external* translational contributions to total energy of the de Broglie wave (see eq. 3.147a) are, correspondingly:

$$\Delta[\mathbf{R} (\mathbf{m}_0 \mathbf{c}^2)_{rot}^{in}] = -2\mathbf{T}_k (\Delta \mathbf{v}/\mathbf{v}) \quad (3.148)$$

$$\Delta(\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext} = \Delta(2\mathbf{T}_k)_{tr}^{ext} = 2\mathbf{T}_k \frac{1 + \mathbf{R}^2}{\mathbf{R}^2} \frac{\Delta \mathbf{v}}{\mathbf{v}} \quad (3.148a)$$

The time derivative of total energy of elementary de Broglie wave, following from (3.60 - 3.60b) is:

$$\frac{d\mathbf{E}_{tot}}{dt} = \frac{2\mathbf{T}_k}{\mathbf{R}^2 \mathbf{v}} \frac{d\mathbf{v}}{dt} = \frac{2\mathbf{T}_k}{\mathbf{R}^2} \frac{d \ln \mathbf{v}}{dt} \quad (3.149)$$

Between the increments of energy of triplets, equal to that of unpaired $\Delta\mathbf{E}_{tot} = \Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^{\pm}}$ and increments of modulated $\mathbf{CVC}_{\mathbf{m}}^+$ and $\mathbf{CVC}_{\mathbf{m}}^-$, emitted by pair $[\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+]$ in the process of $[\mathbf{C} \rightarrow \mathbf{W}]$ transition, the direct correlation is existing.

These cumulative virtual clouds modulated by particle's de Broglie wave ($\lambda_B = \mathbf{h}/\mathbf{m}_{\mathbf{V}}^+ \mathbf{v}$): $\mathbf{CVC}_{\mathbf{m}}^+$ and $\mathbf{CVC}_{\mathbf{m}}^-$ of paired sub-elementary fermions, superimposed with basic virtual pressure waves ($\mathbf{VPW}_0^{\pm}$) of Bivacuum, turn them to the *object waves* ($\mathbf{VPW}_{\mathbf{m}}^{\pm}$), necessary for virtual hologram of the object formation:

$$\Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+}^{\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+} = \frac{\mathbf{h}}{\lambda_B \mathbf{R}^2} \Delta\mathbf{v} = \frac{2\mathbf{T}_k}{\mathbf{R}^2} \Delta \ln \mathbf{v} \xrightarrow{\mathbf{CVC}_{\mathbf{m}}^+} \Delta(\mathbf{VPW}_{\mathbf{m}}^+) \quad (3.150)$$

$$- \Delta\mathbf{E}_{\mathbf{F}_{\downarrow}^-}^{\mathbf{F}_{\uparrow}^- \bowtie \mathbf{F}_{\downarrow}^+} \xrightarrow{\mathbf{CVC}_{\mathbf{m}}^-} \Delta(\mathbf{VPW}_{\mathbf{m}}^-) \quad (3.150a)$$

The virtual pressure waves represent oscillations of corresponding virtual pressure ($\mathbf{VirP}_m^{\pm}$).

The increment of total energy of fermion or antifermion, equal to increment of its unpaired sub-elementary fermion can be presented via increments of paired sub-elementary fermions (3.148 and 3.148a), like:

$$\begin{aligned} \Delta\mathbf{E}_{tot} = \Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+} &= \frac{1}{2} \left(\Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} - \Delta\mathbf{E}_{\mathbf{F}_{\downarrow}^-}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} \right) + \frac{1}{2} \left(\Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} + \Delta\mathbf{E}_{\mathbf{F}_{\downarrow}^-}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} \right) = \\ &= \Delta\mathbf{T}_k^+ + \Delta\mathbf{V}^+ \quad (3.151) \end{aligned}$$

where, the contributions of the kinetic and potential energy increments to the total energy increment, interrelated with increments of positive and negative virtual pressures ($\Delta\mathbf{VirP}^{\pm}$), are, correspondingly:

$$\Delta\mathbf{T}_k = \frac{1}{2} \left(\Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} - \Delta\mathbf{E}_{\mathbf{F}_{\downarrow}^-}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} \right) \sim \frac{1}{2} (\Delta\mathbf{VirP}^+ - \Delta\mathbf{VirP}^-) \sim \alpha \Delta (\mathbf{m}_V^+ \mathbf{v}^2)_{\mathbf{F}_{\uparrow}^{\pm}} \quad (3.152)$$

$$\Delta\mathbf{V} = \frac{1}{2} \left(\Delta\mathbf{E}_{\mathbf{F}_{\uparrow}^+}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} + \Delta\mathbf{E}_{\mathbf{F}_{\downarrow}^-}^{\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-} \right) \sim \frac{1}{2} (\Delta\mathbf{VirP}^+ + \Delta\mathbf{VirP}^-) \sim \beta \Delta (\mathbf{m}_V^+ + \mathbf{m}_V^-) \mathbf{c}_{\mathbf{F}_{\uparrow}^{\pm}}^2 \quad (3.152a)$$

The specific information of any object is imprinted in its Virtual Replica (VR), because cumulative virtual clouds ($\mathbf{CVC}_m^\pm$) of the object's elementary particles and their superposition with Bivacuum pressure waves and Virtual spin waves: $\mathbf{VPW}_m^\pm$ and $\mathbf{VirSW}_m^{\pm 1/2}$ are modulated by frequency, phase and amplitude of the thermal de Broglie waves of molecules, composing this object. Comparing eqs. 3.152a and 3.88a we may see, that the modulated gravitational virtual pressure waves form a part of VR.

H.2 Modulation of the basic Virtual Pressure Waves ($\mathbf{VPW}_q^\pm$) and Virtual Spin Waves ($\mathbf{VirSW}_q^{\pm 1/2}$) of Bivacuum by molecular translations and librations

The external translational/librational kinetic energy of particle $(\mathbf{T}_k)_{tr,lb}$ is directly related to its de Broglie wave length (λ_B), the group ($\mathbf{v}$), phase velocity ($\mathbf{v}_{ph}$) and frequency ($\nu_B = \omega_B/2\pi$):

$$\left(\lambda_B = \frac{\mathbf{h}}{\mathbf{m}_V^+ \mathbf{v}} = \frac{\mathbf{h}}{2\mathbf{m}_V^+ \mathbf{T}_k} = \frac{\mathbf{v}_{ph}}{\nu_B} = 2\pi \frac{\mathbf{v}_{ph}}{\omega_B} \right)_{tr,lb} \quad (3.153)$$

where the de Broglie wave frequency is related to its length and kinetic energy of particle as:

$$\left[\nu_B = \frac{\omega_B}{2\pi} = \frac{h}{2\mathbf{m}_V^+ \lambda_B^2} = \frac{\mathbf{m}_V^+ \mathbf{v}^2}{2h} \right]_{tr,lb} \quad (3.154)$$

It follows from our model, that zero-point frequency of $[C \rightleftharpoons W]$ pulsation $(\omega_0)^i$ of sub-elementary fermions and antifermions, forming triplets of elementary particles $< [\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-] + \mathbf{F}_\uparrow^\pm >^i$, accompanied by $[emission \rightleftharpoons absorbtion]$ of cumulative virtual clouds $\mathbf{CVC}^\pm$ has the same value, as a basic (reference) frequency $(\omega_{q=1} = \omega_0 = \mathbf{m}_0 \mathbf{c}^2 / \hbar)^i$ of Bivacuum.

The total energy of de Broglie wave and resulting frequency of pulsation $(\omega_{\mathbf{C} \rightleftharpoons \mathbf{W}})$ (eq. 3.78) is a result of modulation of the internal frequency, related to the rest mass of particle, by the most probable frequency of de Broglie wave of the whole particle (ω_B), determined by its most probable external momentum: $\mathbf{p} = \mathbf{m}_V^+ \mathbf{v}$.

In a composition of condensed matter this value is different for thermal librations and translation of molecules. The corresponding most probable modulation frequencies of translational and librational de Broglie waves are possible to calculate, using our Hierarchic theory of condensed matter and based on this theory computer program [57, 63, 64].

The interference between the basic $(\mathbf{q} = \mathbf{j} - \mathbf{k} = \mathbf{1})$ reference virtual spin waves $(\mathbf{VirSW}_{q=1}^{\pm 1/2})$ and virtual pressure waves $\mathbf{VPW}_{q=1}^\pm$ of Bivacuum with virtual 'object

waves', modulated by the librational and translational de Broglie waves ($\lambda_{lb,tr} = \mathbf{h}/\mathbf{m}_V^+ \mathbf{v}$) of molecules, produce the holographic-like image or Virtual Replicas (VR) of the object.

The *frequencies* of virtual pressure waves ($\mathbf{VPW}_m^\pm$) and virtual spin waves ($\mathbf{VirSW}_m^{\pm 1/2}$) are modulated by de Broglie waves of the object particles. Corresponding modulation frequencies are related to frequencies of librational (ω_{lb}) and translational (ω_{tr}) de Broglie waves of molecules of matter (3.154) in accordance to rules of combinational resonance:

$$\omega_{\mathbf{VPW}_m^\pm}^i = \mathbf{R} \omega_0^i + \mathbf{g} \omega_{tr} + \mathbf{r} \omega_{lb} \cong [\mathbf{R} \omega_0^i + \mathbf{g} \omega_{tr}] \quad (3.155)$$

$$\omega_{\mathbf{VirSW}_m^{\pm 1/2}}^i = \mathbf{R} \omega_0^i + \mathbf{r} \omega_{lb} + \mathbf{g} \omega_{tr} \cong \mathbf{R} \omega_0^i + \mathbf{r} \omega_{lb} \quad (3.155a)$$

$$\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}; \quad \mathbf{g}, \mathbf{r} = 1, 2, 3 \dots (\text{integer numbers})$$

Each of 24 collective excitations of condensed matter, introduced in our Hierarchic theory [22, 63], has his own characteristic frequency and can be imprinted in Virtual Replica of the object, as a corresponding pattern.

Three kinds of modulations: *frequency, amplitude and phase* of the object ($\mathbf{VPW}_m^\pm$) and ($\mathbf{VirSW}_m^{\pm 1/2}$) by de Broglie waves of the object's molecules may be described by known relations [68]:

1. *The frequencies* of virtual pressure waves ($\omega_{VPW_m^\pm}^M$) and spin waves ($\omega_{VirSW_m^\pm}^M$), *modulated* by translational and librational de Broglie waves of the object's molecules, can be presented as:

$$\omega_{VPW_m^\pm}^M = \mathbf{R} \omega_0^i + \Delta \omega_B^{tr} \cos \omega_B^{tr} t \quad (3.156)$$

$$\omega_{VirSW_m^{\pm 1/2}}^M = \mathbf{R} \omega_0^i + \Delta \omega_B^{lb} \cos \omega_B^{lb} t \quad (3.156a)$$

The Compton pulsation frequency of elementary particles is equal to basic frequency of Bivacuum virtual waves at $\mathbf{q} = \mathbf{j} - \mathbf{k} = \mathbf{1}$:

$$\omega_0^i = \mathbf{m}_0^i \mathbf{c}^2 / \hbar = \omega_{VPW_{q=1}^\pm, VirSW_{q=1}}^i \quad (3.157)$$

Such kind of modulation is accompanied by two satellites with frequencies: $(\omega_0^i + \omega_B^{tr,lb})$ and $(\omega_0^i - \omega_B^{tr,lb}) = \Delta \omega_{tr,lb}^i$. The latter is named frequency deviation. In our case: $\omega_0^e (\sim 10^{21} s^{-1}) \gg \omega_B^{tr,lb} (\sim 10^{12} s^{-1})$ and $\Delta \omega_{tr,lb} \gg \omega_B^{tr,lb}$.

The temperature of condensed matter and phase transitions may influence the modulation frequencies of de Broglie waves of its molecules.

2. The amplitudes of virtual pressure waves ($VPW_m^\pm$) and virtual spin waves $VirSW_m^{\pm 1/2}$ (informational waves) modulated by the object are dependent on translational and librational de Broglie waves frequencies as:

$$\mathbf{A}_{VPW_m^\pm} \approx \mathbf{A}_0 (\sin \mathbf{R}\omega_0^i \mathbf{t} + \gamma \omega_B^{tr} \sin \mathbf{t} \cdot \cos \omega_B^{tr} t) \quad (3.158)$$

$$\mathbf{I}_{VirSW_m^{\pm 1/2}} \approx \mathbf{I}_0 (\sin \mathbf{R}\omega_0^i \mathbf{t} + \gamma \omega_B^{lb} \sin \mathbf{t} \cdot \cos \omega_B^{lb} t) \quad (3.158a)$$

where: the informational/spin field amplitude is determined by the amplitude of Bivacuum fermions $[BVF^\uparrow \rightleftharpoons BVF^\downarrow]$ equilibrium constant oscillation: $\mathbf{I}_S \equiv \mathbf{I}_{VirSW}^{\pm 1/2} \sim \mathbf{K}_{BVF^\uparrow \rightleftharpoons BVF^\downarrow}(\mathbf{t})$

The index of frequency modulation is defined as: $\gamma = (\Delta\omega_{tr,lb}/\omega_B^{tr,lb})$. The carrying zero-point pulsation frequency of particles is equal to the basic frequency of Bivacuum virtual waves: $\omega_{VPW_0^\pm, VirSW_0}^i = \omega_0^i$. Such kind of modulation is accompanied by two satellites with frequencies: $(\omega_0^i + \omega_B^{tr,lb})$ and $(\omega_0^i - \omega_B^{tr,lb}) = \Delta\omega_{tr,lb}$. In our case: $\omega_0^e (\sim 10^{21} s^{-1}) \gg \omega_B^{tr,lb} (\sim 10^{12} s^{-1})$ and $\gamma \gg 1$.

The fraction of molecules in state of mesoscopic molecular Bose condensation (mBC), representing, coherent clusters [22,63] is a factor, influencing the amplitude (A_0) and such kind of modulation of Virtual replica of the object.

3. The phase modulated $VPW_m^\pm$ and $VirSW_m^{\pm 1/2}$ by de Broglie waves of molecules, related to their translations and librations, can be described like:

$$\mathbf{A}_{VPW_m^\pm}^M = \mathbf{A}_0 \sin (\mathbf{R}\omega_0 \mathbf{t} + \Delta\varphi_{tr} \sin \omega_B^{tr} \mathbf{t}) \quad (3.159)$$

$$\mathbf{I}_{VirSW_m^{\pm 1/2}}^M = \mathbf{I}_0 \sin (\mathbf{R}\omega_0 \mathbf{t} + \Delta\varphi_{lb} \sin \omega_B^{lb} \mathbf{t}) \quad (3.159a)$$

The value of phase increment $\Delta\varphi_{tr,lb}$ of modulated virtual waves of Bivacuum ($VPW_m^\pm$ and $VirSW_m^{\pm 1/2}$), contains the information about geometrical properties of the object.

The phase modulation takes place, if the phase increment $\Delta\varphi_{tr,lb}$ is independent on the modulation frequency $\omega_B^{tr,lb}$.

The virtual holographic image, resulting from interference pattern of the virtual object waves: $\mathbf{VPW}_m^\pm$ and $\mathbf{VirSW}_m^{\pm 1/2}$, modulated (scattered) by translations and librations of molecules, with similar reference waves of Bivacuum ($\mathbf{VPW}_q^\pm$ and $\mathbf{VirSW}_q^{\pm 1/2}$) contains full information about the object's internal dynamic and spatial properties and may be named "Virtual Replica (VR)" of the object.

I. Possible mechanism of entanglement between remote elementary particles via Virtual Guides of spin, momentum and energy ($\mathbf{VirG}_{S,M,E}$)

In accordance to our theory, the instant nonlocal quantum entanglement between two or more distant *similar* elementary particles (electrons, protons, neutrons, photons), named [Sender (S)] and [Receiver (R)], revealed in a lot of experiments, started by Aspect and Grangier [69], involves a few stages:

1. Superposition of their nonlocal and distant components of Virtual replicas ($\mathbf{VR}$) or Virtual hologram, formed by interference of modulated by de Broglie waves of object Virtual spin waves: $[\mathbf{VirSW}_{\mathbf{m}}^{\pm 1/2}(S) \leq \Rightarrow \mathbf{VirSW}_{\mathbf{m}}^{\pm 1/2}(R)]$ and Virtual pressure waves: $[\mathbf{VPW}_{\mathbf{m}}^{\pm}(S) \leq \Rightarrow \mathbf{VPW}_{\mathbf{m}}^{\pm}(R)]$ with corresponding *reference waves* of Bivacuum $\mathbf{VirSW}_{\mathbf{q}}^{\pm 1/2}$ and $\mathbf{VPW}_{\mathbf{q}}^{\pm}$, described in Section 3.8.2;

2. Tuning (frequency and phase synchronization) of de Broglie waves of remote interacting identical particles (like electrons, protons) and their complexes in form of atoms and molecules, as a condition of Virtual Guides of spin-momentum-energy ($\mathbf{VirG}_{SME}$) between [S] and [R] formation (see Fig. 3.4).

Superpositions of counterphase ($\mathbf{VirSW}_{\mathbf{m}}^{\pm 1/2}$) of [S] and [R], excited by their unpaired sub-elementary fermions $F_{\uparrow}^{\pm}$ of triplets $< [\mathbf{F}_{\uparrow}^{+} \bowtie \mathbf{F}_{\downarrow}^{-}] + \mathbf{F}_{\uparrow}^{\pm} >^i$ of opposite spins in form of virtual standing waves, may stimulate formation of two kinds of Virtual Guides:

a) single *nonlocal virtual guides* $\mathbf{VirG}_{SME}^{(\mathbf{BVB}^{\pm})^i}$ - virtual microtubules from Bivacuum bosons $(\mathbf{BVB}^{\pm})^i$. In this case the $\mathbf{VirG}_{SME}^{(\mathbf{BVB}^{\pm})^i}$ is not rotating as a whole around its axis and the resulting spin is zero.

b) twin *nonlocal virtual guides* $\mathbf{VirG}_{SME}^{[\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}]^i}$ from Cooper pairs of Bivacuum fermions $[\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}]^i$. In this case each of two adjacent microtubules rotate around their own axes in opposite directions. The resulting angular moment (spin) of such pair is also zero.

Two remote coherent triplets - elementary particles, like (electron - electron) or (proton - proton) with similar frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]_{e,p}$ pulsation and opposite spins (phase) can be connected by Virtual guides ($\mathbf{VirG}_{SME}^i$) of spin (S), momentum (M) and energy (E) from Sender to Receiver of both kinds. The spin - information (qubits), momentum and kinetic energy instant transmission via such $\mathbf{VirG}_{SME}^i$ from [S] and [R] is possible. The same is

true for two synchronized photons (bosons) of opposite spins.

The double $\mathbf{VirG}_{SME}^{[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow]^i}$ can be transformed to single $\mathbf{VirG}_{SME}^{(\mathbf{BVB}^\pm)^i}$ by conversion of opposite Bivacuum fermions: $\mathbf{BVF}^\uparrow = [\mathbf{V}^+ \uparrow\uparrow \mathbf{V}^-]$ and $\mathbf{BVF}^\downarrow = [\mathbf{V}^+ \downarrow\downarrow \mathbf{V}^-]$ to the pair of Bivacuum bosons of two possible alternatives of polarization:

$$\mathbf{BVB}^+ = [\mathbf{V}^+ \uparrow\downarrow \mathbf{V}^-] \quad or \quad \mathbf{BVB}^- = [\mathbf{V}^+ \downarrow\uparrow \mathbf{V}^-]$$

Superposition of two nonlocal virtual spin waves excited by similar elementary particles (electrons or protons) of Sender $(\mathbf{VirSW}_m^{S=+1/2})_S$ and Receiver $(\mathbf{VirSW}_m^{S=-1/2})_R$ of the same pulsation frequency and opposite spins, i.e. opposite phase of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation, forms *Virtual Guide* of spin, momentum and energy $(\mathbf{VirG}_{SME})^i$ (Fig.3.4):

$$\left[< [\mathbf{F}_\downarrow^+ \bowtie \mathbf{F}_\uparrow^-]_C + (\mathbf{F}_\downarrow^-)_W >_S \xrightarrow{\mathbf{VirSW}_S} \underset{\mathbf{BVB}^-}{\overset{\mathbf{BVB}^+}{\rightleftharpoons}} \xleftarrow{\mathbf{VirSW}_R} < (\mathbf{F}_\uparrow^-)_C + [\mathbf{F}_\downarrow^- \bowtie \mathbf{F}_\uparrow^+]_W >_R \right]^i$$

$$or : [\mathbf{n}_+ \mathbf{BVB}^+ (\mathbf{V}^+ \uparrow\downarrow \mathbf{V}^-) + \mathbf{n}_- \mathbf{BVB}^- (\mathbf{V}^+ \downarrow\uparrow \mathbf{V}^-)]^i = (\mathbf{VirG}_{SME}^{ext})^i \quad (3.160)$$

The spin exchange via $\mathbf{VirG}_{SME}^i$ is accompanied either by the instant change of Bivacuum boson polarization: $[\mathbf{BVB}^+ \rightleftharpoons \mathbf{BVB}^-]^i$ or by instant spin state change of both Bivacuum fermions, forming virtual Cooper pairs in the double virtual microtubule:

$$[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow]^i \xrightleftharpoons{(S=+1/2) \rightarrow (S=-1/2)} [\mathbf{BVF}^\downarrow \bowtie \mathbf{BVF}^\uparrow]^i \quad (3.161)$$

The radius of virtual microtubules of $\mathbf{VirG}_{SME}^i$ is dependent on generation of torus and antitorus ($i = e, \mu, \tau$), forming them:

$$\mathbf{L}_V^e = \hbar / \mathbf{m}_0^e \mathbf{c} \gg \mathbf{L}_V^\mu = \hbar / \mathbf{m}_0^\mu \mathbf{c} > \mathbf{L}_V^\tau = \hbar / \mathbf{m}_0^\tau \mathbf{c}$$

The radius of $\mathbf{VirG}_{SME}^e$, connecting two remote electrons, is the biggest one ($\mathbf{L}^e$). The radius of $\mathbf{VirG}_{SME}^\tau$, connecting two protons or neutrons ($\mathbf{L}^\tau$) is about 3.5×10^3 times smaller. The entanglement between similar atoms in pairs, like hydrogen, oxygen, carbon or nitrogen can be realized via complex virtual guides of atoms ($\mathbf{VirG}_{SME}^{at}$), representing *multishell constructions*.

The increments of momentum $\Delta \mathbf{p} = \Delta(\mathbf{m}_V^+ \mathbf{v})_{tr,lb}$ and kinetic $(\Delta \mathbf{T}_k)_{tr,lb}$ energy transmission from [S] to [R] of *selected generation of elementary particles* is determined by the translational and librational velocity variation ($\Delta \mathbf{v}$) of nuclei of (S). This means, that energy/momentum transition from [S] to [R] is possible, if they are in nonequilibrium state.

The variation of kinetic energy of atomic nuclei under external force application, induces nonequilibrium in a system ($\mathbf{S} + \mathbf{R}$) and decoherence of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of protons and neutrons of [S] and [R]. The nonlocal energy transmission from [S] to [R] is possible, if the decoherence is not big enough for disassembly of the virtual microtubules and their systems in the case of atoms. The electronic $\mathbf{VirG}_{SME}^e$, as more coherent (not so dependent on thermal vibrations), can be responsible for stabilization of the complex atomic Virtual Guides $\sum \mathbf{VirG}_{SME}^{e,p,n}$.

The values of the energy and velocity increments of free elementary particles are interrelated by (3.147).

The instantaneous energy flux via $(\mathbf{VirG}_{SME})^i$, is mediated by pulsation of energy and radii of torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$) of Bivacuum bosons: $\mathbf{BVB}^+ = [\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-]$. Corresponding energy increments of the actual torus and complementary antitorus of $\mathbf{BVB}^\pm$, forming $(\mathbf{VirG}_{SME})^i$, are directly related to increments of Sender particle external velocity ($\Delta \mathbf{v}$):

$$\begin{aligned} \Delta E_{V^+} &= +\Delta m_V^+ c^2 = \left(+ \frac{\mathbf{p}^+}{\mathbf{R}^2} (\Delta \mathbf{v})_{\mathbf{F}_\uparrow^+}^{[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]} = m_V^+ c^2 \frac{\Delta \mathbf{L}_{V^+}}{\mathbf{L}_{V^+}} \right)_{N,S} \quad \text{actual} \\ \Delta E_{V^-} &= -\Delta m_V^- c^2 = \left(- \frac{\mathbf{p}^-}{\mathbf{R}^2} (\Delta \mathbf{v})_{\mathbf{F}_\uparrow^-}^{[\mathbf{F}_\uparrow^+ \bowtie \mathbf{F}_\downarrow^-]} = -m_V^- c^2 \frac{\Delta \mathbf{L}_{V^-}}{\mathbf{L}_{V^-}} \right)_{N,S} \quad \text{complementary} \end{aligned} \quad (3.162)$$

$$(3.162a)$$

where: $\mathbf{p}^+ = m_V^+ \mathbf{v}$; $\mathbf{p}^- = m_V^- \mathbf{v}$ are the actual and complementary momenta; $\mathbf{L}_{V^+} = \hbar / m_V^+ c$ and $\mathbf{L}_{V^-} = \hbar / m_V^- c$ are the radii of torus and antitorus of $\mathbf{BVB}^\pm = [\mathbf{V}^+ \uparrow \downarrow \mathbf{V}^-]$, forming $\mathbf{VirG}_{S,M,E}^{in,ext}$.

The nonlocal energy exchange between [S] and [R] is accompanied by the instant pulsation of radii of tori ($\mathbf{V}^+$) and antitori ($\mathbf{V}^-$) of $\mathbf{BVF}^\uparrow$ and $\mathbf{BVB}^\pm$, accompanied by corresponding pulsation $|\Delta \mathbf{L}_{V^\pm} / \mathbf{L}_{V^\pm}|$ of the whole virtual microtubule $\mathbf{VirG}_{SME}$ (Fig.3.4).

Most effectively the proposed mechanism of spin (information), momentum and energy exchange can work between Sender and Receiver, containing coherent molecular clusters - mesoscopic Bose condensate (mBC) [21, 22, 63].

The instantaneous angular momentum (spin) exchange between [S] and [R] does not need the radius pulsation, but only the instantaneous polarization change of Bivacuum dipoles $(\mathbf{BVB}^+ \rightleftharpoons \mathbf{BVB}^-)^i$ or *Cooper pairs* $[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow] \rightleftharpoons [\mathbf{BVF}^\downarrow \bowtie \mathbf{BVF}^\uparrow]$, forming $\mathbf{VirG}_{SME}$.

The Resonance Nonlocal Interaction Between
Distant Electrons of Opposite Spins and Phase of [C $\rightleftharpoons$ W] Pulsation
via Virtual Spin - Momentum - Energy Guide [VirG_{SME}]




Fig.3.4. The mechanism of nonlocal Bivacuum mediated interaction (entanglement) between two distant unpaired sub-elementary fermions of 'tuned' elementary triplets (particles) of the opposite spins $\langle [F_+^+ \otimes F_+^-] + F_+^- \rangle_{\text{Sender}}^i$ and $\langle [F_+^+ \otimes F_+^-] + F_+^- \rangle_{\text{Receiver}}^i$, with close frequency of [C $\rightleftharpoons$ W] pulsation and close de Broglie wave length ($\lambda_B = \hbar/m_v^+ v$) of particles. The tunnelling of momentum and energy increments: $\Delta |m_v^{\pm} c^2| \sim \Delta |\text{VirP}^+| \pm \Delta |\text{VirP}^-|$ from Sender to Receiver and vice-verse via Virtual spin-momentum-energy Guide [VirG_{SME}ⁱ] is accompanied by instantaneous pulsation of diameter ($2\Delta L_{v^{\pm}}$) of this virtual microtubule, formed by Bivacuum bosons **BVB**[±] or double microtubule, formed by Cooper pairs of Bivacuum fermions: $[BVF^{\uparrow} \otimes BVF^{\downarrow}]$. The spin state exchange between [S] and [R] can be realized by the instantaneous change polarization of Cooper pairs: $[BVF^{\uparrow} \otimes BVF^{\downarrow}] \rightleftharpoons [BVF^{\downarrow} \otimes BVF^{\uparrow}]$ and Bivacuum bosons: **BVB**⁺ $\rightleftharpoons$ **BVB**⁻.

In virtual microtubules (**VirG**_{SME}ⁱ)ⁱ the time and its 'pace' are uncertain: **t** = **0/0**, as far the external velocities (**v**) and accelerations (**dv/dt**) of Bivacuum dipoles, composing such virtual Bose condensate, are zero (see eqs. 3.141 and 3.142).

*I.1 The role of tuning force (**F**_{VPW[±]}) of virtual
pressure waves **VPW**_q[±] of Bivacuum in entanglement*

The tuning between **two similar elementary** particles: 'sender (S)' and 'receiver (R)' via $\mathbf{VirG}_{SME}^i$ may be qualitatively described, using well known model of *damped harmonic oscillators*, interacting with all-pervading virtual pressure waves ($\mathbf{VPW}^\pm$) of Bivacuum with fundamental frequency ($\omega_0 = \mathbf{m}_0 \mathbf{c}^2 / \hbar$). The criteria of tuning - synchronization of [S] and [R] is the equality of the amplitude probability of resonant energy exchange of Sender and Receiver with virtual pressure waves ($\mathbf{VPW}_0^\pm$): $\mathbf{A}_{C \rightleftharpoons W}^S = \mathbf{A}_{C \rightleftharpoons W}^R$, resulting from minimization of frequency difference ($\omega_S - \omega_0 \rightarrow 0$ and $(\omega_R - \omega_0) \rightarrow 0$):

$$\mathbf{A}_{C \rightleftharpoons W}^S \sim \left[\frac{1}{(\mathbf{m}_V^+)_S} \frac{\mathbf{F}_{\mathbf{VPW}^\pm}}{(\omega_S^2 - \omega_0^2) + \text{Im } \gamma \omega_S} \right] \quad (3.163)$$

$$[\mathbf{A}_{C \rightleftharpoons W}^R]_{x,y,z} \sim \left[\frac{1}{(\mathbf{m}_V^+)_R} \frac{\mathbf{F}_{\mathbf{VPW}^\pm}}{(\omega_R^2 - \omega_0^2) + \text{Im } \gamma \omega_R} \right] \quad (3.163a)$$

where the frequencies of $\mathbf{C} \rightleftharpoons \mathbf{W}$ pulsation of particles of Sender (ω_S) and Receiver (ω_R) are:

$$\omega_R = \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} = \mathbf{R} \omega_0^{in} + (\omega_B^{ext})_R \quad (3.164)$$

$$\omega_S = \omega_{\mathbf{C} \rightleftharpoons \mathbf{W}} = \mathbf{R} \omega_0^{in} + (\omega_B^{ext})_S \quad (3.164a)$$

γ is a damping coefficient due to *decoherence effects*, generated by local fluctuations of Bivacuum deteriorating the phase/spin transmission via $\mathbf{VirG}_{SME}$; $(\mathbf{m}_V^+)_{S,R}$ are the actual mass of (S) and (R); $[\mathbf{F}_{\mathbf{VPW}}]$ is a *tuning force of virtual pressure waves* $\mathbf{VPW}^\pm$ of Bivacuum with tuning energy $\mathbf{E}_{VPW} = \mathbf{q} \mathbf{m}_0 \mathbf{c}^2$ and wave length $\mathbf{L}_{VPW} = \hbar / \mathbf{m}_0 \mathbf{c}$

$$\mathbf{F}_{\mathbf{VPW}^\pm} = \frac{\mathbf{E}_{VPW}}{\mathbf{L}_{VPW}} = \frac{\mathbf{q}}{\hbar} \mathbf{m}_0^2 \mathbf{c}^3 \quad (3.165)$$

The most probable *tuning force* has a minimum energy, corresponding to $\mathbf{q} = \mathbf{j} - \mathbf{k} = \mathbf{1}$.

The influence of *virtual tuning force* ($\mathbf{F}_{\mathbf{VPW}^\pm}$) stimulates the synchronization of [S] and [R] pulsations, i.e. $\omega_R \rightarrow \omega_S \rightarrow \omega_0$. This fundamental frequency $\omega_0 = \mathbf{m}_0 \mathbf{c}^2 / \hbar$ is the same in any space volume, including those of Sender and Receiver.

The $\mathbf{VirG}_{SME}$ represent quasi **1D** macroscopic virtual Bose condensate with a configuration of single microtubules, formed by Bivacuum bosons ($\mathbf{BVB}^\pm$) or with configuration of double microtubules, composed from Cooper pairs as described in previous section.

The effectiveness of entanglement between two or more similar elementary particles is dependent on synchronization of their $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation frequency and 'tuning' the phase

of these pulsations via nonlocal virtual guide $(\mathbf{VirG}_{SME})^{S,R}$ between Sender and Receiver under the action of the virtual pressure waves $\mathbf{VPW}_{q=1}^{\pm}$ and Tuning energy of Bivacuum.

The mechanism proposed may explain the experimentally confirmed nonlocal interaction between coherent elementary particles [69], atoms and between remote coherent clusters of molecules.

Our theory predicts that the same mechanism, involving a nonlocal net of $\mathbf{VirG}_{SME}$, may provide the entanglement even between macroscopic systems, including biological ones. It is possible, if the frequency and phase of $[C \rightleftharpoons W]$ pulsations of clusters of their particles in state of mesoscopic Bose condensation are 'tuned' to each other [22].

J. Experimental data, confirming Unified theory (UT)

J.1 The electromagnetic radiation of nonuniformly accelerating charges

It follows from our theory, that the charged particle, *nonuniformly* accelerating in cyclotron, synchrotron or in undulator, can be a source of photons. The same is true for dipole radiation (section III.F.1).

From eqs.(3.74, 3.74a and 3.98) we get general expression for electromagnetic radiation, dependent on the doubled kinetic energy $\Delta(2\mathbf{T}_k) = \Delta(\mathbf{m}_V^+ \mathbf{v}^2)$ of alternately accelerated charged particle and related *inelastic* recoil effects, accompanied its $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation:

$$\hbar\omega_{EH} + \Delta 2\mathbf{T}_k = \Delta \{(\mathbf{1} + \mathbf{R})[\mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{m}_0 \mathbf{c}^2]\} = \Delta[(\mathbf{1} + \mathbf{R})\mathbf{TE}] = \quad (3.166)$$

$$\text{or : } \hbar\omega_{EH} + \Delta 2\mathbf{T}_k = \Delta \left[|\alpha \mathbf{m}_V^+ \mathbf{v}^2|_{rec} + \mathbf{m}_V^+ \mathbf{v}^2 \right]^{ext} \quad (3.166a)$$

$$\text{where } \mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \text{ is relativistic factor}$$

We can see from this formula, that the alternation of kinetic energy of charged particle, resulting from its alternating acceleration, can be accompanied by electromagnetic radiation. This effect occurs, if the jump of kinetic energy $\Delta 2\mathbf{T}_k$ and corresponding *inelastic* recoil energy jump: $\Delta |\alpha \mathbf{m}_V^+ \mathbf{v}^2|_{rec}$ exceed the energetic threshold necessary for photon origination. The $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations of all three sub-elementary fermions of triplets of charged elementary particles: $\langle [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-]_W + (\mathbf{F}_{\uparrow}^-)_C \rangle_{p,e} \rightleftharpoons \langle [\mathbf{F}_{\uparrow}^+ \bowtie \mathbf{F}_{\downarrow}^-]_C + (\mathbf{F}_{\uparrow}^-)_W \rangle_{p,e}$, modulated by external translational dynamics, participate in photon creation. In accordance with our model (Fig.3.2) photons are the result of fusion of the electron and positron like triplets. They can be a result of correlated asymmetric excitation of three pairs of Bivacuum Cooper

pairs $3[\mathbf{BVF}^\dagger \bowtie \mathbf{BVF}^\dagger]$ by accelerated electrons or positrons, like in the case of excited atoms and molecules.

It is the Tuning energy (**TE**) of Bivacuum: $\mathbf{TE} = [\mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{q} \mathbf{m}_0 \mathbf{c}^2]$, that induces the transitions of the excited state of particle to its ground state, corresponding to the rest mass energy:

$$\mathbf{m}_V^+ \mathbf{c}^2 \xrightarrow{TE} \mathbf{m}_0 \mathbf{c}^2 \quad \text{at } \mathbf{v}_{tr}^{ext} \rightarrow 0, \text{ if } \mathbf{q} = 1 \quad (3.167)$$

There are huge numbers of experimental data confirming this consequence of our theory for electromagnetic radiation. The gravitational radiation in form of Virtual Pressure Waves (VPW $^\pm$) in similar conditions is also predictable.

At the permanent (uniform) acceleration of the charged elementary particle, moving along the hyperbolic trajectory, the radiation is absent because it does not provide the fulfilment of condition of overcoming of activation barrier, necessary for three Cooper pairs symmetry shift. The photon radiation by charged particles is possible only on the conditions of nonuniform and big enough jumps of particles accelerations.

Some similarity is existing between the mechanisms of inelastic phonon excitation in solids, detected by γ – resonance spectroscopy, and photon excitation in Bivacuum by alternatively accelerated particle.

The one more consequence of Unified Theory is that the radiation of photons, induced by accelerations of charged elementary particle, should be strongly asymmetric and coincide with direction of charged particle propagation in space [21]. This is also well confirmed result by analysis of synchrotron and undulator radiation.

J.2 The double turn (720 $^\circ$), as a condition of the fermions spin state reversibility

It is known fact, that the total rotating cycle for spin of the electrons or positrons is not 360 $^\circ$, but 720 $^\circ$, i.e. *double turn* by external magnetic field of special configuration, is necessary to return elementary fermions to starting state [70]. The correctness of our model of elementary particles was testified by its ability to explain this nontrivial fact [21, 22] on example of the electron or proton:

$$< [F_\uparrow^+ \bowtie F_\downarrow^-]_{x,y} + (\mathbf{F}_\uparrow^-)_z >^{e,\tau} \quad (3.168)$$

The external rotating $\mathbf{H}$ field interact with sub-elementary fermions/antifermions of triplets in two stage manner ($2 \cdot 360^0$), changing their spins. The angle of spin rotation of sub-elementary particle and antiparticles of neutral pairs $[F_{\uparrow}^+ \bowtie F_{\downarrow}^-]$ are the additive parameters. It means that turn of resulting spin of *pair* on 360^0 includes reorientation spins of each $F_{\uparrow}^+$ and $F_{\downarrow}^-$ only on 180^0 . Consequently, the full spin turn of pair $[F_{\uparrow}^+ \bowtie F_{\downarrow}^-]$ resembles that of Möbius transformation.

The spin of unpaired sub-elementary fermion $\mathbf{F}_{\uparrow}^-$, in contrast to paired ones, makes a *full turn* each 360^0 , i.e. twice in 720^0 cycle:

$$\begin{aligned} & \langle [(F_{\uparrow}^+)_x \bowtie (F_{\downarrow}^-)_y] + (\mathbf{F}_{\uparrow}^-)_z \rangle \rightarrow \\ & \xrightarrow{360^0} \langle [(F_{\downarrow}^+)_x \overset{180^0+180^0}{\bowtie} (F_{\uparrow}^-)_y] + (\mathbf{F}_{\uparrow}^-)_z \rangle \rightarrow \\ & \xrightarrow{360^0} \langle [(F_{\uparrow}^+)_x \bowtie (F_{\downarrow}^-)_y] + (\mathbf{F}_{\uparrow}^-)_z \rangle \end{aligned} \quad (3.169)$$

The difference between the intermediate - 2nd stage and the original one in (3.169) is in opposite spin states of paired sub-elementary particle and antiparticle:

$$[(F_{\uparrow}^+)_x \bowtie (F_{\downarrow}^-)_y] \xrightarrow{360^0} [(F_{\downarrow}^+)_x \overset{180^0+180^0}{\bowtie} (F_{\uparrow}^-)_y] \quad (3.170)$$

It follows from our consideration, that the 3D spatial organization of the electron (positron) is asymmetric, and some difference $(F_{\uparrow}^+)_x \neq [(F_{\downarrow}^+)_x \text{ and } (F_{\downarrow}^-)_y \neq (F_{\uparrow}^-)_y]$ is existing. The $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of unpaired sub-elementary fermion $(\mathbf{F}_{\uparrow}^-)_z$ is counterphase and spatially compatible in basic state of the electron triplet with $(F_{\downarrow}^-)_y$ and in the intermediate state - with its partner $(F_{\downarrow}^+)_x$ in triplets (3.169).

One more known "strange" experimental result can be explained by our dynamic model of triplets of elementary particles. The existence of two paired in-phase pulsating sub-elementary fermions (3.170) with opposite parameters, exchanging by spin, charge and energy in the process of their $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation, can be responsible for the *two times stronger magnetic field*, generated by electron, as compared with those, generated by rotating sphere with charge $|e^-|$.

J.3 Michelson-Morley experiment, as a possible evidence for the Virtual Replica of the Earth existing

The experiments, performed in 1887 by Michelson-Morley and similar later experiments of higher precision, has been based on checking the difference of light velocity in the direction of Earth motion on its orbit around the Sun and in the direction normal or opposite to the this one. In the case of fixed aether (vacuum) with condensed matter properties, *independent of the Earth motion*, one may anticipate that the difference in these two light velocities should exist, if the value of refraction index of vacuum/aether is independent on velocity and direction of the Earth propagation in space. The absence of any difference was interpreted, as the absence of the Earth - independent fixed aether. This conclusion was used by Einstein in his postulate of permanency of light velocity, *but different time* in different inertial systems, termed 'the relativity principle'.

The conjecture of virtual replica (VR), following from our corpuscle-wave duality and Bivacuum models, allows the another interpretation of Michelson-Morley experiments. The VR of the Earth or any other material object represents a standing Bivacuum virtual pressure waves (VPW^+ and VPW^-), modulated by the object's particles corpuscle - wave pulsation (see section 8).

The distant component of VR may have at least as big diameter, as the Earth atmosphere and it moves in space together with planet. It is obvious, that in such 'virtual shell' of the Earth the light velocity could be the same in any directions.

I propose the experiment, which may confirm the existence of both: the VR and the Aether/Bivacuum, as a superfluid medium with certain mechanical properties, like compressibility providing the $VPW^\pm$ existing. For this end we assume that the properties of VR on distance of about few hundred kilometers from the planet surface differs from that on the surface.

If we perform one series of the Michelson-Morley like experiments on the satellite, rotating with the same angle velocity as the Earth, i.e. fixed as respect to the Earth surface and another series of experiments on the surface, the difference in results will confirm our Virtual Replica theory and the Bivacuum model with Aether properties. The absence of difference in light velocity in opposite direction as respect to Earth trajectory can be explained in two ways:

1. As a result of equality of light velocity in any directions, independently on direction of Earth translational propagation in space (confirmation of the Einstein relativity principle and the absence of the Aether);

2. As a result of certain correlation between the translational velocity of the material object (Earth or satellite) and Bivacuum perturbation (Bivacuum dipoles symmetry shift) in the same direction, increasing the refraction index of Bivacuum and light velocity (see section 6.6). This explanation is compatible with the Aether concept.

Consequently, the absence of difference in light velocity in Michelson-Morley like experiments, in any case is not a strong evidence for the Aether absence.

*J.4. Interaction of particles with their Virtual Replicas, as a background
of two slit experiments explanation*

In accordance with our model, the electron and proton (Fig.3.1) are the triplets $\langle [\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}] + \mathbf{F}_{\downarrow}^{-} \rangle^{e,\tau}$, formed by two negatively charged sub-elementary fermions of opposite spins ($\mathbf{F}_{\uparrow}^{-}$ and $\mathbf{F}_{\downarrow}^{-}$) and one uncompensated sub-elementary antifermion ($\mathbf{F}_{\downarrow}^{+}$) of μ and τ generation. The symmetric pair of sub-elementary fermion and antifermion: $[\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]$ are pulsing between Corpuscular [C] and Wave [W] states in-phase, compensating the influence of energy, spin and charge of each other.

It follows from our model, that the charge, spin, energy and momentum of the electron and positron are determined just by uncompensated/unpaired sub-elementary fermion ($\mathbf{F}_{\uparrow}^{\pm}$). The parameters of ($\mathbf{F}_{\uparrow}^{\pm}$) are correlated strictly with similar parameters of pair $[\mathbf{F}_{\uparrow}^{-} \bowtie \mathbf{F}_{\downarrow}^{+}]$ due to conservation of symmetry of properties of sub-elementary fermions and antifermions in triplets. It means that energy/momentum and, consequently, de Broglie wave length and frequency of uncompensated sub-elementary fermion ($\mathbf{F}_{\uparrow}^{\pm}$) determine the empirical de Broglie wave properties of the whole particle (electron, positron).

The frequency of de Broglie wave and its length can be expressed from eq. 3.78 as:

$$\nu_B = \frac{(\mathbf{m}_V^+ \mathbf{v}^2)_{tr}^{ext}}{h} = \frac{\mathbf{v}}{\lambda_B} = \nu_{C=W} - \mathbf{R}\nu_0 \quad (3.171)$$

$$or : \nu_B = \frac{\mathbf{m}_V^+ \mathbf{c}^2}{h} - \mathbf{R}\nu_0 \quad (3.171a)$$

where: $\nu_0 = \mathbf{m}_0 \mathbf{c}^2 / h = \omega_0 / 2\pi$; $\lambda_B = h / \mathbf{m}_V^+ \mathbf{v}$

In a nonrelativistic case, when $\mathbf{v} \ll \mathbf{c}$ and the relativistic factor $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \simeq 1$, the energy of de Broglie wave is close to Tuning energy (**TE**) of Bivacuum (3.131):

$$\mathbf{E}_B = h\nu_B \simeq \mathbf{m}_V^+ \mathbf{c}^2 - \mathbf{m}_0 \mathbf{c}^2 = \mathbf{TE} \quad (3.172)$$

The fundamental phenomenon of de Broglie wave is a result of modulation of the carrying internal frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation ($\omega_{in} = \mathbf{R}\omega_0 = \mathbf{R}\mathbf{m}_0\mathbf{c}^2/\hbar$) by the angular frequency of the de Broglie wave: $\omega_B = \mathbf{m}_V^+ \mathbf{v}_{tr}^2/\hbar = 2\pi\mathbf{v}/\lambda_B$, equal to the frequency of beats between the actual and complementary torus and antitorus of the *anchor* Bivacuum fermion ($\mathbf{BVF}_{anc}^\dagger$) of unpaired $\mathbf{F}_\downarrow^\pm$. The Broglie wave length $\lambda_B = \hbar/(\mathbf{m}_V^+ \mathbf{v})$ and mass symmetry shift of $\mathbf{BVF}_{anc}^\dagger$ is determined by the external translational momentum of particle: $\vec{\mathbf{p}} = \mathbf{m}_V^+ \vec{\mathbf{v}}$. For nonrelativistic particles $\omega_B \ll \omega_0$. For relativistic case, when $\mathbf{v}$ is close to $\mathbf{c}$ and $\mathbf{R} \simeq 0$, the de Broglie wave frequency is close to resulting frequency of $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation: $\omega_B \simeq \omega_{\mathbf{C}=\mathbf{W}}$.

In accordance with our model of duality, the reversible $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations of $\mathbf{F}_\downarrow^\pm$ and $\mathbf{BVF}_{anc}^\dagger$ are accompanied by *outgoing and incoming* Cumulative Virtual Cloud ($\mathbf{CVC}^\pm$), composed of subquantum particles of opposite energy. On this point, our understanding of duality and wave properties of particle coincide with that of Bohm and Hiley [17].

Introduced in our theory notion of *Virtual replica (VR) or virtual hologram* of any material object in Bivacuum is a result of interference of basic Virtual Pressure Waves ($\mathbf{VPW}_{q=1}^\pm$) and Virtual Spin Waves ($\mathbf{VirSW}_{q=1}^{\pm 1/2}$) of Bivacuum (reference waves), with virtual "object waves" ($\mathbf{VPW}_m^\pm$) and ($\mathbf{VirSW}_m^{\pm 1/2}$), representing $\mathbf{CVC}^+$ and $\mathbf{CVC}^-$ of pair $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$, modulated by de Broglie waves of the whole particles (see section 3.8).

The feedback influence of Bivacuum *Virtual replica* of the triplet on its original and corresponding momentum exchange may induce the wave - like behavior of even a single separated elementary fermion, antifermion $\langle [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+] + \mathbf{F}_\downarrow^\pm \rangle^{e,p}$ (Fig.3.1) or boson, like the photon (Fig.3.2).

The reason for periodical character of the electron's trajectory in our model (self-interference) can also be a result of periodic momentum oscillation, produced by $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of the anchor $\mathbf{BVF}_{anc}$ of unpaired sub-elementary fermion ($\mathbf{F}_\downarrow^-$) in triplet. In the case of photon, the momentum oscillation is equal to its frequency ($\nu_p = \mathbf{c}/\lambda_p$), as far the relativistic factor $\mathbf{R} = \sqrt{1 - (\mathbf{v}/\mathbf{c})^2}$ in (3.171) is zero or very close to zero near strongly gravitating objects. It is provided by the in-phase $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of the anchor $\mathbf{BVF}_{anc}^\dagger$ of central pair of sub-elementary fermions with similar spin orientations (Fig.3.2). The momentums of two side pairs of sub-elementary fermions of photon with opposite spins compensate each other, because their $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations are counterphase.

The duality properties of elementary particles can be understood, as far they can be

simultaneously in the corpuscle $[\mathbf{C}]$ and wave $[\mathbf{W}]$ phase. Our theory is able to prove this condition in two different ways:

1. In each triplet of elementary particle $\langle [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+] + \mathbf{F}_\uparrow^\pm \rangle$ the $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsations of the pair $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$ is counterphase with unpaired sub-elementary fermion $\mathbf{F}_\uparrow^\pm$. It means, that when $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$ is in the $[\mathbf{C}]$ phase, the $\mathbf{F}_\uparrow^\pm$ is in the $[\mathbf{W}]$ phase and vice versa;

2. In the absence of particle accelerations, meaning its permanent kinetic energy, the internal time of $[\mathbf{C}]$ and $[\mathbf{W}]$ phases is the same. Consequently, in the internal reference system, both phase of unpaired or paired sub-elementary fermions in different de Broglie wave semiperiods can be considered as simultaneous.

This statement follows from our notion of time and its pace for closed system (see section 7.1).

The formula for *pace of time* for any closed coherent system or free particle is determined by the pace of its kinetic energy: $(\mathbf{m}_V^+ \mathbf{v}^2/2)_{x,y,z} = \mathbf{m}_0 \mathbf{v}^2 / \left(2\sqrt{1 - (\mathbf{v}/\mathbf{c})^2} \right)_{x,y,z}$ change, which is anisotropic in a general case:

$$\left[\frac{d\mathbf{t}}{\mathbf{t}} = -d\mathbf{T}_k / \mathbf{T}_k \right]_{x,y,z} \quad (3.173)$$

For the internal time itself it follows from (3.173):

$$\left[\mathbf{t} = -\frac{\mathbf{v}}{d\mathbf{v}/d\mathbf{t}} \frac{1 - (\mathbf{v}/\mathbf{c})^2}{2 - (\mathbf{v}/\mathbf{c})^2} \right]_{x,y,z} \quad (3.174)$$

We can see, that in the absence of acceleration ($d\mathbf{v}/d\mathbf{t} = \mathbf{0}$ and $d\mathbf{v} = \mathbf{0}$), i.e. permanent kinetic energy, momentum and, consequently, permanent de Broglie wave frequency and length, the time for such particle is infinitive and pace of time is zero:

$$\mathbf{t} \rightarrow \infty \quad \text{and} \quad \frac{d\mathbf{t}}{\mathbf{t}} \rightarrow 0 \quad (3.175)$$

These conditions mean that the $[\mathbf{C}]$ and $[\mathbf{W}]$ phases of triplets - elementary particles, moving with permanent velocity, may be considered, as simultaneous. The oscillation of kinetic energy and velocity of triplets occur only in transition states $[\mathbf{C} \rightarrow \mathbf{W}]$ or $[\mathbf{W} \rightarrow \mathbf{C}]$, accompanied by *recoil* $\rightleftharpoons$ *antirecoil* effects.

We can see from the above analysis, that our model does not need the Bohmian "quantum potential" [17] or de Broglie's "pilot wave" for explanation of wave-like behavior of elementary particles.

Scattering of the photon on a free electron will affect its velocity, momentum, mass, wave B frequency, length, its virtual replica (VR) and its feedback influence on the electron, following by change of the interference picture.

Our theory predicts that, applying of the EM field to *singe electrons* with frequency resonant to their de Broglie frequency, should be accompanied by alternative acceleration of the electrons, modulation of their internal time and Virtual Replica with the same frequency, accompanied by 'washing out' the interference pattern in two-slit experiment. This consequence of our explanation of two-slit experiment can be easily verified.

K. The main Conclusions of Unified Theory (section III)

1. A new Bivacuum model, as the infinite dynamic superfluid matrix of virtual dipoles, named Bivacuum fermions $(\mathbf{BVF}^\dagger)^i$ and Bivacuum bosons $(\mathbf{BVB}^\pm)^i$, formed by correlated torus ($\mathbf{V}^+$) and antitorus ($\mathbf{V}^-$), as a collective excitations of subquantum particles and antiparticles of opposite energy, charge and magnetic moments and separated by energy gap, is developed. In primordial non polarized Bivacuum, i.e. in the absence of matter and fields, these parameters of torus and antitorus totally compensate each other. Their spatial and energetic properties correspond to three generations: electrons, muons and tauons ($i = e, \mu, \tau$). The positive and negative Virtual Pressure Waves ($\mathbf{VPW}^\pm$) and Virtual Spin Waves ($\mathbf{VirSW}^{S=\pm 1/2}$) are the result of emission and absorption of positive and negative Virtual Clouds ($\mathbf{VC}^\pm$), resulting from transitions of $\mathbf{V}^+$ and $\mathbf{V}^-$ between different state of excitation.

2. It is demonstrated that symmetry shift between $\mathbf{V}^+$ and $\mathbf{V}^-$ parameters to the left or right, opposite for Bivacuum fermions $\mathbf{BVF}^\dagger$ and antifermions $\mathbf{BVF}^\downarrow$ with relativistic dependence on their external rotational-translational velocity, is accompanied by sub-elementary fermion and antifermion formation. The formation of sub-elementary fermions and their fusion to stable triplets of elementary fermions, like electrons and protons $\langle [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+] + \mathbf{F}_\uparrow^\pm \rangle^{e,p}$, corresponding to the rest mass and charge origination, become possible at the certain velocity of angular rotation of Cooper pairs of $[\mathbf{BVF}^\uparrow \bowtie \mathbf{BVF}^\downarrow]$ around a common axis. It is shown, that this rotational-translational velocity is determined by Golden Mean condition: $(\mathbf{v}/\mathbf{c})^2 = \phi = 0.618$. The photon is a result of fusion (annihilation) of two triplets of particle and antiparticle: [electron + positron] or [proton + antiproton].

The photon represents a rotating sextet of sub-elementary fermions and antifermions with axial structural symmetry and minimum energy $2\mathbf{m}_0^e c^2$.

3. The fundamental physical roots of Golden Mean condition: $(\mathbf{v}/c)^2 = \mathbf{v}_{gr}^{ext}/\mathbf{v}_{ph}^{ext} = \phi$ are revealed as the equality of internal and external group and phase velocities of torus and antitorus of sub-elementary fermions, correspondingly: $\mathbf{v}_{gr}^{in} = \mathbf{v}_{gr}^{ext}$, $\mathbf{v}_{ph}^{in} = \mathbf{v}_{ph}^{ext}$. These equalities are named 'Hidden Harmony Conditions'.

4. The new expressions for total, potential and kinetic energies of de Broglie waves of elementary particles were obtained. The former represents the extended basic Einstein and Dirac formula for free particle: $\mathbf{E}_{tot} = \mathbf{m}_V^+ c^2 = \sqrt{1 - (\mathbf{v}/c)^2} \mathbf{m}_0 c^2 + \mathbf{h}^2/\mathbf{m}_V^+ \lambda_B^2$. The new formulas take into account the contributions of the actual mass/energy of torus ($\mathbf{V}^+$) and those of complementary antitorus ($\mathbf{V}^-$), correspondingly, of asymmetric sub-elementary fermions to the total ones. The shift of symmetry between the mass and other parameters of torus and antitorus of sub-elementary fermions are dependent on their *internal* rotational-translational dynamics in triplets and the *external* translational velocity of the whole triplets.

5. A dynamic mechanism of [corpuscle ($\mathbf{C}$) $\rightleftharpoons$ wave ($\mathbf{W}$)] duality is proposed. It involves the modulation of the internal (hidden) quantum beats frequency between the asymmetric 'actual' (torus) and 'complementary' (antitorus) states of sub-elementary fermions or antifermions by the external - empirical de Broglie wave frequency of the whole particles.

6. It is demonstrated, that the elastic deformations and collective excitations of Bivacuum matrix, providing field origination, are a consequence of reversible [*recoil* $\rightleftharpoons$ *antirecoil*] effects, generated by correlated [*Corpuscle* $\rightleftharpoons$ *Wave*] pulsation of sub-elementary fermions/antifermions of triplets and their fast rotation. The linear and circular alignment of Bivacuum dipoles are responsible for electrostatic and magnetic field origination. The gravitational waves and field are the result of positive and negative virtual pressure waves excitation (VPW^+ and VPW^-) by the in-phase [$\mathbf{C} \rightleftharpoons \mathbf{W}$] pulsation of pairs $[\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+]$ of triplets $\langle [\mathbf{F}_\uparrow^- \bowtie \mathbf{F}_\downarrow^+] + \mathbf{F}_\uparrow^\pm \rangle$, counterphase to that of unpaired $\mathbf{F}_\uparrow^\pm$. Such virtual waves provide the attraction or repulsion between pulsing remote particles, depending on phase shift of pulsations, as in the case of hydrodynamic Bjerknes force action. The zero-point vibrations of particle and evaluated zero-point velocity of these vibrations are also the result of [*recoil* $\rightleftharpoons$ *antirecoil*] effects, accompanied by [$\mathbf{C} \rightleftharpoons \mathbf{W}$] pulsation of triplets.

7. Maxwell's *displacement current* and the *additional instant current* are the consequences of Bivacuum virtual dipoles ($\mathbf{BVF}^\uparrow$ and $\mathbf{BVB}^\pm$) excitations and vibrations, corre-

spondingly. The excitations and vibrations are accompanied by the elastic deformations of secondary Bivacuum matrix, induced by presence of matter and fields. The increasing of the excluded for photons volume of tori and antitori, enhance the refraction index of Bivacuum and decrease the light velocity near strongly gravitating and charged objects. The nonzero contribution of the rest mass energy to photons and neutrino energy is a consequence of the enhanced refraction index of secondary Bivacuum.

8. It is shown that the Principle of least action and realization of 2nd and 3d laws of thermodynamics for closed systems - can be a result of slowing down the dynamics of particles and their kinetic energy decreasing, under the influence of the basic - lower frequency Virtual Pressure Waves ($\mathbf{VPW}_{q=1}^{\pm}$) with minimum quantum number $q = j - k = 1$. This is a consequence of induced combinational resonance between $[\mathbf{C} \rightleftharpoons \mathbf{W}]$ pulsation of particles and basic $\mathbf{VPW}_{q=1}^{\pm}$ of Bivacuum. The new notion of Bivacuum Tuning Energy (TE), responsible for forcing of particles pulsation frequency to resonance conditions with $\mathbf{VPW}_{q=1}^{\pm}$, is introduced.

9. It is demonstrated, that the dimensionless 'pace of time' ($d\mathbf{t}/\mathbf{t} = -d\mathbf{T}_k/\mathbf{T}_k$) and time itself for each closed system are determined by the change of this system kinetic energy. They are positive, if the particles of the system are slowing down under the influence of $\mathbf{VPW}_{q=1}^{\pm}$ and Tuning energy of Bivacuum. The $(d\mathbf{t}/\mathbf{t})$ and $(\mathbf{t})$ are negative in the opposite case. This new concept of time does not contradict the relativistic theory.

10. The notion of Virtual Replica (VR) or virtual hologram of any material object is developed. The $\mathbf{VR}$ is a result of interference of all-pervading quantized Virtual Pressure Waves ($\mathbf{VPW}_q^+$ and $\mathbf{VPW}_q^-$) and Virtual Spin waves ($\mathbf{VirSW}_q^{\mathbf{S}=\pm 1/2}$) of Bivacuum, working as "reference waves" in holograms formation, with modulated by de Broglie waves of matter atoms and molecules - "object waves" $\mathbf{VPW}_m^{\pm}$ and $\mathbf{VirSW}_m^{\pm 1/2}$. In our Unified theory the virtual pressure waves are identified with gravitational waves. The mechanism of gravitation may have common features with the hydrodynamic Bjerknes force between pulsating spheres. Their attraction or repulsion is dependent on the phase shift between pulsating spheres.

11. A possible Mechanism of Quantum entanglement between remote elementary particles via Virtual Guides of spin, momentum and energy ($\mathbf{VirG}_{\mathbf{S},\mathbf{M},\mathbf{E}}$) is proposed. The $\mathbf{VirG}_{\mathbf{S},\mathbf{M},\mathbf{E}}$, connecting similar and coherent electrons and nuclears of atoms of Sender(S) and Receiver(R). The single $\mathbf{VirG}_{\mathbf{S},\mathbf{M},\mathbf{E}}^{\mathbf{BVB}^{\pm}}$ can be assembled from Bivacuum bosons ($\mathbf{BVB}^{\pm}$)ⁱ and the twin $\mathbf{VirG}_{\mathbf{S},\mathbf{M},\mathbf{E}}^{\mathbf{BVF}^{\uparrow} \bowtie \mathbf{BVF}^{\downarrow}}$ from adjacent microtubules, rotating in opposite directions,

are formed by Cooper pairs of Bivacuum fermions $[\mathbf{BVF}^\dagger \bowtie \mathbf{BVF}^\dagger]^i$. The spin/information transmission via Virtual Guides is accompanied by reorientation of spins of tori and anti-tori of conjugated Bivacuum dipoles. The momentum and energy transmission from S to R is realized by the instant pulsation of diameter of such virtual microtubule. The Virtual Guides of both kinds represent the quasi 1D virtual Bose condensate with nonlocal properties, similar to that of 'wormholes'.

12. It is demonstrated that the consequences of Unified theory (UT) of Bivacuum, matter and fields are in good accordance with known experimental data. Among them: electromagnetic radiation of nonuniformly accelerating charges and the double turn ($2 \times 360 = 720^\circ$) of the electron's spin in external magnetic field, necessary for total reversibility of spin state of fermion. The two slit experiment also get its explanation even in a single particles case, as a consequence of interference of particle in the wave phase with its own virtual replica (VR).

IV. GENERAL CONCLUSIONS

There are several important physical features being common to the two present theories, thus reinforcing their results. Among the basic properties there are the similarities with the Dirac theory, the symmetry with respect to the concepts of matter and antimatter, the non-appearance of magnetic monopoles, and a space-charge vacuum current in addition to the displacement current. Further there are the models of the leptons having an internal vortex-like structure and force balance, the photon models with a small but nonzero rest mass and a slightly reduced velocity of propagation, and the simultaneous particle-wave behavior.

There are further results by which the two theories can be considered to become complementary to each other. Among the characteristic results of Theory(I), one can thus mention the deduced point-charge-like state of the electron model, being associated with a very small but nonzero effective radius. This removes the self-energy problem and it presents a more surveyable alternative to the renormalization procedure. Further results are the deduced electronic charge which deviates by only one percent from its experimental value, the photon wave-packet models having limited spatial extensions associated with a nonzero angular momentum, and the deduced needle-radiation under certain conditions. Theory(I) has so

far been limited to applications on leptons, individual photons, and certain types of light beams.

A number of characteristic results from Theory(II) are also of special interest. This Unified theory includes the following new issues:

- the new model of vacuum, named Bivacuum;
- the extension of special theory of relativity, unifying it with quantum mechanics;
- the dynamic mechanism of corpuscle-wave duality;
- the mechanism of the rest mass and charge origination, as a result of Bivacuum dipoles symmetry shift, determined by Golden mean conditions;
- the mechanism of elementary particles fusion from sub-elementary particles;
- the new approach to quantum gravity and to electric and magnetic phenomena, related to specific perturbation of Bivacuum by corpuscle-wave pulsation of particles;
- the new approach to time problem;
- the mechanism of nonlocality, as a result of Bose condensation, the virtual one in Bivacuum and real in superconductors and superfluids;
- the concept of virtual replica or virtual hologram of any material object, making possible the direct and feedback reaction of matter with Bivacuum. The resonance interaction of basic virtual pressure waves (VPW^+ and VPW^-) with pulsing particles, slowing down their dynamics, display itself as a Principle of least action, 2nd and 3d law of thermodynamics.

Acknowledgements

The authors are indebted to Miss Anna Forsell for valuable and skillful help with the manuscript of this paper.

REFERENCES

-
- [1] R.P. Feynman, "Lectures in Physics: Mainly Electromagnetism and Matter", Addison-Wesley, Reading, Massachusetts (1964), p.28–1.

- [2] J.D. Jackson, “Classical Electrodynamics”, John Wiley and Sons, Inc., New York-London-Sydney (1962), Ch.6, p.201 and Ch.17,p.589.
- [3] L.H. Ryder, “Quantum Field Theory”, Cambridge Univ.Press (1996), Ch.10, p.406, Ch.1, Ch.3.4; see also edition of 1987.
- [4] W. Heitler, “The Quantum Theory of Radiation”, Third Edition, Clarendon Press, Oxford (1954), p.401 and 409.
- [5] J.J. Thomson, *Nature* *137*, 232 (1936).
- [6] Y. Tsuchiya, E. Inuzuka, T. Kuroono, and M. Hosoda, *Adv.Electron Phys.* *64A* 21(1985).
- [7] B. Lehnert, *Spec.Sci.Technol.*, *9*177(1986); *17*, 259(1994); and *17*, 267(1994).
- [8] B. Lehnert, *Optik* *99*113(1995).
- [9] B. Lehnert and S. Roy, “Extended Electromagnetic Theory”, World Scientific Publ., Singapore (1998).
- [10] B. Lehnert, in “Modern Nonlinear Optics”, Part 2, Second Edition, “Advances in Chemical Physics”, Vol.*119*, Edited by M.W.Evans, I.Prigogine, and S.A.Rice, John Wiley & Sons, Inc., New York (2001), p.1.
- [11] B. Lehnert, *Physica Scripta* *66*, 105(2002); in press (2005).
- [12] B. Lehnert and J. Scheffel, *Physica Scripta* *65* 200(2002).
- [13] B. Lehnert, in “Dirac Equation, Neutrinos and Beyond”, Edited by V.V.Dvoeglazov, Ukrainian Journal: “Electromagnetic Phenomena”, T.3, No.1(9), 35(2003).
- [14] B. Lehnert, *Physica Scripta* *T113* 41(2004).
- [15] B. Lehnert, Report TRITA-ALF-2004-02, Fusion Plasma Physics, Royal Institute of Technology, Stockholm (2004).
- [16] P.A.M. Dirac, “The Principles of Quantum Mechanics”, Clarendon Press, Oxford (1947).
- [17] D. Bohm and B.J. Hiley , “ The Undivided Universe. An Ontological Interpretation of Quantum Theory”, Routledge, London–New York (1993).
- [18] J.A. Wheeler, “Superspace and Quantum Geometrodynamics”, In “Battle Recontres”, Edited by C.M. De Witt and J.A. Wheeler, Benjamin, New York (1968), p.242–307.
- [19] B.G. Sidharth, “The Universe of Fluctuations”, <http://xxx.lanl.gov/abs/quant-ph/980831>, (1998).
- [20] B.G. Sidharth, “The Universe of Chaos and Quanta”, <http://xxx.lanl.gov/990228>, (1999).
- [21] A. Kaivarainen, “Bivacuum as a Matrix for Matter, Fields and Time Origination. Vir-

- tual Pressure Waves, Virtual Replicas and Overunity Devices”, Archives of Los Alamos, <http://arXiv.org/abs/physics/0207027>, (2004).
- [22] A. Kaivarainen, “New Approach to Entanglement and Quantum Psi Phenomena, Based on Unified Theory of Bivacuum, Matter, Fields and Time”, Archives of Los Alamos, <http://arXiv.org/abs/physics/0103031>, (2005).
 - [23] A. Kaivarainen, “Unified Theory of Bivacuum, Particles Duality, Fields and Time. Virtual Replicas of Material Objects and Possible Mechanisms of Nonlocality”, Archives of Los Alamos, <http://arXiv.org/abs/physics/003001>, (2005).
 - [24] H. Aspden, “The Physics of Creation”, Sabberton Publications, Southampton, England (2003).
 - [25] A.O. Barut and A.J. Bracken, *Phys.Rev. D*, *23*, 10(1981).
 - [26] A. Einstein, in “The Born-Einstein Letters”, edited by I.Born, Macmillan, London (1971), p.158.
 - [27] A. Einstein, “Ideas and Opinions”, Crown Publishers, Inc., New York (1982).
 - [28] J. Stachler, Editor, “The Collected Papers of Albert Einstein”, Volume 2, “The Swiss Years: Writings, 1900–1909”, Princeton University Press, Princeton, New Jersey (1989).
 - [29] E. Schrödinger, *Proc. Am. Phil. Soc.* *124*, 323 (1980).
 - [30] D. Hestness, *Found. of Phys.* *20*, 10(1990).
 - [31] H.E. Puthoff, “Polarizable-vacuum(PV) Representation of General Relativity”, Archives of Los Alamos, <http://xxx.lanl.gov/abs/gr-qc/990937>, (1991).
 - [32] V. Fock, “The theory of space, time and gravitation” Pergamon Press, London (1964).
 - [33] S.L. Glashov, *Nucl.Phys.*, *B22*, 579(1961).
 - [34] S. Weinberg, *Phys.Rev.Lett.*, *19*, 1264(1967).
 - [35] A. Salam, in “Elementary Particle Theory: Relativistic Groups and Analyticity”, Nobel Symposium No.8, Ed. by N.Svartholm, Almquist and Wiksell, Stockholm (1968), p.367.
 - [36] A. Rueda, and B. Haish, “A Vacuum-generated Inertia Reaction Force”, in “Computing Anticipatory Systems”, CASY’s 2000, 4th International Conference, Ed. by D.M.Dubois, AIP, Conference Proceedings, *573*, 89(2001).
 - [37] A. Kaivarainen, “Unified Model of Matter - Fields duality and Bivacuum Mediated Electromagnetic and Gravitational interactions”, In “Frontiers in Quantum Physics Research”, Edited by F. Columbus and V. Krasnoholvets, Nova Science Publ., Inc., (2004) p.83.

- [38] P.A.M. Dirac, “Directions in Physics”, Wiley-Interscience, New York (1978).
- [39] G. Feldman and P.T. Mathews, Phys.Rev. *130*, 1633(1963).
- [40] P.A.M. Dirac, Proc.Roy.Soc. *117*, 610(1928) and *118*, 351(1928).
- [41] P.M. Morse and H. Feshbach, “Methods of Theoretical Physics”, McGraw-Hill Book Comp., Inc., New York (1953), Part I, p.260 and 331; Part II, p.1320.
- [42] R.B. Leighton, “Principles of Modern Physics”, McGraw-Hill Book Comp., Inc., New York-Toronto-London (1959), Ch.20.
- [43] A. Gersten, Preprints CERN-TH.4687/87 and CERN-TH.4688, Geneva: CERN (1987).
- [44] A. Gersten, Found.Phys.Lett., *12*, 291(1999).
- [45] J.A. Stratton, “Electromagnetic Theory”, First Edition, McGraw-Hill Book Comp., Inc., New York and London (1941), Chs. II and VI.
- [46] J. Schwinger, Phys.Rev. *76*, 790(1949).
- [47] R.P. Feynman, “QED: The Strange Theory of Light and Matter”, Penguin, London (1990).
- [48] E.R. Cohen and B.N. Taylor, Physics Today, BG 7, August (1998).
- [49] G. Hunter and R.L.P. Wadlinger, Physics Essays *2*, 154(1989).
- [50] M.W. Evans and J.-P. Vigiér, “The Enigmatic Photon”, Kluwer Academic Publishers, Dordrecht, Vols. 1 (1994) and 2 (1995).
- [51] M.W. Evans and J.-P. Vigiér, “The Enigmatic Photon”, Kluwer Academic Publishers, Dordrecht, Vol. 3 (1996).
- [52] A. Einstein, Ann.Phys.(Leipzig), *7*, 132(1905) and *18*, 121(1917).
- [53] L. Bass and E. Schrödinger, Proc.Roy.Soc.A, *232*, 1 (1955).
- [54] L. de Broglie and J.-P. Vigiér, Phys.Rev.Lett., *28*, 1001(1972).
- [55] A. Einstein, “On the Quantum Theory of Radiation” (1917) in “Sources of Quantum Mechanics”, Ed. by B.L. van der Waerden, Amsterdam (1967).
- [56] H. Tauber, Physics Essays, *16*, 200(2003).

REFERENCES to Section III

57. Kaivarainen A. (1995). Hierarchic Concept of Matter and Field. Water, biosystems and elementary particles. New York, NY, pp. 485, ISBN 0-9642557-0-7.

58. Kiehn R.M. (1998). The Falaco Soliton: Cosmic strings in a swimming pool; Coherent structures in fluids are deformable topological torsion defects. At: IUTAM-SMFLO conf. at DTU, Denmark, May 25, 1997; URL: <http://www.uh.edu/~rkiehn>
59. Winterberg F. (2003). Plank mass plasma vacuum conjecture. *Z. Naturforsch*, 58a, 231-267.
60. Jin D.Z., Dubin D.H. E. (2000). Characteristics of two-dimensional turbulence that self-organizes into vortex crystals. *Phys. Rev. Lett.*, 84(7), 1443-1447.
61. Okun' L.B. (1998). *Physics of Elementary particles*. "Nauka", Moscow, pp. 272.
62. Berstetski V., Lifshhitz M., Pitaevski L. (1989). *Course of theoretical physics, Physics, vol. IV: Quantum electrodynamics*. Nauka, Moscow.
63. Kaivarainen A. (2001). New Hierarchic theory of condensed matter and its computerized application to water and ice. In the Archives of Los-Alamos: <http://arXiv.org/abs/physics/0102086>.
64. Kaivarainen A. (2001a). Hierarchic theory of matter, general for liquids and solids: ice, water and phase transitions. *American Institute of Physics (AIP) Conference Proceedings* (ed. D.Dubois), vol. 573, 181-200.
65. Dicke R. H. (1957). "Gravitation without a principle of equivalence," *Rev. Mod. Phys.* 29, 363-376.
66. Glansdorf P., Prigogine I. (1971). *Thermodynamic theory of structure, stability and fluctuations*. Wiley and Sons, N.Y.
67. Naudin J-L. website "The Quest For Overunity": <http://members.aol.com/jnaudin509/>
- Okun' L.B. (1998). *Physics of Elementary particles*. "Nauka", Moscow, pp. 272.
68. Prochorov A.M. *Physics. Big Encyclopedic Dictionary*. Moscow, 1999.
69. Aspect A. and Grangier P. (1983). Experiments on Einstein-Podolsky-Rosen type correlations with pairs of visible photons. In: *Quantum Theory and Measurement*. Eds. Wheeler J.A., Zurek W.H. Princeton University Press.
70. Davies P. (1985). *The search for a grand unified theory of nature*. A

Touchstone book, published by Simon & Schuster, Inc. NY.